



III P

0-

0

Caution Lang Part Caution





How To Order

What information do we need?

Please quote your:

- 1 Name
- 2 Address
- 3 Telephone Number
- 4 Fax Number
- 5 Email Address
- 6 Purchase Order Number
- 7 Product Description
- 8 Catalogue Number
- 9 Quantity Required
- **10** Any Specific Delivery Instructions

Where should you send your quote?

Please send your quote to us using one of the following options:



Through Our Website fortressdiagnostics.com/contact-us



Over the Phone +44 (0)28 9448 7676



By Email info@fortressdiagnostics.com

Advancing Global Diagnostics

Fortress Diagnostics is a well-established, multi award-winning IVD manufacturer based in the United Kingdom, supplying an extensive portfolio of products to over 100 countries worldwide.

We have been providing our distributors and end users with high quality products for over 20 years. The objective of Fortress Diagnostics is to provide the highest level of diagnostic solutions, globally, through continued investment in research and development.

Why choose Fortress Diagnostics?



Our proven product performance, matched with our selection of reputable distribution partners, has been instrumental in our continued success and growth. Built on a culture of quality and performance, we are committed to providing our customers with a reliable service and results they can trust.



Our ISO 13485:2016 certification for the design, development, manufacturing and distribution of in vitro diagnostics (IVD) medical reagents and instrumentation endorses our product range and ensures that the highest quality standards are constantly maintained.



Through our global distribution network, we provide highly accurate medical testing solutions to immunology, haematology and serological laboratories in hospitals, medical centres, clinics, blood banks and research institutions in over 100 international markets.



Hosting an in-house Research and Development team, we are committed to creating the next innovative IVD solutions to exceed our customer's requirments and advance the healthcare industry. We reinvest over 30% of our turnover in R&D every year.

Our Product Range

Fortress Diagnostics supplies an extensive portfolio of IVD products spanning the whole diagnostics field:

Clinical Chemistry Analyser Specific Reagents Quality Controls & Calibrators Immunoassays (ELISA) Immunoassays (CLIA) Syphilis Febrile Antigens Latex Serology Blood Grouping Rapid Tests Haematology Reagents Haemostasis Urine Tests Food & Feed Safety Instrumentation EQA & IQC Programmes Neonatal Screening Molecular Diagnostics Western Blot Assays Stains Antibiotic Discs



Find Out More





Website

fortressdiagnostics.com

The Fortress Diagnostics website details our full product range, allowing users to add products of interest to a "Quote Request" which is sent directly to our sales team. We regularly post blogs, event information, resources and company updates.

Quality Control Inserts Portal

insert.fortressdiagnostics.com

Fortress Diagnostics are excited to launch the new Fortress QC Inserts Portal. The portal allows customers to easily search our database of quality control inserts by product name, catalogue number or batch number in our easy-to-use filter tool. Inserts can be viewed online or downloaded in PDF format.





Brochures & Catalogues

fortressdiagnostics.com/resources

All Fortress Diagnostics brochures and catalogues can be viewed and downloaded on the Fortress Resource Hub in PDF format.

Videos

youtube.com/@fortressdiagnostics

Subscribe to our YouTube channel to explore information videos, tutorial guides for Seraqual 365, interviews for World Awareness Days and more.





Social Media

Explore and follow our social media pages across **Instagram**, **LinkedIn**, **Facebook** and **Twitter**. We regularly post updates on product launches, events Fortress are exhibiting at, news from the world of diagnostics and company developments.

SECTION 1	6	Clinical Chemistry & Analyser Specific Reagents	07
SECTION 2	Ī	Quality Controls & Calibrators	33
SECTION 3	H	Immunoassays (ELISA)	49
SECTION 4	H	Immunoassays (CLIA)	55
SECTION 5	R	Syphilis	59
SECTION 6	1	Febrile Antigens	63
SECTION 7	\odot	Latex Serology	67
SECTION 8	Q	Blood Grouping	70
SECTION 9	A	Rapid Tests	77
SECTION 10		Haematology Reagents	81
SECTION 11	$\boldsymbol{\otimes}$	Haemostasis	83
SECTION 12		Urinalysis	87
SECTION 13	×	Food & Feed Safety	89
SECTION 14	Ø	Instrumentation	95
SECTION 15	I	IQC & EQA Programmes	105
SECTION 16	-	Neonatal Screening	111
SECTION 17	~	Molecular Diagnostics	113
SECTION 18		Western Blot Assays	115
SECTION 19		Stains	117
SECTION 20		Antibiotic Discs	121

CLINICAL CHEMISTRY REAGENTS & ANALYSER SPECIFIC REAGENTS

SECTION 1

BII II

Clinical Chemistry

Clinical chemistry, also known as chemical pathology or clinical biochemistry, is a branch of laboratory medicine that focuses on the analysis of bodily fluids, primarily blood and urine, to diagnose and monitor disease. Clinical chemistry reagents are vital for accurate and reliable diagnostic testing, as they are the substances used to detect and measure specific analytes in biological samples.

High-quality reagents ensure the precision of tests, influencing the results that guide clinical decision-making. They enable the identification of various health conditions by reacting with substances in blood and urine, allowing for the assessment of organ function, metabolic processes, and disease states. Furthermore, the development and standardisation of reagents support advancements in technology and research, leading to improved diagnostic methods.

Fortress Diagnostics manufacturers, develops and supplies an extensive range of high quality clinical chemistry reagents. Our reagents are manufactured to the highest quality standards in an ISO 13485:2016–certified laboratory, ensuring excellent stability, accuracy and results you can trust. We also supply a large portfolio of analyser-specific reagents.

Ordering Information

Description	Method	Format	Cat No.	Size	Storage
Acid Phosphatase	Fast Red	Lyo	BXC0401A	R1: 1 x 65ml R2: 6 x 10ml R3:3 x 10ml R4: 1 x 5ml	2-8°C
Acid Phosphatase	Fast Red	Lyo	BXC0401B	R1: 1 x 105ml R2: 10 x 10ml R3:5 x 10ml R4: 1 x 5ml	2-8°C
Acid Phosphatase	Fast Red	Lyo	BXC0401C	R1: 1 x 35ml R2: 10 x 3ml R3:5 x 3ml R4: 1 x 5ml	2-8°C
ADA	Enzymatic	LS	BXC0208A	R1: 1 x 20ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Albumin	BCG (Concentrate)	LS	BXC0221A	R1: 6 x 60ml R4: 1 x 5ml	2-8°C
Albumin	BCG (Ready to Use)	Mono LS	BXC0222A	R1: 2 x 60ml R4: 1 x 5ml	Room Temp
Albumin	BCG (Ready to Use)	Mono LS	BXC0222B	R1: 6 x 60ml R4: 1 x 5ml	Room Temp
Aldolase	GDH/TIM (UV)	Lyo	BXC0391A	R1: 5 x 20ml R2: 2 x 1ml R3:1 x 0.5ml	2-8°C
Alkaline Phosphatase	Colorimetric	LS	BXC0183A	R1: 1 x 50ml R2: 2 x 125ml R4: 1 x 5ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	BXC0184A	R1: 5 x 25ml R2: 1 x 25ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	BXC0184B	R1: 5 x 50ml R2: 1 x 50ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	BXC0184C	R1: 5 x 100ml R2: 1 x 100ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	BXC0185A	R1: 5 x 25ml R2: 1 x 25ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	BXC0185B	R1: 5 x 50ml R2: 1 x 50ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	BXC0185C	R1: 5 x 100ml R2: 1 x 100ml	2-8°C
Alkaline Phosphatase	DEA (DGKC) 4+1	LS	BXC0187A	R1: 1 x 100ml R2: 1 x 25ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	Lyo	BXC0181A	R1: 5 x 20ml R2: 5 x 20ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	Lyo	BXC0181C	R1: 1 x 105ml R2: 10 x 10ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	Lyo	BXC0181D	R1: 1 x 65ml R2: 20 x 3ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	Lyo	BXC0182A	R1: 1 x 105ml R2: 5 x 20ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	Lyo	BXC0182D	R1: 1 x 105ml R2: 10 x 10ml	2-8°C
Alpha-1-Acid Glycoprotein		LS	BXC0890A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	Stool	BXC0189A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0190A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0190B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C

Clinical Chemistry | SECTION 1

Simple Hereinalden Simple	Description	Method	Format	Cat No.	Size	Storage
Update frame or gample Pre-dilation)DisDisc declassesNo. 12.2.0000 Ref. 1.2000CPC or gampleLT (GPT)color/metric (Slandard)LSBXC0212AStrin X: 10000 R2:1.X2.8°CLT (GPT)IFCCLSBXC0213AStrin X: 10000 R2:1.X2.8°CLT (GPT)IFCCLSBXC0214BTit X: 10000 R2:1.X2.8°CLT (GPT)Color/metric (Tablea)LSBXC0214BTit X: 10000 R2:1.X2.8°CLT (GPT)Color/metric (Tablea)LSBXC0214BTit X: 10000 R2:1.X2.8°CLT (GPT)DGKC 4+1LSBXC0214BTit X: 10000 R2:1.X2.8°CLT (GPT)DGKC 4+1LSBXC0214BTit X: 10000 R2:1.X2.8°CLT (GPT)IFCCLSOBXC0211BTit X: 10000 R2:1.X2.8°CLT (GPT)IFCCLSOBXC0215BTit X: 1	Alpha-1-Microglobulin	Sample Pre-dilution)	LS	BXC0888A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
LIT (GPT)Colorimetric (Standard)LSBK201216StandBr8-1 Karlinger24°CLIT (GPT)IFCCSKSKC0114Stat 100m R2.1 × Stat 00m R2.1 × Arr (GPT)IFCCSKC0114SKC0114Stat 100m R2.1 × Stat 00m R2.1 × Arr (GPT)SKC0114SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114SKC0114SKC0114Stat 100m R2.1 × Arr (GPT)SKC0114SKC	Alpha-1-Microglobulin		LS	BXC0888B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lit (PT)PCC<	ALT (GPT)	Colorimetric (Standard)	LS	BXC0212A	125ml R4: 1 x 5ml R6:1	2-8°C
Lit (orf)inc.Ls900/21920ml24° CLT (GPT)iFCCi.s8XC0213FR15.5 X100m1 R2.1 x24°C.1LT (GPT)Calorimetri (Table)i.s8XC0213FR1.5 X100m1 R2.1 x24°C.1LT (GPT)DCK C 4-1i.s8XC0213FR1.5 X100m1 R2.1 x24°C.1LT (GPT)DCK C 4-1i.s8XC0213FR1.5 X100m1 R2.2 x24°C.1LT (GPT)IFCCi.so8XC0211FR1.5 X100m1 R2.1 x24°C.1LT (GPT)IFCCi.so8XC0211FR1.5 X100m1 R2.2 x24°C.1LT (GPT)IFCCi.so8XC0211FR1.5 X50m1 R2.1 x24°C.1IT (GPT)IFCCi.so8XC0211FR1.5 X50m1 R2.1 x24°C.1IT (GPT)IFCCi.so8XC0211FR1.5 X50m1 R2.1 x24°C.1IT (GPT)IFCCi.so8XC0210FR1.5 X50m1 R2.1 x24°C.1IT (GPT)IFCCI.so8XC0210FR1.1 X20m1 R2.1 x24°C.1IT (GPT)IFCCI.so8XC0210FR1.1 X20m1 R2.1 x24°C.1IT (GPT)IFCCI.so8XC0520FR1.1 X20m1 R2.1 x24°C.1IT (GPT)IFCCI.so <td>ALT (GPT)</td> <td>IFCC</td> <td>LS</td> <td>BXC0213A</td> <td></td> <td>2-8°C</td>	ALT (GPT)	IFCC	LS	BXC0213A		2-8°C
Lit L(PT)IPCCLSBXC02114Thomal PCCall orimetric (Table)LSBXC02114R11:11 X100H1R2:11, X10FHCall CLT (GPT)DGKC 4+1LSBXC02158R1:12 X10HR1R2:11, X10FHCall CLT (GPT)DGKC 4+1LSBXC02158R1:12 X10FHCall CLT (GPT)DGKC 4+1LSBXC02116R1:12 X10FHCall CLT (GPT)IFCCLyoBXC02110R1:5 X0FHR2:20 X201LT (GPT)IFCCLyoBXC0211CR1:5 X0FHCall CLT (GPT)IFCCLyoBXC0211CR1:5 X0FHCall CumplaseDES/NPC7 IFCCLSBXC05630R1:12 X20HR24°CumplaseDES/NPC7 IFCCLSBXC05610R1:12 X0HR24°CumplaseDES/NPC7 IFCCLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620R1:12 X0HR24°CumplaseCNPG3MonLSBXC05620 <td< td=""><td>ALT (GPT)</td><td>IFCC</td><td>LS</td><td>BXC0213D</td><td></td><td>2-8°C</td></td<>	ALT (GPT)	IFCC	LS	BXC0213D		2-8°C
Lit (UP1) Coormetric (able) LS BX.021/N NO.001 P32 LTT (GPT) DGKC 4-11 LS BXC0218 Stril X 100m R22 X 2 Stri	ALT (GPT)	IFCC	LS	BXC0213F		2-8°C
Lit (GPT) DKK. 4*1 LS BXU215% 25m1 25m1 25m1 LT (GPT) DGKC 4+1 LS BXC02158 Stril 2 × 105m1 R2 2 × 30 28°C LT (GPT) IFCC Lyo BXC02110 R1 2 × 105m1 R2 1 × 3 28°C LT (GPT) IFCC Lyo BXC02110 R1 × 210m1 R2 2 × 50m 28°C LT (GPT) IFCC Lyo BXC02100 R1 × 105m1 R2 1 × 5 28°C INT (GPT) IFCC Lyo BXC02100 R1 × 105m1 R2 1 × 5 28°C INT (GPT) IFCC Lyo BXC05300 R1 1 × 25m1 R2 1 × 5m1 28°C INT MASIN EFC LYO BXC05400 R1 1 × 25m1 R2 1 × 5m1 28°C INT MASIN EFC LYO LYO BXC05630 R1 1 × 25m1 R2 1 × 5m1 28°C INT MASIN EFS-PNPG7 IFCC LS BXC0520 R1 + X 10m1 R2 1 × 5m1 28°C INT MASIN CAPC3 Mono LS BXC0520 R1 + X 10m1 R2 + X m1 28°C INT MASIN CAPC3	ALT (GPT)	Colorimetric (Table)	LS	BXC0214A		2-8°C
Lit (GPT)DURC #F1LSRAC 02167SPAID2.8 CLT (GPT)IFCCLyoRXC02116R1.2 X 105ml R2: 10 X 10ml 100ml R2: 10 X 10ml R2: 10 X<	ALT (GPT)	DGKC 4+1	LS	BXC0215A		2-8°C
LLI (GPT)IFCCUgoBX.02118Zom24°CLLT (GPT)IFCCUgoBX.02110R1:1 × 105ml R2:1 × x52-8°CLLT (GPT)IFCCUgoBX.02110R1:1 × 55ml R2: × x50ml2-8°CLLT (GPT)IFCCUgoBX.02110R1:1 × 55ml R2: × x50ml2-8°CummoniaGLDHMone L5BX.0376AR1:1 × 100ml R2: 1 × x50ml2-8°CummylaseEPS-PNFG7 IFCCI.SuBX.00563BR1:1 × 25ml R2: x 5ml2-8°CumylaseEPS-PNFG7 IFCCI.SuBX.05653BR1:1 × 25ml R2: 1 × 5ml2-8°CumylaseBenzylidene Blocked-PNPG7LyoBX.05654BR1:1 × 100ml R2: 1 × 2-8°C2-8°CumylaseCNPG3Mone L5BX.05654BR1:1 × 100ml R2: 1 × 2-8°C2-8°CumylaseBenzylidene Blocked-PNPG7LyoBX.05654BR1:1 × 100ml R2: 1 × 2-8°C2-8°CumylaseCNPG3Mone L5BX.0567AR1:1 × 30ml R2: 1 × 2-8°C2-8°CumylaseEPS-PNFG7 IFCCLSBX.0567AR1:1 × 30ml R2: 1 × 2-8°C2-8°CumylasePancreaticEPS-PNFG7 IFCCLSBX.0567AR1:1 × 30ml R2: 1 × 2-8°C	ALT (GPT)	DGKC 4+1	LS	BXC0215B		2-8°C
Lit (GPT)FCC(yoBACU TO(mon)2.8 °CLit (GPT)IFCCLyoBX C02110R1: 5 X SOM R2: 5 X SOM2.8°CLit (GPT)IFCCLyoBX C02120R1: 14 X SAM R2: 20 × 2012.8°CumpolaeCDBHMono LSBX C02760R1: 14 X SAM R2: 20 × 2012.8°CumplaeEPS pNPG7 IFCCLSBX C05630R1: 12 X SIM R2: 1 X SI2.8°CumplaeEPS-pNPG7 IFCCLSBX C05630R1: 12 X SIM R2: 1 X SI2.8°CumplaeCNPG3Mono LSBX C05630R1: 1 X 30m1 R2: 1 X SI2.8°CumplaeCNPG3Mono LSBX C05640R1: 10 × 10m1 R2: 1 X SI2.8°CumplaeCNPG3Mono LSBX C05640R1: 10 × 10m1 R2: 1 X SI2.8°CumplaeCNPG3Mono LSBX C05640R1: 10 × 10m1 R2: 1 X SI2.8°CumplaeCNPG3Mono LSBX C05640R1: 5 × 10m1 R2: 1 X SI2.8°CumplaeCNPG3Kan CMono LSBX C03700R1: 5 × 10m1 R2: 1 X SI2.8°CumplaeConverting Enzyme (ACE)Mono LSBX C0370R1: 5 × 10m1 R2: 1 X SI2.8°CumplaeTurbilatexLSBX C0370R1: 1 X 30m1 R2: 1 X SI2.8°Cumplae </td <td>ALT (GPT)</td> <td>IFCC</td> <td>Lyo</td> <td>BXC0211B</td> <td></td> <td>2-8°C</td>	ALT (GPT)	IFCC	Lyo	BXC0211B		2-8°C
LT (0PT)IFCCLyoBXC0211ER1:1 × 45m1R2: 20 × 2012.8°CAnnoniaGLDHMono L5BXC0376AR1:1 (0 × 10m1 R4: 1.1 × 25m12.8°CAmylaseEP5-NPC7/FCCL5BXC0563AR1:1 × 25m1 R2: 1 × 5m12.8°CAmylaseEP5-NPC7/FCCL5BXC0563AR1:1 × 25m1 R2: 2 × 5m12.8°CAmylaseEP5-NPC7/FCCL5BXC0563AR1:1 × 10m1 R2: 1 × 30m12.8°CAmylaseBenzylidene Blacked-pNPC7LyoBXC0561AR1:1 × 10m1 R2: 1 × 30m12.8°CAmylaseCNPC3Mono L5BXC0562AR1: 6 × 10m12.8°CAmylase, PancreaticCNPG3Mono L5BXC0564AR1: 5 × 10m1 R2: 5 × 5m2.8°CAmylase, PancreaticCNPG3Mono L5BXC057AR1: 5 × 10m1 R2: 5 × 5m2.8°CAmylase, PancreaticL5BXC057AR1: 5 × 10m1 R2: 1 × 102.8°CAmilaeImmunoturbidimetric (Without)L5BXC0379CR1: 1 × 30m1 R2: 1 × 30m1	ALT (GPT)	IFCC	Lyo	BXC0211C		2-8°C
Mono LS BXC0376A R1:10 × 10ml R4:1 x 1ml Controls: 2 x 1ml 2.4°C Impliese EPS-pNPC7 IFCC LS BXC0563B R1:1 x 25ml R2:1 x 5ml 2.4°C Impliese EPS-pNPC7 IFCC LS BXC0563B R1:1 x 25ml R2:1 x 5ml 2.4°C Impliese EPS-pNPC7 IFCC LS BXC0563B R1:1 x 100ml R2:1 x 2.4°C Impliese EPS-pNPC7 IFCC LS BXC0563B R1:1 x 100ml R2:1 x 2.4°C Impliese EPS-pNPC7 IFCC LS BXC0563B R1:1 x 100ml R2:1 x 2.4°C Impliese ENSPNG7 IFCC LS BXC0562B R1:1 x 100ml R2:1 x 2.4°C Impliese CNPC3 Mono LS BXC0562B R1:1 x 10ml R2:1 x 2.4°C Impliese CNPC3 Mono LS BXC042PA R1:5 x 10ml R2:1 x 2.4°C Impliese Immunoturbidimetric (Without) LS BXC037PA R1:1 x 30ml R2:1 x 2.4°C Impliese Turbilatex LS BXC037PB R1:1 x 30ml R2:1 x 2.4°C Impliese Turbilatex </td <td>ALT (GPT)</td> <td>IFCC</td> <td>Lyo</td> <td>BXC0211D</td> <td>R1: 5 x 50ml R2: 5 x 50ml</td> <td>2-8°C</td>	ALT (GPT)	IFCC	Lyo	BXC0211D	R1: 5 x 50ml R2: 5 x 50ml	2-8°C
mmmonia CLDH Mono LS BCUS140 Tim L Controls: 2 x 1ml 24°C smylase EPS-pNPG7 IFCC LS BC05630 R1:1 x 25ml R2: 2 x 5ml 2.4°C smylase EPS-pNPG7 IFCC LS BC05630 R1:1 x 100ml R2: 1 x 5ml 2.4°C smylase Benzylidene Blockd-pNPG7 Lyo BC05630 R1:1 x 100ml R2: 1 x 5ml 2.4°C smylase CMPG3 Mono LS BXC0568 R1: 6 x 10ml 2.4°C smylase CMPG3 Mono LS BXC0560 R1: 5 x 10ml R2: 5 x 5ml 2.4°C smylase CMPG3 Mono LS BXC0564 R1: 5 x 10ml R2: 5 x 2ml 2.4°C smylase CMPG3 Mono LS BXC04764 R1: 5 x 10ml R2: 1 x 3ml 2.4°C smylase CMPG3 LS BXC04764 R1: 5 x 10ml R2: 1 x 3ml 2.4°C smylase Parti Informativitimetric (Without LS BXC03790 R1: 1 x 30ml R2: 1 x 3ml 2.4°C smit-CCP Urbilatex LS BXC03708 R1: 1 x 30ml R2: 1 x 3ml 2.4°C	ALT (GPT)	IFCC	Lyo	BXC0211E	R1: 1 x 45ml R2: 20 x 2ml	2-8°C
winglaseEPS-pNPG7 IFCCLSBXC0563BR1: 2 x 2 sml R2: 2 x 5ml2-8°CxinglaseEPS-pNPG7 IFCCLSBXC0563DR1: 1 x 100ml R2: 1 x 20ml2-8°CxinglaseCNPG3Mono LSBXC0564R1: 6 x 10ml2-8°CxinglaseCNPG3Mono LSBXC0564R1: 6 x 10ml2-8°CxinglaseCNPG3Mono LSBXC0564R1: 6 x 10ml2-8°Cxinglase, PancreaticEPS-pNPG7 IFCCLSBXC0564R1: 5 x 10ml R2: 5 x 2ml2-8°Cxinglase, PancreaticEPS-pNPG7 IFCCLSBXC0176AR1: 5 x 10ml R2: 1 x 2-8°C2-8°Cxinglatensin Converting Enzyme (ACE)Mono LSBXC0429AR1: 2 x 20ml R2: 1 x 2-8°C2-8°Cxinti Thrombin-IIIImmunoturbidimetric (Without 	Ammonia	GLDH	Mono LS	BXC0376A		2-8°C
ImplaseEPS-pNPG7 IFCCLSBXC0560R11 x 100ml R2: 1 x 2 are constrained by the Blocked pNPG7L9BXC0561R11 x 30ml R2: 1 x 3 constrained by the Blocked pNPG7L9BXC0562R11 x 30ml R2: 5 x 5ml2.8°CImplaseCNPG3Mono LSBXC0562R1: 6 x 10ml2.8°CR1: 1 x 30ml R2: 5 x 2ml2.8°CImplase, PancreaticCNPG3Mono LSBXC0562R1: 1 x 10ml R2: 5 x 2ml2.8°CImplase, PancreaticEPS-pNPG7 IFCCLSBXC0564R1: 5 x 10ml R2: 5 x 2ml2.8°CImplase, PancreaticEPS-pNPG7 IFCCLSBXC04204R1: 2 x 20ml R2: 1 x 8ml2.8°CImplase, PancreaticImplase Pre-dilutionLSBXC03794R1: 1 x 30ml R2: 1 x 3ml2.8°CImplace, PanceTurbilatexLSBXC03794R1: 1 x 30ml R2: 1 x 3ml2.8°CImplace, Pre-dilutionLSBXC03795R1: 1 x 30ml R2: 1 x 3ml2.8°CImplace, Pre-dilutionLSBXC03795R1: 1 x 30ml R2: 1 x 3ml2.8°CImplase, Pre-dilutionLSBXC03796R1: 1 x 30ml R2: 1 x 3ml2.8°C	Amylase	EPS-pNPG7 IFCC	LS	BXC0563A	R1: 1 x 25ml R2: 1 x 5ml	2-8°C
Implifies EPS/PAPP/FICE ES BR/0503 20ml 24 R Implies Benzylidene Blocked-pNPG7 Lyo BXC05614 R1:1 x 30ml R2: S x50 2.8°C Implies CNPG3 Mono LS BXC05624 R1: 6 x 10ml 2.8°C Implies parceatic CPS-pNPG7 IFCC Mono LS BXC05404 R1: 5 x 10ml R2: 5 x 30ml R2: 1 x	Amylase	EPS-pNPG7 IFCC	LS	BXC0563B	R1: 2 x 25ml R2: 2 x 5ml	2-8°C
mylaseCNPG3Mon LSBXC0562AR1: 6 x 10ml2-8°Cmylase, PancreaticEPS-pNPG7 IFCCLSBXC0562BR1: 10 x 10ml2-8°Cunglotensin Converting Enzyme (ACE)Mono LSBXC0176AR1: 5 x 10ml R2: 5 x 2ml2-8°Cunti Thrombin-IIIImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0379AR1: 1 x 30ml R2: 1 x 10ml R2: 5 x 10ml R4: 5 x 10ml R4: 1 x 1ml2-8°Cunti-CCPTurbilatexLSBXC0379CR1: 1 x 30ml R2: 1 x 10ml R4: 5 x 10ml R2: 1 x 10ml R4: 1 x 1ml(1 x1ml 1-8°C2-8°Cunti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml 1-8°C2-8°Cunti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml2-8°Cupolipoprotein A1Immunoturbidimetric (Without Sample Pre-dilution)LSBXC011AR1: 2 x 20ml R2: 1 x Bml 2-8°C2-8°CusoTurbidimetric Single PointLSBXC0501AR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC0501AR1: 1 x 10ml R2: 1 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC0501AR1: 1 x 10ml R2: 1 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC0501AR1: 1 x 10ml R2: 1 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC02	Amylase	EPS-pNPG7 IFCC	LS	BXC0563D		2-8°C
windlageCNPG3Mono LSBXC0562BR1:10 x 10ml2.4°Cwindlage, PancreaticEPS-pNPG7 IFCCLSBXC0176AR1:5 x 10ml R2: 5 x 20ml R2: 1 x 8ml2.4°Cwindlage, PancreaticImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0176AR1: 2 x 20ml R2: 1 x 8ml2.4°Cwinti Thrombin-IIIImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0379AR1: 1 x 30ml R2: 1 x 10ml	Amylase	Benzylidene Blocked-pNPG7	Lyo	BXC0561A	R1: 1 x 30ml R2: 5 x 5ml	2-8°C
And unspiceEPS-pNPG7 IFCCLSBXC0364AR1: S x 10ml R2: S x 2ml2-8°CunspiceImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0176AR1: 2 x 20ml R2: 1 x 8ml2-8°Cunti Thrombin-IIIImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0379AR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml2-8°Cunti-CCPTurbilatexLSBXC0379AR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml2-8°Cunti-CCPTurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml2-8°Cunti-CCPTurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°Cunti-CCPTurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°Cunti-CCPTurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°Cupsplipoprotein A1Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8°CusoTurbidimetric Single PointLSBXC051BR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC051CR1: 1 x 40ml R2: 1 x 10ml R4: 2 x 10ml R4: 1 x 1ml2-8°CusoColorimetric Single PointLSBXC0501BR1: 1 x 10ml R4: 2 x 10ml R4: 2 x 10ml R4: 1 x 1ml2-8°CusoTurbidimetric Single PointLSBXC0501CR1: 1 x 10ml R4: 1 x 1ml2-8°CusoColorimetric Single PointLSBXC0501BR1: 1 x 10ml R2: 1 x 10ml	Amylase	CNPG3	Mono LS	BXC0562A	R1: 6 x 10ml	2-8°C
Angiotensin Converting Enzyme (ACE)Mono LSBXC0176AR1: 5 x 10ml R4: 1 x 1ml2-8°CAnti Thrombin-IIIImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0379AR1: 2 x 20ml R2: 1 x 8ml2-8°CAnti-CCPTurbilatexLSBXC0379CR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml2-8°CAnti-CCPTurbilatexLSBXC0379CR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1 ml2-8°CAnti-CCPTurbilatexLSBXC0412AR1: 2 x 20ml R2: 1 x 8ml2-8°CApplipoprotein BImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0501AR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501CR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501CR1: 1 x 10ml R2: 1 x 10ml R4: 1 x 1ml2-8°C<	Amylase	CNPG3	Mono LS	BXC0562B	R1: 10 x 10ml	2-8°C
Intri Thrombin-IIIImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0429AR1: 2 x 20ml R2: 1 x 8ml 2 *8°C2 *8°CIntri-CCPTurbilatexLSBXC0379AR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml 2.6°C2 *8°CIntri-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml(1x1ml 2.6°C2 *8°CIntri-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml(1x1ml 2.6°C2 *8°CIntri-CCP **USA ONLY**TurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2 *8°CIntroductorImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml 2 *8°C2 *8°CIntroductorImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0412AR1: 2 x 20ml R2: 1 x 8ml 2 *8°C2 *8°CIssoTurbidimetric Single PointLSBXC0501AR1: 1 x 12 ml 10ml R4: 5 x 1ml 2 *8°C2 *8°CIssoTurbidimetric Single PointLSBXC0501AR1: 1 x 12 ml 10ml R4: 1 x 1ml 2 *8°C2 *8°CIssoTurbidimetric Single PointLSBXC0501CR1: 4 x 40ml R2: 2 x 10ml R1: 1 x 12 ml 2 *8°C2 *8°CIssoTurbidimetric Single PointLSBXC0202AR1: 1 x 12 ml 2 *8°C2 *8°CIssoTurbidimetric Single PointLSBXC0202AR1: 1 x 12 ml R2: 1 x 10ml R4: 1 x 1ml2 *8°CIssoTurbidimetric Single PointLSBXC0202AR1: 1 x 10ml R2: 2 x 10ml 2 *8°C<	Amylase, Pancreatic	EPS-pNPG7 IFCC	LS	BXC0564A	R1: 5 x 10ml R2: 5 x 2ml	2-8°C
Anti-Infombin-IIISample Pre-dilutionLSBXC0429AR1: 2 x 20mt R2: 1 x 8ml2-8 cAunti-CCPTurbilatexLSBXC0379AR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1.0.5ml2-8°CAunti-CCPTurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml(1x1ml)2-8°CAunti-CCP **USA ONLY**TurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°CAunti-CCP **USA ONLY**TurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°CAupolipoprotein A1Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8°CAugolipoprotein BImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0501AR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0501AR1: 2 x 20ml R2: 1 x 8ml2-8°CAusoTurbidimetric Single PointLSBXC0501AR1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0202AR1: 1 x 25ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0202AR1: 1 x 125ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0202AR1: 1 x 125ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLS <td>Angiotensin Converting Enzyme (ACE)</td> <td></td> <td>Mono LS</td> <td>BXC0176A</td> <td>R1: 5 x 10ml R4: 1 x 1ml</td> <td>2-8°C</td>	Angiotensin Converting Enzyme (ACE)		Mono LS	BXC0176A	R1: 5 x 10ml R4: 1 x 1ml	2-8°C
Intr-CCPIurbilatexLSBXC0379A10ml R4: 6 x 1 x 0.5ml2-8°CAunti-CCPTurbilatexLSBXC0379CR1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml(1x1ml L6)2-8°CAunti-CCP **USA ONLY**TurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°CAunti-CCP **USA ONLY**TurbilatexLSBXC0379DR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml2-8°CAunti-CCP **USA ONLY**TurbilatexLSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8°CAununoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8°CAupolipoprotein BImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0501AR1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0501BR1: 2 x 20ml R2: 1 x 8ml2-8°CAusoTurbidimetric Single PointLSBXC0501BR1: 1 x 40ml R2: 2 x 10ml R4: 1 x 1ml2-8°CAusoTurbidimetric Single PointLSBXC0501BR1: 1 x 125ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAusoColorimetric (Standard)LSBXC0202AR1: 1 x 100ml R2: 1 x 10ml R4: 1 x 10ml2-8°CAust (GOT)IFCCLSBXC0203AR1: 1 x 100ml R2: 1 x 20ml2-8°CAust (GOT)IFCCLSBXC0203AR1: 2 x 100ml R2: 2 x 20ml2-8°CAust (GOT)IFCCLSBXC0203AR1: 2 x 100ml R2: 2 x 20ml2-8°C	Anti Thrombin-III		LS	BXC0429A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Nuti-CCPTurbilatexLSBXC0379C10ml R4: 1 x 1ml(1x1ml) L62-8°CNuti-CCPTurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml)2-8°CNuti-CCP **USA ONLY**TurbilatexLSBXC0379BR1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml)2-8°CNuti-CCP **USA ONLY**Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml 2-8°C2-8°CNuti-CCP **USA ONLY**Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0412AR1: 2 x 20ml R2: 1 x 8ml 2-8°C2-8°CNuti-Copinportein A1Immunoturbidimetric Single PointLSBXC0501AR1: 2 x 40ml R2: 1 x 8ml 2-8°C2-8°CNuti-CopinTurbidimetric Single PointLSBXC0501AR1: 2 x 40ml R2: 1 x 1ml 10ml R4: 1 x 1ml2-8°CNuti-CopinTurbidimetric Single PointLSBXC0501AR1: 2 x 40ml R2: 2 x 10ml R2: 1 x 10ml R4: 1 x 1ml 2-8°C2-8°CNuti-CopinTurbidimetric Single PointLSBXC0501BR1: 1 x 125ml R4: 1 x 1ml 2-8°C2-8°CNuti-CopinColorimetric (Standard)LSBXC0202AR1: 1 x 125ml R4: 1 x 1ml 2-8°C2-8°CNuti-CopinIFCCLSBXC0203AR1: 1 x 100ml R2: 1 x 2-8°C2-8°CNuti-CopinIFCCLSBXC0203AR1: 2 x 10ml R2: 1 x 2-8°C2-8°CNuti-CopinIFCCLSBXC0203AR1: 1 x 100ml R2: 1 x 2-8°C2-8°CNuti-CopinIFCCLSBXC0203AR1: 5 x 100ml R	Anti-CCP	Turbilatex	LS	BXC0379A		2-8°C
IntractorIntroductor<	Anti-CCP	Turbilatex	LS	BXC0379C	10ml R4: 1 x 1ml(1x1ml	2-8°C
Number CCD ***USA ONLY**TurbitatexLSBXC0379B10ml R4: 6 x 1 x 1ml2-8° CApolipoprotein A1Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8° CApolipoprotein BImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0501AR1: 1 x 40ml R2: 1 x 8ml2-8° CASOTurbidimetric Single PointLSBXC0501AR1: 1 x 40ml R2: 1 x 1ml2-8° CASOTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml2-8° CASOTurbidimetric Single PointLSBXC0501BR1: 4 x 40ml R2: 4 x 10ml R2: 2 x 10ml R4: 1 x 1ml2-8° CASOTurbidimetric Single PointLSBXC0501CR1: 4 x 40ml R2: 4 x 10ml R2: 4 x 10ml R4: 1 x 1ml2-8° CAST (GOT)Colorimetric (Standard)LSBXC0202AR1: 1 x 125ml R2: 1 x 125ml R4: 1 x 5ml R6: 1 x 10ml R2: 1 x 20ml2-8° CAST (GOT)IFCCLSBXC0203AR1: 1 x 100ml R2: 1 x 20ml R2: 2 x 20ml2-8° CAST (GOT)IFCCLSBXC0203AR1: 2 x 100ml R2: 2 x 20ml2-8° C	Anti-CCP	Turbilatex	LS	BXC0379D		2-8°C
Spolipoprotein A1Sample Pre-dilution)LSBXC0411AR1: 2 x 20ml R2: 1 x 8ml2-8 CApolipoprotein BImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0412AR1: 2 x 20ml R2: 1 x 8ml2-8°CASOTurbidimetric Single PointLSBXC0501AR1: 1 x 40ml R2: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R2: 2 x 10ml R4: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R2: 2 x 10ml R4: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501CR1: 4 x 40ml R2: 4 x 10ml R4: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501CR1: 1 x 125ml R2: 1 x 10ml R4: 1 x 1ml2-8°CAST (GOT)IFCCLSBXC0202AR1: 1 x 100ml R2: 1 x 20ml R2: 2 x 20ml R2: 1 x 20ml R2: 2 x 20ml R2: 1 x 20ml R2: 1 x 20ml R2: 2 x 20ml R2: 2 x 20ml R2: 1 x 20ml R2: 2 x 20ml R2: 1 x 20ml R2: 2 x 20ml R2: 1 x 20ml R2: 1 x 20ml R2: 1 x 20ml R2:	Anti-CCP **USA ONLY**	Turbilatex	LS	BXC0379B		2-8°C
Sample Pre-dilution)LSBXC0412AR1: 2 x 20mt R2: 1 x 8mt2-8 °CASOTurbidimetric Single PointLSBXC0501AR1: 1 x 40ml R2: 1 x 1ml2-8°CASOTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R2: 1 x 10ml R2: 2 x 10ml R2: 2 x 10ml R2: 2 x 10ml R2: 2 x 10ml R2: 1 x	Apolipoprotein A1		LS	BXC0411A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
SOTurbidimetric Single PointLSBXC0501AToml R4: 1 x 1ml2-8 °CASOTurbidimetric Single PointLSBXC0501BR1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml2-8 °CASOTurbidimetric Single PointLSBXC0501CR1: 4 x 40ml R2: 4 x 10ml R4: 1 x 1ml2-8 °CASOTurbidimetric Single PointLSBXC0501CR1: 4 x 40ml R2: 4 x 10ml R4: 1 x 1ml2-8 °CAST (GOT)Colorimetric (Standard)LSBXC0202AR1: 1 x 125ml R2: 1 x 125ml R4: 1 x 5ml R6: 1 x 100ml2-8 °CAST (GOT)IFCCLSBXC0203AR1: 1 x 100ml R2: 1 x 20ml2-8 °CAST (GOT)IFCCLSBXC0203DR1: 2 x 100ml R2: 2 x 20ml2-8 °C	Apolipoprotein B		LS	BXC0412A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Independence LS BXC0501B Index 1 x 1ml 2-8 °C ASO Turbidimetric Single Point LS BXC0501C R1: 4 x 40ml R2: 4 x 10ml R4: 1 x 1ml 2-8 °C AST (GOT) Colorimetric (Standard) LS BXC0202A R1: 1 x 125ml R2: 1 x 125ml R4: 1 x 5ml R6: 1 x 100ml 2-8 °C AST (GOT) IFCC LS BXC0203A R1: 1 x 100ml R2: 1 x 20ml 2-8 °C AST (GOT) IFCC LS BXC0203B R1: 2 x 100ml R2: 2 x 20ml 2-8 °C	ASO	Turbidimetric Single Point	LS	BXC0501A		2-8°C
Introdumetric single Point LS BXC0501C R4: 1 x 1ml 2-8 C NST (GOT) Colorimetric (Standard) LS BXC0202A R1: 1 x 125ml R2: 1 x 125ml R4: 1 x 5ml R6: 1 x 100ml 2-8°C NST (GOT) IFCC LS BXC0203A R1: 1 x 100ml R2: 1 x 20ml 2-8°C NST (GOT) IFCC LS BXC0203D R1: 2 x 100ml R2: 2 x 20ml 2-8°C	450	Turbidimetric Single Point	LS	BXC0501B		2-8°C
XST (GOT) Colorimetric (Standard) LS BXC0202A 125ml R4: 1 x 5ml R6:1 2-8°C 2-8°C XST (GOT) IFCC LS BXC0203A R1: 1 x 100ml R2: 1 x 2-8°C 2-8°C XST (GOT) IFCC LS BXC0203A R1: 2 x 100ml R2: 2 x 20ml 2-8°C XST (GOT) IFCC LS BXC0203D R1: 2 x 100ml R2: 2 x 20ml 2-8°C	ASO	Turbidimetric Single Point	LS	BXC0501C		2-8°C
IFCC LS BXC0203A 20ml 20ml 2-8°C IST (GOT) IFCC LS BXC0203D 20ml R1: 2 x 100ml R2: 2 x 20ml 2-8°C IST (GOT) IFCC LS BXC0203E R1: 5 x 100ml R2: 1 x 2.8°C	AST (GOT)	Colorimetric (Standard)	LS	BXC0202A	125ml R4: 1 x 5ml R6:1	2-8°C
IFCC LS BXC0203D 20ml 2-8°C VST (GOT) IFCC LS BXC0203E R1: 5 x 100ml R2: 1 x 2-8°C	AST (GOT)	IFCC	LS	BXC0203A		2-8°C
	AST (GOT)	IFCC	LS	BXC0203D		2-8°C
	AST (GOT)	IFCC	LS	BXC0203F	R1: 5 x 100ml R2: 1 x 100ml	2-8°C

SECTION 1 | Clinical Chemistry

Description	Method	Format	Cat No.	Size	Storage
AST (GOT)	Colorimetric (Table)	LS	BXC0204A	R1: 1 x 100ml R2: 1 x 100ml	2-8°C
AST (GOT)		LS	BXC0205A	R1: 1 x 100ml R2: 1 x 25ml	2-8°C
AST (GOT)		LS	BXC0205B	R1: 2 x 100ml R2: 2 x 25ml	2-8°C
AST (GOT)	IFCC	Lyo	BXC0201B	R1: 2 x 105ml R2: 10 x 20ml	2-8°C
AST (GOT)	IFCC	Lyo	BXC0201C	R1: 1 x 105ml R2: 10 x 10ml	2-8°C
AST (GOT)	IFCC	Lyo	BXC0201D	R1: 5 x 50ml R2: 5 x 50ml	2-8°C
AST (GOT)	IFCC	Lyo	BXC0201E	R1: 1 x 45ml R2: 20 x 2ml	2-8°C
Beta-2-Microglobulin	Turbilatex	LS	BXC0090A	R1: 2 x 11ml R2: 2 x 4.3ml CAL: 4 x 1 x0.5ml	2-8°C
Beta-Hydroxybutyrate	3-HBDH/NAD	Lyo	BXC0545A	R1: 1 x 50ml R2: 5 x 10ml R4: 1 x 5ml	2-8°C
Bile Acids	Enzymatic	Lyo	BXC0581A	R1: 1 x 65ml R2: 1 x 20ml R3:6 x 10ml	2-8°C
Bile Acids	3-Alpha-HSD	LS	BXC0584A	R1: 1 x 30ml R2: 1 x 10ml R4:1 x 2ml	2-8°C
Bilirubin	Direct	LS	BXC0191A	R1: 2 x 50ml R2: 1 x 25ml	Room Temp
Bilirubin	Direct	LS	BXC0191B	R1: 2 x 100ml R2: 1 x 50ml	Room Temp
Bilirubin	Total	LS	BXC0192A	R1: 2 x 50ml R2: 1 x 25ml	Room Temp
Bilirubin	Total	LS	BXC0192B	R1: 2 x 100ml R2: 1 x 50ml	Room Temp
Bilirubin	Total & Direct Manual	LS	BXC0193A	R1: 1 x 100ml R2: 1 x 100ml R3: 1 x 50ml / 1 x 10ml	Room Temp
Bilirubin	Total & Direct Manual	LS	BXC0193B	R1: 2 x 100ml R2: 2 x 100ml R3:1 x 100ml /1 x 20ml	Room Temp
Bilirubin Direct	Vanadate Oxidation	LS	BXC0104A	R1: 2 x 70ml R2: 2 x 18ml	12-25°C
Bilirubin Total	Vanadate Oxidation	LS	BXC0105A	R1: 2 x 70ml R2: 2 x 18ml	2-8°C
Blood Alcohol	UV	LS	BXC0491A	R1: 3 x 20ml R2: 2 x 5ml R4: 1 x 2ml	2-8°C
C1 Inactivator	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0170A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
Calcium	AMP/CPC	LS	BXC0291A	R1: 1 x 125ml R2: 1 x 125ml R3:1 x 12.5ml R4: 1 x 5ml	15-25°C
Calcium	Arsenazo	Mono LS	BXC0292A	R1: 2 x 125ml R4: 1 x 5ml	15-25°C
Ceruloplasmin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0765A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
Chloride	Thiocyanate	Mono LS	BXC0281A	R1: 2 x 60ml STD L:1 x 5ml STD H:1 x 5ml	15-25°C
Cholesterol	CHOD-PAP	Lyo	BXC0262A	R1: 2 x 105ml R2: 10 x 20ml R4: 1 x 5ml	2-8°C
Cholesterol	CHOD-PAP	Lyo	BXC0262B	R1: 10 x 50ml R2: 10 x 50ml R4: 1 x 5ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	BXC0261A	R1: 2 x 60ml R4: 1 x 5ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	BXC0261B	R1: 6 x 60ml R4: 1 x 5ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	BXC0261C	R1: 8 x 250ml R4: 1 x 5ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	BXC0261D	R1: 4 x 250ml R4: 1 x 5ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	BXC0801A	R1: 5 x 20 R2: 2 x 10ml	2-8°C
Citrate (Urinary)	Enzymatic Colorimetric	LS	BXC0120A	R1: 2 x 10ml R2: 1 x 2ml R3: 2 x 10ml R4: 1 x 5ml Controls: 2 x 3ml	2-8°C
СК-МВ	Immunoinhibition	LS	BXC0452A	R1: 1 x 25ml R2: 1 x 5ml	2-8°C
СК-МВ	Immunoinhibition	LS	BXC0452B	R1: 2 x 25ml R2: 2 x 5ml	2-8°C
СК-МВ	Immunoinhibition	LS	BXC0452C	R1: 10 x 3ml R2: 1 x 6ml	2-8°C
СК-МВ	Immunoinhibition	Lyo	BXC0451A	R1: 1 x 35ml R2: 10 x 2.5ml R3:1 x 2ml	2-8°C

Clinical Chemistry | SECTION 1

Description	Method	Format	Cat No.	Size	Storage
СК-МВ	Immunoinhibition	Lyo	BXC0451B	R1: 1 x 105ml R2: 10 x 10ml R3:1 x 2ml	2-8°C
СК-МВ	Immunoinhibition	Lyo	BXC0451C	R1: 1 x 15ml R2: 5 x 2.5ml R3:1 x 2ml	2-8°C
СК-МВ 4+1	Immunoinhibition	LS	BXC0458A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
СК-МВ 4+1	Immunoinhibition	LS	BXC0458B	R1: 2 x 20ml R2: 1 x 10ml	2-8°C
CK-nac	Nac Activation, Serum Start, DGKC	LS	BXC0252A	R1: 2 x 25ml R2: 2 x 5ml	2-8°C
CK-nac	Nac Activation, Serum Start, DGKC	LS	BXC0252B	R1: 2 x 50ml R2: 2 x 10ml	2-8°C
CK-nac	Nac Activation, Serum Start, DGKC	LS	BXC0252C	R1: 6 x 10ml R2: 1 x 12ml	2-8°C
CK-nac	Nac Activation, Serum Start, IFCC	Lyo	BXC0251B	R1: 1 x 105ml R2: 10 x 10ml	2-8°C
CK-nac	Nac Activation, Serum Start, IFCC	Lyo	BXC0251D	R1: 1 x 35ml R2: 10 x 2.5ml	2-8°C
CK-nac 4+1	Nac Activation, Serum Start, DGKC	LS	BXC0253A	R1: 2 x 20ml R2: 1 x 10ml	2-8°C
CK-nac 4+1	Nac Activation, Serum Start, DGKC	LS	BXC0253B	R1: 2 x 40ml R2: 1 x 20ml	2-8°C
CO2 (Bicarbonate)	Enzymatic	Mono LS	BXC0152A	R1: 10 x 5ml R4: 1 x 1ml	2-8°C
CO2 (Bicarbonate)	Enzymatic	Mono LS	BXC0152B	R1: 10 x 10ml R4: 1 x 1ml	2-8°C
Complement C3	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0851A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Complement C4	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0861A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Copper	DBP (Colorimetric)	Mono LS	BXC0341A	R1: 2 x 25ml R4: 1 x 5ml	2-8°C
Copper (Urinary)	DBP (Colorimetric)	LS	BXC0342A	R1: 2 x 25ml R2: 1 x 10ml R4: 1 x 5ml	2-8°C
Copper (Urinary)	DBP (Colorimetric)	Mono LS	BXC0343A	R1: 2 x 25ml R4: 1 x 5ml	2-8°C
Creatinine	Jaffe without Deproteinization	LS	BXC0111A	R1: 1 x 120ml R2: 1 x 120ml R4: 1 x 5ml	Room Temp
Creatinine	Jaffe without Deproteinization	LS	BXC0111B	R1 3 x 60ml R2 3 x 60ml R4 1 x 5ml	Room Temp
Creatinine	Jaffe with Deproteinization	LS	BXC0112A	R1: 1 x 120ml R2: 1 x 120ml R4: 1 x 35ml	15-25°C
Creatinine	Jaffe without Deproteinization (One Month Stability)	LS	BXC0117A	R1: 1 x 120ml R2: 1 x 120ml R4: 1 x 5ml	2-8°C
Creatinine	Enzymatic, SOX	LS	BXC0113A	R1: 1 x 50ml R2: 1 x 25ml R4: 1 x 1ml	2-8°C
Creatinine	Enzymatic, SOX	LS	BXC0113B	R1: 2 x 50ml R2: 2 x 25ml R4: 1 x 1ml	2-8°C
Creatinine	Enzymatic, SOX	LS	BXC0113C	R1: 2 x 100ml R2: 2 x 50ml R4: 1 x 1ml	2-8°C
Creatinine TCA Concentrate	For Deproteinization	LS	BXC0114A	R1: 1 x 120ml	2-8°C
CRP	Immunoturbidimetric, (Linearity 280mg/L)	LS	BXC0384A	R1: 2 x 20ml R2: 1 x 8ml R4: 1 x 1ml(L6)	2-8°C
CRP	Immunoturbidimetric, Multipoint (Linearity 280mg/L)	LS	BXC0384B	R1: 2 x 20ml R2: 1 x 8ml R4: 1 x 6 x1ml	2-8°C
CRP (Ultra Sensitive)	Immunoturbidimetric, Multipoint	LS	BXC0383A	R1: 1 x 16ml R2: 1 x 4ml R4: 1 x 200ul	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	BXC0382A	R1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml(L6)	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	BXC0382B	R1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml(L6)	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	BXC0382C	R1: 4 x 40ml R2: 4 x 10ml R4: 1 x 1ml(L6)	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L) No Calibrator	LS	BXC0382D	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L) No Calibrator	LS	BXC0382E	R1: 2 x 40ml R2: 2 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L) No Calibrator	LS	BXC0382F	R1: 4 x 40ml R2: 4 x 10ml	2-8°C
Cystatin-C	Turbidimetric	LS	BXC0777A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
D3-Hydroxybutyrate	3-HBDH/NAD	LS	BXC0542A	R1: 1 x 50ml R2: 1 x 8ml R4: 1 x 5ml	2-8°C
D3-Hydroxybutyrate	3-HBDH/NAD	Lyo	BXC0541A	R1: 1 x 50ml R2: 5 x 10ml R4: 1 x 5ml	2-8°C

SECTION 1 | Clinical Chemistry

Description	Method	Format	Cat No.	Size	Storage
D-Dimer	Immunoturbidimetric, Multipoint	LS	BXC0787A	R1: 1 x 15ml R2: 1 x 5ml; 1x1ml calibrator	2-8°C
D-Dimer	Immunoturbidimetric, Multipoint	LS	BXC0787B	R1: 1 x 30ml R2: 1 x 10ml;1x1ml calibrator	2-8°C
Drabkins Solution (For use with BXC0551A)		LS	BXC0552A	R1: 4 x 200ml	Room Temp
Ferritin	Turbidimetric	LS	BXC0441A	R1: 1 x 20ml R2: 1 x 5ml R4: 1 x 1ml	2-8°C
Fibrinogen	Immunoturbidimetric, Multipoint	LS	BXC0442A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Fructosamine	Nitroblue Tetrazolium (NBT)	LS	BXC0591A	R1: 3 x 6ml R2: 3 x 14ml	2-8°C
GGT	Carboxy	LS	BXC0362A	R1: 1 x 60ml R2: 1 x 12ml	2-8°C
GGT	Carboxy	LS	BXC0362D	R1: 2 x 60ml R2: 2 x 12ml	2-8°C
GGT	Carboxy	Lyo	BXC0361A	R1: 1 x 60ml R2: 5 x 10ml	2-8°C
Glucose	Hexokinase	LS	BXC0103A	R1: 5 x 25ml R2: 1 x 25ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Lyo	BXC0102A	R1: 4 x 250ml R2: 4 x 250ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Lyo	BXC0102B	R1: 10 x 500ml R2: 10 x 500ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXC0101A	R1: 6 x 60ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXC0101B	R1: 6 x 125ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXC0101C	R1: 2 x 500ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXC0101D	R1: 4 x 250ml R4: 1 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXC0101E	R1: 10 x 500ml R4: 1 x 5ml	2-8°C
Glucose-6-Phosphate Dehydrogenase	Digitonin (UV)	Lyo	BXC0571A	R1: 2 x 50ml R2: 1 x 2ml R3:1 x 2ml R4: 1 x 20ml	2-8°C
Glucose-6-Phosphate Dehydrogenase	UV VISUAL	Lyo	BXC0574A	25T	2-8°C
Glucose-6-Phosphate Dehydrogenase	UV VISUAL	Lyo	BXC0574B	50T	2-8°C
Glucose-6-Phosphate Dehydrogenase	UV VISUAL	Lyo	BXC0574C	100T	2-8°C
Glucose-6-Phosphate Dehydrogenase	UV VISUAL	Lyo	BXC0574D	500T	2-8°C
Glucose-6-Phosphate Dehydrogenase	UV VISUAL	Lyo	BXC0574E	800T	2-8°C
Glucose-Fructose Kit			BXC0100A	TBC R1: 1 x 70ml R2: 5 x	2-8°C
Glutathione Peroxidase	Cumene Hydroperoxide	Lyo	BXC0551A	10ml R3:1 x 1ml R6:2 x 200ml R7:1 x 200ml	2-8°C
Glutathione Reductase	UV	Lyo	BXC0853A	R1: 1 x 50ml R2: 5 x 5ml R3: 5 x 3ml	2-8°C
Glycohaemoglobin	Colorimetric	LS	BXC0650A	R1: 1 x 60ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Glycohaemoglobin	Colorimetric	LS	BXC0650B	R1: 3 x 100ml R2: 1 x 50ml R4: 1 x 1ml	2-8°C
Haemoglobin	Drabkin's (Concentrate)	Mono LS	BXC0482A	R1: 1 x 1000ml R4: 1 x 1ml	2-8°C
Haemoglobin	Drabkin's (Concentrate)	Mono LS	BXC0482B	R1: 3 x 1000ml R4: 1 x 1ml	2-8°C
Haptoglobin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0496A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
HbA1c	Direct Enzymatic	LS	BXC0672A	R1: 1 x 21ml R2: 1 x 9ml R3:1 x 20 ml	2-8°C
HbA1c Modified Enzymatic	Modified Enzymatic	LS	BXC0670A	R1: 1 x 30ml R2: 1 x 10ml R3: 2 x 120ml R4: 2 x 0.5ml	2-8°C
HbA1c Modified Enzymatic	Modified Enzymatic	LS	BXC0670C	R1: 1 x 30ml R2: 1 x 10ml R3: 2 x 120ml	2-8°C
HDL Cholesterol	Direct (without Calibrator)	LS	BXC0421A	R1: 3 x 10ml R2: 1 x 10ml	2-8°C
HDL Cholesterol	Direct (without Calibrator)	LS	BXC0421B	R1: 6 x 30ml R2: 3 x 20ml	2-8°C
HDL Cholesterol	Direct (without Calibrator)	LS	BXC0421C	R1: 4 x 70ml R2: 2 x 50ml	2-8°C
HDL Cholesterol	Direct with Calibrator	LS	BXC0421D	R1: 3 x 10ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C

Clinical Chemistry | SECTION 1

HR.CholesteryInterminationInterm	Description	Method	Format	Cat No.	Size	Storage
Filt ConsisteredDirect with CalibratorBABALLARIEBALLARIE28 / 28 / 28 / 28 / 28 / 28 / 28 / 28 /	HDL Cholesterol	Direct with Calibrator	LS	BXC0421E		2-8°C
Hu ConsisterioriDescriptionDescriptionDescriptionDescriptionDescriptionDescriptionHoncosysteinEnsymatic U/VLSBXC03924R1: 26 (6m) Reit StarDescriptionHuman Free Kappa Light ChainsTurbiatesLSBXC03924R1: 26 (2m) Reit StarDescriptionIghtTurbiatesTurbiatesLSBXC03924R1: 26 (2m) Reit StarDescriptionIghtTurbiatesTurbiatesLSBXC03924R1: 26 (2m) Reit StarDescriptionIghtTurbiatesTurbiatesLSBXC03940R1: 22 (2m) Reit StarDescriptionIghtImmanotribulinetic (VM)LSBXC0714R1: 22 (2m) Reit StarDescriptionIghtImmanotribulinetic (VM)LSBXC0214R1: 22 (2m) Reit StarDescriptionIghtImmanotribulinetic (VM)LSBXC0214R1: 22 (2m) Reit StarDescriptionIghtImmanotribulinetic (VM)LSBXC0214R1: 24 (2m) Reit StarDescriptionIghtFerreineLSBXC0214R1: 24 (2m) Reit StarDescriptionIghtFerreineLSBXC0214R1: 25 (2m) Reit StarDesc	HDL Cholesterol	Direct with Calibrator	LS	BXC0421F		2-8°C
HonocytelineEngrandic U/LSBXC09404R11 x 1 cm R2: 1 x 4m2 drCHuman Free Kappa Light ChainsTurblatexLSBXC0320R11 x 2 cm R2: 1 x 7m2 drCIgASimple ProductionLSBXC0710R12 x 2 cm R2: 1 x 8m2 drCIgGSimple ProductionLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGSimple ProductionLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGSimple ProductionLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 1 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 1 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 2 cm R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720R12 x 10m R2: 1 x 8m2 drCIgGImmonobilinetic (WholeLSBXC0720	HDL Cholesterol	Direct with Calibrator	LS	BXC0421G		2-8°C
Human Free Kappa Light ChainsTurbilatesLSBXC0322Site SectorSite SectorSectorHuman Free Lambda Light ChainsTurbilatesLSBXC0400Rith SectorSectorSectorIgASample PredilationLSBXC0701Rith SectorSectorSectorSectorIgGSample PredilationLSBXC0713Rith SectorSectorSectorSectorIgGSample PredilationLSBXC0721ARith SectorSectorSectorSectorIgGSample PredilationLSBXC072ARith SectorSectorSectorSectorIgGSample PredilationLSBXC022ARith SectorSectorSectorIgGFerreacLSBXC023ARith SectorSectorSectorIronFerreacineLSBXC023ARith SectorSectorSectorIronChromazurol BManuco LiggBXC023ARith SectorSectorIronChromazurol BManuco LiggBXC023ARith SectorSectorIronChromazurol BManu Chromazurol BBXC023ARith SectorSect	HDL Cholesterol	Precipitant	Mono LS	BXC0422A	R1: 2 x 60ml R4: 1 x 5ml	15-25°C
Human Pree Kappa Light ChainsTurblatexLsNUMBER <t< th=""><th>Homocysteine</th><th>Enzymatic UV</th><th>LS</th><th>BXC0690A</th><th>R1: 1 x 16ml R2: 1 x 4ml</th><th>2-8°C</th></t<>	Homocysteine	Enzymatic UV	LS	BXC0690A	R1: 1 x 16ml R2: 1 x 4ml	2-8°C
Human Pred Lamoda Light ChainsLutroblackLob BAC 34 millARE 5 A millARE 5 willAPE 6 millIgASample Pred ditationLSBAC 0701AR1: 2 x 20m R2: 1 x 8ml2.4°CIgESample Pred ditationLSBAC 071AR1: 2 x 20m R2: 1 x 8ml2.4°CIgGSample Pred ditationLSBAC 071AR1: 2 x 20m R2: 1 x 8ml2.4°CIgGSample Pred ditationLSBAC 073AR1: 2 x 20m R2: 1 x 8ml2.4°CIgGSample Pred ditationLSBAC 073AR1: 2 x 20m R2: 1 x 8ml2.4°CIronFerrozineLSBAC 073AR1: 2 x 20m R2: 1 x 8ml2.4°CIronFerrozineLSBAC 073AR1: 2 x 20m R2: 2 x 8ml2.4°CIronFerrozineLSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronChromazurol BMon LSBAC 073AR1: 2 x 50m R4: 1 x 5ml2.4°CIronMarge Pred ditationLSBAC 074	Human Free Kappa Light Chains	Turbilatex	LS	BXC0332A		2-8°C
ignignignignigninteractio	Human Free Lambda Light Chains	Turbilatex	LS	BXC0380A		2-8°C
igegample Pre-ditutionLSBACUYAtotal R4:: tx imil24°Cigglimit mutualimetric (without sample Pre-ditution/without sample Pre-ditution/without isample Pre-ditution/wit	IgA		LS	BXC0701A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
BoSample Pre-dilution)ISBKU721APERCURNAL PLANS2.48 CIgMSample Pre-dilution)LSBKC0731APEL2 X201H R21 X BMI2.46°CIronFerreeLSBKC023AC12 X201H R21 X BMI2.40°CIronFerreaLSBKC023AC12 X201H R21 X SMI2.40°CIronFerreaLSBKC023AC12 X201H R21 X SMI2.40°CIronChromazno BMono LSBKC023AC12 X001H R21 X SMI2.40°CIron/UBCFerreneLSBKC023AC11 X001H R22 X SMI2.40°CIron/UBCImmunoturbidimetric (WithoutLSBKC023AC11 X1001H R22 X SMI2.40°CIron/UBCImmunoturbidimetric (WithoutLSBKC023AC11 X1001H R22 X SMI2.40°CIron/UBCImmunoturbidimetric (WithoutLSBKC023AAC11 X1001H R22 X SMI2.40°CIron/UBCImmunoturbidimetric (WithoutLSBKC023AAC11 X1001H R22 X SMI2.40°CIron/UBCImmunoturbidimetric (WithoutLSBKC023AAC11 X1001H R22 X SMI2.40°CIron Cate DebydrogenaseLactate DefyragenaseLSBKC024AAR15 X201H R21 X SMI2.40°CIron Cate Debydrogenase (LD)Pyruwate-A Lactate, DGKCLSBKC024AAR15 X201H R21 X SMI2.40°CIron Cate Debydrogenase (LD)Pyruwate-A Lactate, DGKCLSBKC024AAR15 X201H R21 X SMI2.40°CIron Cate Debydrogenase (LD)Pyruwate-A Lactate, DGKCLSBKC043AIR15 X101H R21 X SMI2.4	IgE		LS	BXC0751A		2-8°C
igmsample Pre-dilutionLsBKCU2314R12 x 123ml R21 x 12ml2.4 °CironFerreneLsBKC0232AST12 x 123ml R21 x 5ml2.4 °CironFerrozineLsBKC0235AST12 x 40ml R21 x 5ml2.4 °CironFerrozineLsBKC0235AST12 x 40ml R21 x 5ml2.4 °CironChromazurol BMonol SBKC0235AST12 x 40ml R21 x 5ml2.4 °Ciron/UBCChromazurol BMonol SBKC0235AST12 x 40ml R21 x 5ml2.4 °Ciron/UBCEerreneLsBKC0235AST12 x 40ml R21 x 5ml2.4 °Ciron/UBCEerreneLsBKC0235AST12 x 40ml R21 x 5ml2.4 °CLactateEerreneLsBKC0235AST12 x 10ml R21 x 10ml2.4 °CLactateEerreneLsBKC0235AST12 x 10ml R21 x 10ml2.4 °CLactateEerreneLsBKC0235AST12 x 10ml R21 x 2ml2.4 °CLactateDehydrogenaseLactate > Pyruvate, DCKCLsBKC0235AST13 x 10ml R21 x 2ml2.4 °CLactate Dehydrogenase (LD)Ipyruate-> Lactate, DCKCLsBKC0234AST2 x 10ml R21 x 2ml2.4 °CLactate Dehydrogenase (LD)Ipyruate-> Lactate, DCKCLsBKC033HST2 x 10	lgG		LS	BXC0721A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
IronJerreneLSBAL0220GAL01240GAL01240JerIronFerrozineFerrozineLSBXC0230R1:2 x40ml R2: 2 SMR4:00ml R2: 1 x SmlR4:00ml R2: 1 x Sml <th>IgM</th> <th>•</th> <th>LS</th> <th>BXC0731A</th> <th>R1: 2 x 20ml R2: 1 x 8ml</th> <th>2-8°C</th>	IgM	•	LS	BXC0731A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
IronLSEXCU334Tom R4: 1 x Sm128 CIronFerrozineFerrozineLSBXC02345Cit X MM R2: 2 X28 CIronChomazurol BMono LSBXC02345Cit X SMI R2: 1 X SMI28 CIron/UBCFerreneLSBXC02345Cit X SMI R2: 1 X SMI28 CKappa Light ChainsImmunoturbidimetric (Mithout Sample Pre-dilution)LSBXC02344Cit X SMI R2: 1 X SMI28 CLactateEnzymatic (Colorimetric)VoBXC06240R1: 2 X SMI R2: 1 X SMI28 CLactate DehydrogenaseLactate > Pyruvate, DGXCLSBXC02344R1: S XOMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC02344R1: S XOMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC02345R1: S XOMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC02442R1: S XOMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC03341R1: X SMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC03341R1: X SMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC03341R1: X SMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC03341R1: X SMI R2: 1 X SMI28 CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGXCLSBXC03341R1: X SMI	Iron	Ferrene	LS	BXC0232A		2-8°C
IronLisEXCU33620m R4: 1 x 5m124 CIronChomazuol BMono LS8XC0236A1:2 x 50m R4: 1 x 5m12.4°CIron/UIBCErreneLisBxC0334AC1: 2 x 50m R4: 1 x 5m12.4°CKappa Light ChainsImmunoturbidimetric (Without)LisBxC0334AR1: 2 x 20m R2: 1 x 5m12.4°CLactateErrymatic (Colorimetric)LisBxC0621AR1: x 105m LB2 x 15m12.4°CLactate DehydrogenaseErzymatic (Colorimetric)LisBxC0621AR1: x 105m LB2 x 15m12.4°CLactate DehydrogenaseEatate > Pyruvate, DGKCLisBXC0624AR1: s x 20m R2: 1 x 20m2.4°CLactate DehydrogenaseEatate > Pyruvate, DGKCLisBXC024AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLisBXC024AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLisBXC024AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Direct (without Calibrator)LisBXC034AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Direct (without Calibrator)LisBXC034AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Direct (without Calibrator)LisBXC034AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Direct (without Calibrator)LisBXC034AR1: s x 20m R2: 1 x 20m2.4°CLactate Dehydrogenase (LD)Direct (without Calibrator)Li	Iron	Ferrozine	LS	BXC0235A		2-8°C
Iron/UBCFereneISBCC024 C_{12}^{11} Mind B2.31 Mind S1.51 Mind B1.51 Mind S1.51 Mind B1.51 Mind	Iron	Ferrozine	LS	BXC0235B		2-8°C
iron/UBCErreneLSBXC02344C12 stom Ert stom C13 stom Ert stom C13 stom Ert stom C13 stom Ert stomSeveKapa Light Chainsmanuferdenilution/C000LSBXC0314R1× a Xon R21 x and R21 x and <br< th=""><th>Iron</th><th>Chromazurol B</th><th>Mono LS</th><th>BXC0236A</th><th>R1: 2 x 50ml R4: 1 x 5ml</th><th>2-8°C</th></br<>	Iron	Chromazurol B	Mono LS	BXC0236A	R1: 2 x 50ml R4: 1 x 5ml	2-8°C
Kappa Light ChainsSample Pre-dilutionISBX.03314R1: 2x 20mit R2: 1x 20mitZ-8 CLactateEnzymatic (Colorimetric)LyoBX.06214R1: 1x 105mit R2: 1x 5 mitZ-8°CLactateEnzymatic (Colorimetric)Mono LSBX.06224R1: 2x Somt R4: 1x 5 mitZ-8°CLactateDehydrogenaseLactate > Pyruvate, DGKCLSBX.002430R1: 5x 20mit R2: 1x 20mitZ-8°CLactate DehydrogenaseLactate > Pyruvate, DGKCLSBX.002420R1: 10x 10ml R2: 2xZ-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBX.002420R1: 10x 10ml R2: 2xZ-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBX.00314R1: 2x 20mit R2: 1x 20mitZ-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBX.00314R1: 2x 20mit R2: 1x 20mitZ-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBX.00314R1: 2x 20mit R2: 1x 20mitZ-8°CLactate Dehydrogenase (LD)Direct (without Calibrator)LSBX.00314R1: 2x 20mit R2: 1x 20mitZ-8°CLDL CholesterolDirect (without Calibrator)LSBX.004316R1: 4x 70mit R2: 1x 20mitZ-8°CLDL CholesterolDirect (with Calibrator)LSBX.004316R1: 4x 70mit R2: 1x 20mitZ-8°CLDL CholesterolDirect (with Calibrator)LSBX.004316R1: 4x 70mit R2: 1x 20mitZ-8°CLDL CholesterolDirect (with Calibrator)LSBX.004316R1: 4x 70mit R2: 1x 20mitZ	Iron/UIBC	Ferrene	LS	BXC0234A	C:1 x 10ml D:1 x 10ml E:1 x 10ml F:1 x 100ml	2-8°C
LactateEnzymatic (colorimetric)EyoBXC0621AGmL R4:1 x Sml2-8°CLactateEnzymatic (Colorimetric)Mano LSBXC0622AR1:2 x Soml R4:1 x Sml2-8°CLactate DehydrogenaseLactate > Pyruvate, DGKCLSBXC0243AR1:5 x 20ml R2:1 x 20ml2-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC0242AR1:5 x 20ml R2:1 x 20ml2-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC0242AR1:5 x 20ml R2:1 x 20ml2-8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC0381AR1:2 x 20ml R2:1 x 80ml2-8°CLactate Dehydrogenase (LD)Direct (without Calibrator)LSBXC0381AR1:3 x 10ml R2: x2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431AR1:6 x 30ml R2: 1 x2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1:6 x 30ml R2: 3 x2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431CR1:4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431CR1:4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431CR1:4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431CR1:4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431CR1:4 x 70ml R2: 1 x 50ml R2: 1	Kappa Light Chains		LS	BXC0331A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate DehydrogenaseLactate > Pyruvate, DGKCLSBXC0243AR1: 5 x 20ml R2: 1 x 20ml 10ml2.8°CLactate DehydrogenaseLactate > Pyruvate, DGKCLSBXC0243BR1: 1 x 10ml R2: 2 x 10ml2.8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC0242BR1: 5 x 20ml R2: 1 x 20ml 10ml2.8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC0243BR1: 2 x 20ml R2: 1 x 20ml 	Lactate	Enzymatic (Colorimetric)	Lyo	BXC0621A		2-8°C
Lactate DehydrogenaseLactate -> Pyruvate, DGKCLSBXC0243BR1: 10 x 10ml R2: 2 x 10ml2-8°CLactate Dehydrogenase (LD)Pyruvate -> Lactate, DGKCLSBXC0242AR1: 5 x 20ml R2: 1 x 20ml2-8°CLactate Dehydrogenase (LD)Pyruvate -> Lactate, DGKCLSBXC0243BR1: 10 x 10ml R2: 2 x 10 x 10ml R2: 2 x 2-8°C2-8°CLambda Light ChainsImmunoturbidimetric (Without Sample Pre-dilution)LSBXC031AR1: 2 x 20ml R2: 1 x 8ml 2 x 9°C2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml 20ml R2: 3 x 20ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431BR	Lactate	Enzymatic (Colorimetric)	Mono LS	BXC0622A	R1: 2 x 50ml R4: 1 x 5ml	2-8°C
Lactate behydrogenaseLactate > Pyruvate, DLKCLSBXC0243810ml2.8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC02424R1: 5 x 20ml R2: 1 x 20ml2.8°CLactate Dehydrogenase (LD)Pyruvate > Lactate, DGKCLSBXC03418R1: 10 x 10ml R2: 2 x2.8°CLambda Light ChainsImmunoturbidimetric (Without Sample Pre-dilution)LSBXC03104R1: 3 x 10ml R2: 1 x2.8°CLDL CholesterolDirect (without Calibrator)LSBXC04318R1: 3 x 10ml R2: 1 x2.8°CLDL CholesterolDirect (without Calibrator)LSBXC04318R1: 4 x 70ml R2: 2 x 20ml2.8°CLDL CholesterolDirect (without Calibrator)LSBXC04318R1: 4 x 70ml R2: 3 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 3 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04310R1: 4 x 70ml R2: 3 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 3 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 2 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 2 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 1 x 20ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC04318R1: 4 x 70ml R2: 1 x 20ml2.8°CLDL Cho	Lactate Dehydrogenase	Lactate -> Pyruvate, DGKC	LS	BXC0243A	R1: 5 x 20ml R2: 1 x 20ml	2-8°C
Lactate Dehydrogenase (LD)Pyruvate -> Lactate, DGKCLSBXC0242BR1: 10 x 10ml R2: 2 x 10ml2-8°CLambda Light ChainsImmunoturbidimetric (Without Sample Pre-dilution)LSBXC0381AR1: 2 x 20ml R2: 1 x 8ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431AR1: 5 x 30ml R2: 3 x 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1: 6 x 30ml R2: 3 x 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431DR1: 3 x 10ml R2: 1 x 10ml 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431DR1: 3 x 10ml R2: 1 x 10ml R2: 1 x 10ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431AR1: 2 x 10ml R2: 1 x 10ml R2: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R1: 1 x 10ml R2: 1 x 10ml R2: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 1 x	Lactate Dehydrogenase	Lactate -> Pyruvate, DGKC	LS	BXC0243B		2-8°C
Lactate Dehydrogenase (LD)Pyrtvate-> Lactate, DCRCLSBC0242B10ml248 C10mlImmunoturbidimetric (WithoutLSBXC0381AR1: 2 x 20ml R2: 1 x 8ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431AR1: 3 x 10ml R2: 1 x 10ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1: 6 x 30ml R2: 3 x 20ml2-8°CLDL CholesterolDirect (without Calibrator)LSBXC0431CR1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1: 3 x 10ml R2: 1 x 10ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 1 x 70ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 4 x 70ml R2: 2 x 50ml2-8°CLG CholesterolPrecipitantK	Lactate Dehydrogenase (LD)	Pyruvate -> Lactate, DGKC	LS	BXC0242A	R1: 5 x 20ml R2: 1 x 20ml	2-8°C
Lambda Light ChainsSample Pre-dilutionLSBXC0381AR1:2 X 20m1 R2:1 X 8m12.8 CLDL CholesterolDirect (without Calibrator)LSBXC0431AR1:3 X 10m1 R2:1 X 10m1 R2:1 X 20m12.8 °CLDL CholesterolDirect (without Calibrator)LSBXC0431BR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (without Calibrator)LSBXC0431CR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1:3 X 10m1 R2:1 X 10m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431DR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:2 X 50m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:1 X 10m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:1 X 10m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:1 X 10m12.8 °CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1:4 X 70m1 R2:1 X 10m12.8 °CLigase (UV)Triacylglycerol Li	Lactate Dehydrogenase (LD)	Pyruvate -> Lactate, DGKC	LS	BXC0242B		2-8°C
LDL CholesterolDirect (without Calibrator)LSBXC0431A10ml2.8 °CLDL CholesterolDirect (without Calibrator)LSBXC0431B $R_{1:6 \times 30ml R2: 3 \times 20ml}$ 2.8°CLDL CholesterolDirect (without Calibrator)LSBXC0431CR1:4 × 70ml R2: 2 × 50ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431D $R_{1:3 \times 10ml R2: 1 \times 10ml}$ 2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431F $R_{1:4 \times 70ml R2: 2 \times 50ml}$ 2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431F $R_{1:4 \times 70ml R2: 2 \times 50ml}$ 2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431F $R_{1:4 \times 70ml R2: 2 \times 50ml}$ 2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431F $R_{1:4 \times 70ml R2: 2 \times 50ml}$ 2.8°CLDL CholesterolPrecipitantMono LSBXC0431FR1:4 × 70ml R2: 2 × 50ml}2.8°CLGutamatePrecipitantKSBXC0431FR1:4 × 70ml R2: 1 × 50ml R2: 1 × 50ml2.8°CLipase (UV)Triacylglycerol LipaseLSBXC0511AR1:4 × 70ml R2: 1 × 30ml2.8°CLipoprotein (a)Immunoturbidimetric (WithoutLSBXC0130AR1:2 × 20ml R2: 1 × 80ml2.8°CLipoprotein (a)EnzymaticLSBXC0130AR1:2 × 20ml R2: 1 × 80ml2.8°CLipoprotein (a)EnzymaticLSBXC0130AR1:2 × 20ml R2: 1 × 80ml2.8°CLipoprotein (a)EnzymaticLSBXC0130AR1:1 × 20ml	Lambda Light Chains		LS	BXC0381A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
LDL CholesterolDirect (without Calibrator)LSBXC0431B20ml248 CLDL CholesterolDirect (without Calibrator)LSBXC0431CR1:4 x 70ml R2: 2 x 50ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1:6 x 30ml R2: 1 x 10ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1:6 x 30ml R2: 3 x 20ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1:4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolPrecipitantMono LSBXC0431ER1:4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolColorimetricMono LSBXC0431ER1:2 x 60ml R4: 1 x 5ml 2 x 8°C2-8°CLipase (UV)Triacylglycerol LipaseLSBXC0511AR1:1 x 35ml R2: 10 x 2 5ml R4: 1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1:2 x 20ml R2: 1 x 8ml 2 x 8ml2-8°CLithiumEnzymaticEnzymaticLSBXC013CAR1:1 x 20ml R2: 1 x 8ml 2 x 8ml2-8°C	LDL Cholesterol	Direct (without Calibrator)	LS	BXC0431A		2-8°C
LDL CholesterolDirect (with Calibrator)LSBXC0431DR1: 3 x 10ml R2: 1 x 10ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 6 x 30ml R2: 3 x 20ml R4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolPrecipitantMono LSBXC0431FR1: 2 x 60ml R4: 1 x 5ml 2 8°C2-8°CLGutamateColorimetricLSBXC0061AR1: 2 x 10ml R2: 1 x 10ml R4: 1 x 3ml 2 8°C2-8°CLipase (UV)Triacylglycerol LipaseLyoBXC0130AR1: 2 x 20ml R2: 1 x 8ml 2 5ml R4: 1 x 3ml 2 8°C2-8°CLipoprotein (a)Immunoturbidimetric (Without) Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml 2 1 x 3ml R2: 10 x 2 5ml R4: 1 x 3ml 2 8°C2-8°CLiphiumImmunoturbidimetric (Without) Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml 2 1 x 3ml 2 1 x 3ml R2: 10 x 2 5ml R4: 1 x 3ml 2 8°C2-8°C	LDL Cholesterol	Direct (without Calibrator)	LS	BXC0431B		2-8°C
LDL CholesterolDirect (with Calibrator)LSBXC0431DR4: 1 x 1ml28°CLDL CholesterolDirect (with Calibrator)LSBXC0431ER1: 6 x 30ml R2: 3 x 20ml R4: 1 x 1ml2.8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2.8°CLDL CholesterolPrecipitantMono LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2.8°CLGlutamatePrecipitantKISBXC041AFR1: 2 x 60ml R4: 1 x 5ml S001AF2.8°CLipase (UV)ColorimetricLSBXC051AFR1: 2 x 10ml R2: 1 x 3ml S0ml R4: 1 x 3ml2.8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml S001AF2.8°CLiphiumEnzymaticLSBXC0130AR1: 2 x 20ml R2: 1 x 8ml S0ml R4: 1 x 3ml2.8°C	LDL Cholesterol	Direct (without Calibrator)	LS	BXC0431C	R1: 4 x 70ml R2: 2 x 50ml	2-8°C
LDL CholesterolDirect (with Calibrator)LSBXC0431FR4: 1 x 1ml2-8°CLDL CholesterolDirect (with Calibrator)LSBXC0431FR1: 4 x 70ml R2: 2 x 50ml R4: 1 x 1ml2-8°CLDL CholesterolPrecipitantMono LSBXC0432AR1: 2 x 60ml R4: 1 x 5ml2-8°CL-GlutamateLSBXC0061AR1: 4x20ml2-8°CLipaseColorimetricLSBXC0511AR1: 2 x 10ml R2: 1 x 10ml R4: 1 x 3ml2-8°CLipase (UV)Triacylglycerol LipaseLyoBXC0512AR1: 2 x 20ml R2: 1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 3ml2-8°CLithiumEnzymaticLSBXC0125AR1: 1 x 20ml R2: 1 x 3ml2-8°C	LDL Cholesterol	Direct (with Calibrator)	LS	BXC0431D	R4: 1 x 1ml	2-8°C
LDL CholesterolDirect (with Calibrator)LSBXC0431FR4: 1 x 1ml2-8°CLDL CholesterolPrecipitantMono LSBXC0432AR1: 2 x 60ml R4: 1 x 5ml2-8°CL-GlutamateLSBXC0061AR1: 4x20ml2-8°CLipaseColorimetricLSBXC0511AR1: 2 x 10ml R2: 1 x 3ml2-8°CLipase (UV)Triacylglycerol LipaseLyoBXC0512AR1: 1 x 35ml R2: 10 x 2.5ml R4: 1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml2-8°CLithiumEnzymaticLSBXC0125AR1: 1 x 20ml R2: 1 x 10ml2-8°C	LDL Cholesterol	Direct (with Calibrator)	LS	BXC0431E		2-8°C
L-GlutamateLSBXC0061AR1:4x20ml2-8°CLipaseColorimetricLSBXC0511AR1:2 x 10ml R2:1 x 3ml2-8°CLipase (UV)Triacylgycerol LipaseLyoBXC0512AR1:1 x 35ml R2:10 x 2.5ml R4:1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1:2 x 20ml R2:1 x 8ml2-8°CLithiumEnzymaticLSBXC0125AR1:1 x 20ml R2:1 x 8ml2-8°C	LDL Cholesterol	Direct (with Calibrator)	LS	BXC0431F		2-8°C
LipaseColorimetricLSBXC0511AR1: 2 x 10ml R2: 1 x 10ml R4: 1 x 3ml2-8°CLipase (UV)Triacylglycerol LipaseLyoBXC0512AR1: 1 x 35ml R2: 10 x 2.5ml R4: 1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml2-8°CLithiumEnzymaticLSBXC0125AR1: 1 x 20ml R2: 1 x 8ml2-8°C	LDL Cholesterol	Precipitant	Mono LS	BXC0432A	R1: 2 x 60ml R4: 1 x 5ml	2-8°C
LipaseColorimetricLisBXC0511A10ml R4: 1 x 3ml2-8 CLipase (UV)Triacylglycerol LipaseLyoBXC0512AR1: 1 x 35ml R2: 10 x 2.5ml R4: 1 x 3ml2-8°CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml2-8°CLithiumEnzymaticLSBXC0125AR1: 1 x 20ml R2: 1 x 10ml2-8°C	L-Glutamate		LS	BXC0061A	R1:4x20ml	2-8°C
Lipose (UV)Inacylgiscerot LipaseLyoBXC0512A2.5ml R4: 1 x 3ml2-8 CLipoprotein (a)Immunoturbidimetric (Without Sample Pre-dilution)LSBXC0130AR1: 2 x 20ml R2: 1 x 8ml2-8°CLithiumEnzymaticLSBXC0125AR1: 1 x 20ml R2: 1 x 10ml2-8°C	Lipase	Colorimetric	LS	BXC0511A		2-8°C
Lipoprotein (a) Sample Pre-dilution) LS BXC0130A R1: 2 x 20ml R2: 1 x 8ml 2-8 C Lithium Enzymatic LS BXC0125A R1: 1 x 20ml R2: 1 x 10ml 2-8 C	Lipase (UV)	Triacylglycerol Lipase	Lyo	BXC0512A		2-8°C
LICHIUM ENZYMATIC LS BXCUT25A 10ml 2-8 C	Lipoprotein (a)		LS	BXC0130A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lithium CalibratorOnly for Enzymatic LithiumLSBXC0141A1 x2x1ml2-8°C	Lithium	Enzymatic	LS	BXC0125A		2-8°C
	Lithium Calibrator	Only for Enzymatic Lithium	LS	BXC0141A	1 x2x1ml	2-8°C

SECTION 1 | Clinical Chemistry

Description	Method	Format	Cat No.	Size	Storage
L-Lactic Acid	UV	Lyo	BXC0119A	R1: 1 x 50ml R2: 4 x 10ml R3: 1 x 1ml R4: 1 x 5ml	2-8°C
L-Malic Acid	UV	LS	BXC0118A	R1: 1 x 50ml R2: 4 x 10ml R3: 1 x 1ml R4: 1 x 5ml	2-8°C
Magnesium	Calmagite	LS	BXC0351A	R1: 1 x 60ml R2: 1 x 60ml R4: 1 x 5ml	15-25°C
Magnesium	Calmagite	LS	BXC0351B	R1: 2 x 20ml R2: 2 x 20ml R4: 1 x 5ml	15-25°C
Magnesium	Xylidyl Blue	Mono LS	BXC0352A	R1: 2 x 60ml R4: 1 x 5ml	15-25°C
Magnesium	Xylidyl Blue	Mono LS	BXC0352B	R1: 6 x 60ml R4: 1 x 5ml	15-25°C
Magnesium	Arsenazo	Mono LS	BXC0353A	R1: 2 x 60ml R4: 1 x 5ml	15-25°C
Methanol in Body Fluid	UV	LS	BXC0495A	R1: 3 x 20ml R2: 2 x 5ml R4: 1 x 2ml	2-8°C
Microalbumin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0471A	R1: 1 x 50ml R2: 1 x 10ml R4: 5 x 1ml(C1-C5)	2-8°C
Myoglobin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0485A	R1: 1 x 20ml R4: 1 x 5ml	2-8°C
NEFA	UV Colorimetric	LS	BXC0477A	R1: 1 x 40 ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
NEFA	UV Colorimetric	LS	BXC0477B	R1: 2 x 40 ml R2: 1 x 20ml R4: 1 x 1ml	2-8°C
Oxalate (Urinary)	Enzymatic Colorimetric	LS	BXC0129A	R1: 1x10ml R2: 1x1ml R3: 1x20ml R4: 1x5ml Controls: 2x3ml (Charcoal 1 Vial)	2-8°C
Phosphorus (Inorganic)	Molybdate	LS	BXC0301A	R1: 2 x 30ml R2: 2 x 70ml R4: 1 x 5ml	15-25°C
Phosphorus (Inorganic)	Molybdate	Mono LS	BXC0302A	R1: 5 x 20ml R4: 1 x 5ml	15-25°C
Potassium	TPB (Colorimetric)	LS	BXC0132A	R1: 1 x 30ml R2: 1 x 30ml R3:1 x 60ml R4: 1 x 5ml	2-8°C
Potassium	Enzymatic	LS	BXC0135A	R1: 4 x 20ml R2: 2 x 10ml R4: 2 x 5ml	2-8°C
Potassium	TPB (Colorimetric)	Mono LS	BXC0138A	R1: 5 x 20ml R4: 1 x 5ml	15-25°C
Potassium	TPB (Colorimetric)	Mono LS	BXC0138B	R1: 10 x 20ml R4: 1 x 5ml	15-25°C
Pre-Albumin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0449A	R1: 2 x 20ml R4: 1 x 8ml	2-8°C
Procalcitonin	Turbilatex	LS	BXC0590A	R1: 1 x 30ml R2: 1 x 10ml	2-8°C
Renal Calculi (Kidney Stone Analysis)		LS	BXC0951A	100T	Room Temp
RF	Turbidimetric, Multi-point	LS	BXC0611A	R1: 1 x 40ml R2: 1 x 10ml R4: 1 x 2ml	2-8°C
RF	Turbidimetric, Multi-point	LS	BXC0611B	R1: 2 x 40ml R2: 2 x 10ml R4: 1 x 2ml	2-8°C
RF	Turbidimetric, Multi-point	LS	BXC0611C	R1: 4 x 40ml R2: 4 x 10ml R4: 1 x 2ml	2-8°C
Sodium	Enzymatic	LS	BXC0142A	R1: 2 x 20ml R2: 2 x 10ml R4: 2 x 5ml	2-8°C
Sodium	Colorimetric, Enzymatic	Mono LS	BXC0146A	R1: 5 x 20ml R4: 2 x 5ml	2-8°C
Superoxide Dismutase	Xanthine	Lyo	BXC0531A	R1: 5 x 20ml R2: 1 x 100ml R3:3 x 10ml R4: 5 x 10ml R6:2 x 125ml	2-8°C
ТІВС	Ion Saturation Method	LS	BXC0233A	R1: 1 x 100ml R2: 1 x 20g	Room Temp
ТІВС	DIRECT	LS	BXC0237A	R1: 1 x 50ml R2: 1 x 15ml	2-8°C
Total Antioxidant Status (TAS)	Enzymatic Colorimetric	LS	BXC0553A	R1: 1 x 50ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Total Lipid (Colorimetric)	Sulpho Phospho Vanillin	LS	BXC0263A	R1: 2 x 100ml R4: 1 x 10ml	Room Temp
Total Protein	Biuret (Concentrate)	LS	BXC0171A	R1: 4 x 125ml R2: 1 x 50ml R4: 1 x 5ml	Room Temp
Total Protein	Biuret (Ready To Use, Monoliquid)	Mono LS	BXC0173A	R1: 2 x 60ml R4: 1 x 5ml	Room Temp
Total Protein	Biuret (Ready To Use, Monoliquid)	Mono LS	BXC0173B	R1: 6 x 60ml R4: 1 x 5ml	Room Temp
Transferrin	Immunoturbidimetric (Without Sample Pre-dilution)	LS	BXC0741A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C

Clinical Chemistry | SECTION 1

Description	Method	Format	Cat No.	Size	Storage
Triglycerides	GPO-PAP	Lyo	BXC0272A	R1: 1 x 105ml R2: 10 x 10ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Lyo	BXC0272B	R1: 2 x 105ml R2: 10 x 20ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Lyo	BXC0272C	R1: 4 x 50ml R2: 4 x 50ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	BXC0271A	R1: 2 x 60ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	BXC0271B	R1: 4 x 60ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	BXC0271C	R1: 5 x 20ml R4: 1 x 5ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	BXC0271D	R1: 12 x 60ml R4: 1 x 5ml	2-8°C
Troponin-I	Turbidimetric	LS	BXC0469A	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
Troponin-I	Turbidimetric	LS	BXC0469B	R1: 1 x 80ml R2: 1 x 20ml	2-8°C
UIBC	Direct		BXC0220A	R1: 1 x 50ml R2: 1 x 10ml R3: 1 Vial	Room Temp
Urea	Modified Berthelot	LS	BXC0122A	R1: 2 x 125ml R2: 1 x 50ml R3:2 x 6.5ml R4: 1 x 5ml	2-8°C
Urea	Modified Berthelot	LS	BXC0122B	R1: 4 x 125ml R2: 2 x 50ml R3:4 x 6.5ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint	LS	BXC0123A	R1: 1 x 100ml R2: 1 x 20ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint	LS	BXC0123B	R1: 3 x 100ml R2: 1 x 60ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint	LS	BXC0123C	R1: 5 x 100ml R2: 1 x 100ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint (DGKC) 4+1	LS	BXC0124A	R1: 1 x 100ml R2: 1 x 25ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint	Lyo	BXC0121A	R1: 1 x 105ml R2: 5 x 20ml R4: 1 x 5ml	2-8°C
Urea	GLDH Kinetic & Endpoint	Lyo	BXC0121B	R1: 2 x 105ml R2: 10 x 20ml R4: 1 x 5ml	2-8°C
Urea	GLDH	Mono LS	BXC0126A	R1: 1 x 100ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	LS	BXC0602A	R1: 4 x 50ml R2: 1 x 50ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	LS	BXC0602B	R1: 4 x 20ml R2: 1 x 20ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	Lyo	BXC0601A	R1: 1 x 105ml R2: 5 x 20ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	Lyo	BXC0601B	R1: 2 x 105ml R2: 10 x 20ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	BXC0603A	R1: 5 x 20ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	BXC0603B	R1: 1 x 100ml R4: 1 x 5ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	BXC0603C	R1: 5 x 100ml R4: 1 x 5ml	2-8°C
Urinary & CSF Protein	Benzethonium Chloride	LS	BXC0174A	R1: 1 x 50ml R2: 1 x 10ml R4: 1 x 5ml	2-8°C
Urinary & CSF Protein	Pyrogallol Red	Mono LS	BXC0172A	R1: 2 x 60ml R4: 1 x 5ml	Room Temp
Urinary & CSF Protein	Pyrogallol Red	Mono LS	BXC0172B	R1: 4 x 60ml R4: 1 x 5ml	Room Temp
Vitamin-D	Turbidimetric	LS	BXC0742A	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
Vitamin-D	Turbidimetric	LS	BXC0742L	R1: 1 x 800ml R2: 1 x 200ml	2-8°C
Vitamin-D Calibrator Set		LS	BXC0475A	5x1ml	2-8°C
Zinc	5-Bromo-PAPS (Colorimetric, with deproteinisation)	LS	BXC0461A	R1: 2 x 50ml R2: 1 x 25ml R3: 1 x 25ml R4: 1 x 10ml	2-8°C
Zinc	5-Bromo-PAPS (Colorimetric, without deproteinisation)	Mono LS	BXC0462A	R1: 2 x 40ml R4: 1 x 5ml	2-8°C

Drugs of Abuse & Therapeutic Reagents

Drugs of abuse and therapeutic reagents in diagnostics are crucial for identifying substance use and managing therapeutic interventions. Drugs of abuse, including opioids, stimulants, and alcohol, can significantly impact health and require accurate detection for effective treatment and monitoring.

Diagnostic tests employ specialised reagents to effectively screen for drugs of abuse in biological samples, such as urine or blood. This process is essential not only in clinical environments, where immediate health assessments are critical, but also in rehabilitation programs that monitor patients' substance use and recovery progress. Conversely, therapeutic reagents are crucial for managing medication therapies, as they allow healthcare providers to monitor drug levels in patients. This ensures that patients receive the appropriate dosages, thereby preventing potential toxicity and optimizing treatment effectiveness.

Collectively, these diagnostic tools empower healthcare professionals to make well-informed decisions, promote safe prescribing practices, and provide comprehensive support for patients throughout their recovery journeys, ultimately enhancing overall health outcomes.

Ordering Information

Description	Method	Cat No.	Size	Storage
Barbiturates	Turbilatex	BXC0992A	R1:2x16.9ml; R2:2x8ml	2-8°C
Benzodiazepines	Turbilatex	BXC0993A	R1:2x16.9ml; R2:2x8ml	2-8°C
Cannabinoids	Turbilatex	BXC0994A	R1:2x16.9ml; R2:2x8ml	2-8°C
Carbamazepine	Turbilatex	BXC0996A	R1:2x16.9ml; R2:2x8ml	2-8°C
Cocaine Metabolite	Turbilatex	BXC0997A	R1:2x16.9ml; R2:2x8ml	2-8°C
Digoxin	Turbilatex	BXC0998A	R1:2x8ml; R2:2x6ml	2-8°C
Ecstasy	Turbilatex	BXC0999A	R1:2x16.9ml; R2:2x8ml	2-8°C
EDDP	Turbilatex	BXC0893A	R1:2x16.9ml; R2:2x8ml	2-8°C
Gentamicin	Turbilatex	BXC0895A	R1:2x16.9ml; R2:2x8ml	2-8°C
Methadone	Turbilatex	BXC0896A	R1:2x16.9ml; R2:2x8ml	2-8°C
Opiates	Turbilatex	BXC0897A	R1:2x16.9ml; R2:2x8ml	2-8°C
Phenobarbital	Turbilatex	BXC0898A	R1:2x17ml; R2:2x6ml	2-8°C
Valproic Acid	Turbilatex	BXC0899A	R1:2x15ml; R2:2x6ml	2-8°C



Analyser Specific Reagents

Analyser-specific reagents are formulated to be compatible with specific types or brands of clinical chemistry analysers. These reagents ensure accuracy and consistency in test results, allowing for reliable detection and quantification of various analytes, such as enzymes, hormones, and drugs.

Specifically formulated for each analyser, these reagents enhance the efficiency of diagnostic processes, reduce the likelihood of errors, and improve the overall quality of patient care. Their importance extends to standardization, as they help maintain comparability of results across different laboratories and testing environments. Ultimately, analyser-specific reagents are essential for enabling precise diagnostics, supporting effective treatment decisions, and enhancing patient outcomes in healthcare settings.

Fortress Diagnostics manufacturers, develops and supplies an extensive range of high quality analyser specific clinical chemistry reagents. Our reagents are manufactured to the highest quality standards in an ISO 13485:2016–certified laboratory, ensuring excellent stability, accuracy and results you can trust.

Compatible Systems

- Hitachi: 704, 717, 902, 904, 911, 912 & 917
- Monarch: 240+, 400, 480, 680, 1200 & 1600
- Olympus: AU400, AU600, AU640, AU680 & AU2700
- Beckman: DxC 700
- Furuno: CA180, CA270 & CA400

Hitachi System Reagents

Suitable for Hitachi 704, 717, 902, 904, 911 & 912

Description	Method	Format	Cat No.	Size	Storage
Albumin RTU	BCG	LS	HIT0222A	R1: 18 x 50ml	2-8°C
Albumin RTU	BCG	LS	HIT0222C	R1: 5 x 50ml	2-8°C
Albumin RTU	BCG	LS	HIT0222D	R1: 10 x 100ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	HIT0184A	R1: 12 x 50ml R2: 3 x 44ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	HIT0184D	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	HIT0185A	R1: 12 x 50ml R2: 3 x 44ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	HIT0185D	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	LS	HIT0190A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Alpha-1-Microglobulin	Immunoturbidimetry without sample pre-dilution	LS	HIT0241A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
ALT	IFCC	LS	HIT0213A	R1: 12 x 50ml R2: 3 x 44ml	2-8°C
ALT	IFCC	LS	HIT0213E	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Ammonia	GLDH, Enzymatic, UV	LS	HIT0372B	R1: 5 x 10ml R2: 1 x 10ml	2-8°C
Amylase	EPSG7	LS	HIT0563A	R1: 19 x 22ml R2: 4 x 21ml	2-8°C
Amylase	EPSG7	LS	HIT0563C	R1: 2 x 22ml R2: 1 x 9ml	2-8°C
Anti-CCP	Turbilatex	LS	HIT0379A	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	HIT0379B	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml	2-8°C
Apolipoprotein A1	Immunoturbidimetry without sample pre-dilution	LS	HIT0411A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Apolipoprotein B	Immunoturbidimetry without sample pre-dilution	LS	HIT0412A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
ASO	Turbidimetric	LS	HIT0501D	R1: 2 x 20ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
ACT	IFCC	LS	HIT0203A	R1: 12 x 50ml R2: 3 x	2-8°C
AST				44ml	

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
Bilirubin (Direct)	Jendrassik-Grof	LS	HIT0191A	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Bilirubin (Total)	Jendrassik-Grof	LS	HIT0192A	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
С3	Immunoturbidimetry without sample pre-dilution	LS	HIT0851A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
C4	Immunoturbidimetry without sample pre-dilution	LS	HIT0861A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Calcium	Arsenazo	LS	HIT0292C	R1: 3 x 50ml	2-8°C
Calcium	CPC	LS	HIT0291A	R1: 12 x 50ml R2: 6 x 43ml	2-8°C
Calcium	CPC	LS	HIT0291B	R1: 6 x 50ml R2: 3 x 43ml	2-8°C
Chloride	Thiocyanate	LS	HIT0281B	R1: 3 x 50ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	HIT0261H	R1: 12 x 50ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	HIT02611	R1: 10 x 100ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	HIT0261J	R1: 5 x 100ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	HIT0801C	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
СК	NAC-Activated	LS	HIT0252A	R1: 7 x 23ml R2: 4 x 8.4ml	2-8°C
СК	NAC-Activated	LS	HIT0252D	R1: 2 x 21ml R2: 1 x 10ml	2-8°C
СК-МВ	Immunoinhibition	LS	HIT0452A	R1: 6 x 21ml R2: 2 x 14.6ml	2-8°C
СК-МВ	Immunoinhibition	LS	HIT0452D	R1: 2 x 21ml R2: 1 x 10ml	2-8°C
CO2	Enzymatic	LS	HIT0152C	R1: 6 x 20ml	2-8°C
Creatinine	Jaffe	LS	HIT0111A	R1: 12 x 50ml R2: 3 x 45ml	2-8°C
Creatinine	Jaffe	LS	HIT0111C	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Creatinine	Enzymatic, SOX	LS	HIT0113B	R1: 2 x 50ml R2: 1 x 50ml R4: 2 x 1ml	2-8°C
CRP	Immunoturbidimetric, (Linearity 280mg/L)	LS	HIT0384A	R1: 2 x 20ml R2: 1 x 8ml R4: 1 x 1ml(L6)	2-8°C
CRP (Ultra Sensitive)	Immunoturbidimetric, Multipoint	LS	HIT0383A	R1: 1 x 16ml R2: 1 x 4ml R4: 1 x 200ul	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	HIT0382D	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	HIT0382E	R1: 2 x 40ml R2: 2 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	HIT0382F	R1: 4 x 40ml R2: 4 x 10ml	2-8°C
Cystatin-C	Turbidimetric	LS	HIT0777A	R1: 2 x 20ml R2: 1 x 10ml	2-8°C
Detergent 33	For HITACHI™ Systems	LS	HIT33100	10 x 100ml	2-8°C
Fructosamine	NBT	LS	HIT0591B	R1: 2 x 6ml R2: 2 x 14ml	2-8°C
GGT	Carboxy	LS	HIT0362A	R1: 12 x 50ml R2: 6 x 22ml	2-8°C
GGT	Carboxy	LS	HIT0362E	R1: 2 x 50ml R2: 1 x 22ml	2-8°C
Glucose	GOD-PAP	Mono LS	HIT0101H	R1: 12 x 50ml	2-8°C
Glucose	GOD-PAP	Mono LS	HIT0101J	R2: 5 x 100ml	2-8°C
Glucose	Hexokinase	LS	HIT0103B	R1: 12 x 50 R2: 6 x 22ml	2-8°C
HbA1c	Enzymatic	LS	HIT0672C	R1: 1 x 21ml R2: 1 x 9ml R3:1 x 12.5ml R4: 1 x 45ml	2-8°C
HbA1c Modified Enzymatic	Modified Enzymatic	LS	HIT0670C	R1: 1 x 30ml R2: 1 x 10ml R3: 2 x 120ml	2-8°C
HDL Cholesterol	Direct	LS	HIT0421A	R1: 6 x 40ml R2: 4 x 20ml	2-8°C
HDL Cholesterol	Direct	LS	HIT0421G	R1: 6 x 20ml R2: 2 x 20ml	2-8°C
Homocysteine	Turbidimetric	LS	HIT0690A	R1: 2 x 16ml R2: 1 x 8ml	2-8°C

Description	Method	Format	Cat No.	Size	Storage
IgA	Immunoturbidimetry without sample pre-dilution	LS	HIT0701A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
IgG	Immunoturbidimetry without sample pre-dilution	LS	HIT0721A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
IgM	Immunoturbidimetry without sample pre-dilution	LS	HIT0731A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Iron	Ferrozine	LS	HIT0235B	R1: 12 x 50ml R2: 6 x 22ml	2-8°C
Iron	Ferrozine	LS	HIT0235C	R1: 6 x 50ml R2: 3 x 22ml	2-8°C
Kappa Light Chains	Immunoturbidimetry without sample pre-dilution	LS	HIT0331A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate	Enzymatic	LS	HIT0621B	R1: 1 x 100ml R2: 16 x 6ml	2-8°C
Lactate	Enzymatic	Mono LS	HIT0622A	R1: 5 x 20ml	2-8°C
Lactate Dehydrogenase (LD)	Pyruvate -> Lactate	LS	HIT0242A	R1: 12 x 20ml R2: 3 x 20ml	2-8°C
Lactate Dehydrogenase (LD)	Pyruvate -> Lactate	LS	HIT0242C	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate Dehydrogenase (LD)	Lactate -> Pyruvate	LS	HIT0243A	R1: 12 x 20ml R2: 3 x 20ml	2-8°C
Lactate Dehydrogenase (LD)	Lactate -> Pyruvate	LS	HIT0243C	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lambda Light Chains	Immunoturbidimetry without sample pre-dilution	LS	HIT0381A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
LDL Cholesterol	Direct	LS	HIT0431G	R1: 2 x 20ml R2: 1 x 14ml	2-8°C
Lipase	Colorimetric	LS	HIT0511B	R1: 2 x 10ml R2: 1 x 10ml	2-8°C
Lipoprotein (a)	Immunoturbidimetry without sample pre-dilution	LS	HIT0130A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Magnesium	Xylidyl Blue	LS	HIT0352A	R1: 6 x 20ml	2-8°C
Microalbumin	Immunoturbidimetry without sample pre-dilution	LS	HIT0471B	R1: 2 x 20ml R2: 1 x 8ml R4: 5x1ml	2-8°C
Multiclean Cell Wash Solution	For HITACHI™ 704, 902, 904, & 911	LS	HITWS002	2 LITRES	2-8°C
NaOH with Detergent	For HITACHI™ 912, & 917	LS	HITNAOHD	2 LITRES	2-8°C
Phosphorus Inorganic	Molybdate	LS	HIT0301A	R1: 12 x 50ml R2: 6 x 43ml	2-8°C
Phosphorus Inorganic	Molybdate	LS	HIT0301B	R1: 3 x 50ml R2: 2 x 35ml	2-8°C
RF	Turbidimetric	LS	HIT0611D	R1: 2 x 20ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
TIBC (Direct)	Colorimetric	LS	HIT0237B	R1: 6 x 10ml R2: 6 x 5ml	2-8°C
TIBC (Direct) Total Protein RTU	Colorimetric Biuret	LS LS	HIT0237B HIT0173A	R1: 6 x 10ml R2: 6 x 5ml R1: 3 x 40ml R2: 3 x 45ml	2-8°C 2-8°C
				R1: 3 x 40ml R2: 3 x	
Total Protein RTU	Biuret	LS	HIT0173A	R1: 3 x 40ml R2: 3 x 45ml	2-8°C
Total Protein RTU Total Protein RTU	Biuret Biuret Immunoturbidimetry without	LS LS	HIT0173A HIT0173B	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml	2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin	Biuret Biuret Immunoturbidimetry without sample pre-dilution	LS LS LS	HIT0173A HIT0173B HIT0741A	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml	2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP	LS LS LS Mono LS	HIT0173A HIT0173B HIT0741A HIT0271A	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml	2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP	LS LS Mono LS Mono LS	HIT0173A HIT0173B HIT0741A HIT0271A HIT0271C	R1: 3 × 40ml R2: 3 × 45ml R1: 6 × 50ml R1: 2 × 20ml R2: 1 × 8ml R1: 8 × 50ml R1: 18 × 50ml R1: 12 × 50ml R1: 5 × 100ml	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP	LS LS Mono LS Mono LS	HIT0173A HIT0173B HIT0741A HIT0271A HIT0271C HIT0271E	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml R1: 18 x 50ml R1: 12 x 50ml R1: 5 x 100ml R1: 6 x 82ml R2: 3 x 100ml	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP GPO-PAP	LS LS Mono LS Mono LS Mono LS	HIT0173A HIT0173B HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml R1: 18 x 50ml R1: 12 x 50ml R1: 5 x 100ml R1: 6 x 82ml R2: 3 x	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides Urea	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP GPO-PAP GPO-PAP GDD-PAP	LS LS Mono LS Mono LS Mono LS	HIT0173A HIT0173B HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml R1: 18 x 50ml R1: 12 x 50ml R1: 5 x 100ml R1: 6 x 82ml R2: 3 x 100ml R1: 6 x 40ml R2: 3 x	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides Urea	Biuret Biuret Immunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP GPO-PAP GPO-PAP GDDH Kinetic & End Point GLDH Kinetic & End Point	LS LS Mono LS Mono LS Mono LS LS	HIT0173A HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F HIT0123A HIT0123D	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml R1: 18 x 50ml R1: 12 x 50ml R1: 5 x 100ml R1: 6 x 82ml R2: 3 x 100ml R1: 6 x 40ml R2: 3 x 49ml R1: 12 x 50ml R2: 6 x	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides Urea Urea	Biuret Biuret Dimmunoturbidimetry without sample pre-dilution GPO-PAP GPO-PAP GPO-PAP GPO-PAP GLDH Kinetic & End Point GLDH Kinetic & End Point	LS LS Mono LS Mono LS Mono LS LS LS	HIT0173A HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F HIT0123A HIT0123D HIT0123D	R1: 3 x 40ml R2: 3 x 45ml R1: 6 x 50ml R1: 2 x 20ml R2: 1 x 8ml R1: 8 x 50ml R1: 18 x 50ml R1: 12 x 50ml R1: 5 x 100ml R1: 6 x 82ml R2: 3 x 100ml R1: 6 x 40ml R2: 3 x 49ml R1: 12 x 50ml R2: 6 x 22ml R1: 6 x 50ml R2: 3 x	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides Urea Urea Urea	Biuret Biuret Dimmunoturbidimetry without Sample pre-dilution GPO-PAP GPO-PAP GPO-PAP GPO-PAP GLDH Kinetic & End Point GLDH Kinetic & End Point Uricase PAP	LS LS Mono LS Mono LS Mono LS LS LS	HIT0173A HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F HIT0123A HIT0123D HIT0602A HIT0602C	R1: 3 × 40ml R2: 3 × 45ml R1: 6 × 50ml R1: 2 × 20ml R2: 1 × 8ml R1: 8 × 50ml R1: 18 × 50ml R1: 12 × 50ml R1: 6 × 82ml R2: 3 × 100ml R1: 6 × 40ml R2: 3 × 49ml R1: 12 × 50ml R2: 6 × 22ml	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C
Total Protein RTU Total Protein RTU Transferrin Triglycerides Triglycerides Triglycerides Urea Urea Uric Acid Uric Acid	BiuretBiuretBiuretCPO-PAPGPO-PAPGPO-PAPGPO-PAPGPO-PAPGLDH Kinetic & End PointGLDH Kinetic & End PointUricase PAPUricase PAPUricase PAP	 LS LS Mono LS Mono LS Mono LS LS LS Mono LS 	HIT0173A HIT0741A HIT0271A HIT0271C HIT0271E HIT0271F HIT0123A HIT0123A HIT0123D HIT0602A HIT0603D	R1: 3 × 40ml R2: 3 × R1: 6 × 50ml R1: 2 × 20ml R2: 1 × 8ml R1: 8 × 50ml R1: 18 × 50ml R1: 12 × 50ml R1: 5 × 100ml R1: 6 × 82ml R2: 3 × 100ml R1: 6 × 40ml R2: 3 × 2ml R1: 6 × 50ml R2: 6 × 2ml R1: 6 × 50ml R2: 3 × R1: 6 × 50ml R2: 3 ×	2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C 2-8°C

Hitachi System Reagents

Suitable for Hitachi 917

Description	Method	Format	Cat No.	Size	Storage
Albumin RTU	BCG	LS	HIT7222A	R1: 6 x 64ml R2: 6 x 16ml	2-8°C
Albumin RTU	BCG	LS	HIT7222B	R1: 6 x 64ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	HIT7184A	R1: 6 x 66ml R2: 6 x 16ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	HIT7185A	R1: 6 x 66ml R2: 6 x 16ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	LS	HIT7190A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
Alpha-1-Microglobulin	Immunoturbidimetry without sample pre-dilution	LS	HIT7241A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
ALT	IFCC	LS	HIT7213A	R1: 4 x 66ml R2: 4 x 16ml	2-8°C
Ammonia	GLDH, Enzymatic, UV	LS	HIT7376A	R1: 6 x 10ml	2-8°C
Amylase	EPSG7	LS	HIT7563A	R1: 6 x 66ml R2: 6 x 16ml	2-8°C
Anti-CCP	Turbilatex	LS	HIT7379A	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	HIT7379B	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml	2-8°C
Apolipoprotein A1	Immunoturbidimetry without sample pre-dilution	LS	HIT7411A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
Apolipoprotein B	Immunoturbidimetry without sample pre-dilution	LS	HIT7412A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
ASO	Turbidimetric	LS	HIT7501A	R1: 2 x 20ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
AST	IFCC	LS	HIT7203A	R1: 4 x 66ml R2: 4 x 16ml	2-8°C
Bilirubin, Direct		LS	HIT7191A	R1: 6 x 62ml R2: 6 x 19ml	2-8°C
Bilirubin, Total		LS	HIT7192A	R1: 6 x 62ml R2: 6 x 19ml	2-8°C
C3	Immunoturbidimetry without sample pre-dilution	LS	HIT7851A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
C4	Immunoturbidimetry without sample pre-dilution	LS	HIT7861A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
Calcium	CPC	LS	HIT7291A	R1: 6 x 63ml R2: 6 x 29ml	2-8°C
Chloride	Thiocyanate	LS	HIT7281A	R1: 6 x 66ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	HIT7280A	R1: 12 x 66ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	HIT7801A	R1: 6 x 66ml R2: 6 x 16ml	2-8°C
ск	NAC-Activated	LS	HIT7252A	R1: 6 x 60ml R2: 6 x 15ml	2-8°C
СК-МВ	Immunoinhibition	LS	HIT7452A	R1: 6 x 20ml R2: 6 x 5ml	2-8°C
CO2	Enzymatic	LS	HIT7152A	R1: 6 x 20ml	2-8°C
Creatinine	Jaffe	LS	HIT7111A	R1: 6 x 64ml R2: 6 x 35ml	2-8°C
Creatinine	Enzymatic, SOX	LS	HIT7113B	R1: 2 x 50ml R2: 1 x 50ml R4: 2 x 1ml	2-8°C
CRP	Immunoturbidimetry without sample pre-dilution	LS	HIT7382A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
Detergent 33	For HITACHI™ Systems	LS	HIT7310A	R1: 12 x 66ml	2-8°C
Fructosamine	NBT	LS	HIT7591A	R1: 2 x 14ml R2: 2 x 6ml	2-8°C
GGT	Carboxy	LS	HIT7362A	R1: 6 x 66ml R2: 6 x 16ml	2-8°C
Glucose	GOD-PAP	Mono LS	HIT7101A	R2: 12 x 66ml	2-8°C
HbA1c	Enzymatic	LS	HIT7672A	R1: 1 x 21ml R2: 1 x 9ml R3:1 x 12.5ml R4: 1 x 45ml	2-8°C
HDL Cholesterol	Direct	LS	HIT7421A	R1: 6 x 54ml R2: 6 x 20ml	2-8°C
IgA	Immunoturbidimetry without sample pre-dilution	LS	HIT7701A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C

igGimmunoperiodimetry without immunoperiodimetry without imm	Description	Method	Format	Cat No.	Size	Storage
jpMsample pre-dilutionLSHI /737ATomin24°CIronFerrozineLSHI /737ATomin24°CKappa Light ChainsImmunoturbidimetry withouts ample pre-dilutionLSHI 77331AR1: 2: X form R2: 1xx R2: X form R2: 1xx R2: X form R2: 1xx R2: X form R2: 1xx R2: X form R2: 1xx24°CLactateEnzymaticLyoHIT721AR1: 2: X form R2: 1xx R2: X form R2: 1xx R2: X form R2: 1xx R2: X form R2: 1xx24°CLactate Dehydrogenase (LD)Vyruxt => LactateLSHIT721AR1: 6: X form R2: 6: X R1: 6: X form R2: 6: X R2: 1xx24°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT721AR1: 6: X form R2: 6: X R1: 6: X form R2: 6: X24°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT731AR1: 2: X form R2: 6: X R2: 1xx24°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT731AR1: 2: X form R2: 1xx R2: 1xx24°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT731AR1: 2: X form R2: 1xx R2: 1xx24°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT731AR1: 2: X form R2: 1xx R2: 1xx24°CLactate Dehydrogenase (LD)ColorimetricLSHIT731AR1: 6: X form R2: 1xx R2: 1xx24°CLactate Dehydrogenase (LD)ColorimetricLSHIT731AR1: 6: X form R2: 1xx R3: 1xx24°CLactate Dehydrogenase (LD)Rimonoturbidimetry without ample pre-dilutionLSHIT731AR1: 6: X form	IgG		LS	HIT7721A		2-8°C
IronFerrozineISHTT23Afom include 24°CKappa Light ChainsImmunoturbidimetry without sample pre-dilutionISHTT731AR1:2 x 10ml R2: 1 x2.8°CLactateEnzymaticVorHTT721AR1:2 x 20ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Purvate > LactateISHTT721AR1: 6 x 20ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Lactate > PyrvateISHTT724AR1: 6 x 60ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Lactate > PyrvateISHTT734AR1: 6 x 60ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Lactate > PyrvateISHTT734AR1: 6 x 60ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Lactate > PyrvateISHTT734AR1: 6 x 60ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)Lactate > PyrvateISHTT734AR1: 6 x 20ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)DirectISHTT731AR1: 6 x 20ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)DirectISHT731AR1: 6 x 20ml R2: 6 x 20ml2.8°CLactate Dehydrogenase (LD)DirectISHT731AR1: 6 x 20ml R2: 6 x 20ml2.8°CLastate Dehydrogenase (LD)Sample pre-dilutionISHT731AR1: 6 x 20ml R2: 6 x 20ml2.8°CLastate Dehydrogenase (LD)For HTACH™ 912, 8917ISHT731AR1: 6 x 20ml R2: 6 x 20ml2.8°CNaGensuin Maine DereeditutionISHT731AR1: 6 x 20ml R2: 6 x 20ml	IgM	5	LS	HIT7731A		2-8°C
Kappa Ligne Chainssample pre-dilutionISFIT 731AJondZei CLactateExzymaticLyoHT7621AR1:2 x 60ml R2:6 x 20ml2-8°CLactate Dehydrogenase (LD)Pyruvate > LactateIsHT7742AR1:6 x 66ml R2:6 x2-8°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHT7743AR1:6 x 66ml R2:6 x2-8°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHT731AR1:6 x 66ml R2:6 x2-8°CLambda Light ChainsImmonoturbidimetry without sample pre-dilutionLSHT731AR1:6 x 20ml R2:6 x2-8°CLipseColorimetricLSHT731AR1:6 x 20ml R2:6 x2-8°CLipseImmonoturbidimetry without sample pre-dilutionLSHT731AR1:6 x 20ml R2:6 x2-8°CLipseColorimetricLSHT731AR1:6 x 20ml R2:6 x2-8°CMagnesiumSingle pre-dilutionLSHT731AR1:6 x 20ml R2:6 x2-8°CNobydateLSHT731AR1:6 x 20ml R2:6 x2-8°CMagnesiumS	Iron	Ferrozine	LS	HIT7235A		2-8°C
LactateEnzymaticMono LSHIT7622AR1:6 x 20ml2.8°CLactate Dehydrogenase (LD)Pyruvate → LactateLSHIT7242AR1:6 x 66ml R2: 6 x 16ml2.8°CLactate Dehydrogenase (LD)Lactate → PyruvateLSHIT7243AR1:6 x 66ml R2: 6 x 16ml2.8°CLambda Light ChainsImmunoturbidimetry without smple pre-dilutionLSHIT731AR1:6 x 20ml R2: 6 x 10ml2.8°CLD CholesterolDirectLSHIT713AR1:6 x 20ml R2: 6 x 10ml2.8°CLipporbein (a)DirectLSHIT713AR1:6 x 20ml R2: 6 x 10ml2.8°CMagnesiumXylidy BlueLSHIT713AR1:6 x 20ml R2: 6 x 10ml2.8°CMagnesiumSylidy BlueLSHIT713AR1:6 x 20ml R2: 6 x 10ml2.8°CMono LD bidimetry without smple pre-dilutionLSHIT713AR1: 6 x 20ml R2: 6 x 10ml2.8°CNoADH with DetergentGorimetricLSHIT713AR1: 6 x 20ml R2: 1 x 10ml2.8°CNoBelporter RTUMolydateLSHIT731AR1: 6 x 20ml R2: 1 x 31ml2.8°CRFTurbidimetricLSHIT713AR1: 6 x 20ml R2: 1 x 31ml2.8°CTasferrinSurget pre-dilutionLSHIT731AR1: 6 x 20ml R2: 1 x 31ml2.8°CTasferrinGorimetricLSHIT713AR1: 6 x 20ml R2: 1 x 31ml2.8°CTasferrinSurget pre-dilutionLSHIT713AR1: 6 x 20ml R2: 1 x 31ml2.8°CTasferrinSurg	Kappa Light Chains		LS	HIT7331A		2-8°C
Lactate Dehydrogenase (LD)Pyruvate > LactateLSHIT7242AR1:6 x 66ml R2: 6 x time2.8°CLactate Dehydrogenase (LD)Lactate > PyruvateLSHIT7243AR1:6 x 66ml R2: 6 x time2.8°CLambda Light ChainsImmunoturbidimetry without sample pre-dilutionLSHIT7381AR1:6 x 20ml R2: 6 x model2.8°CLD CholesterolDirectLSHIT7130AR1:6 x 20ml R2: 6 x model2.8°CLipaseColorimetricLSHIT710AR1:6 x 20ml R2: 6 x model2.8°CMagnesiumXyldy BlueLSHIT7130AR1:5 x 20ml R2: 6 x model2.8°CMicroalbuminImmunoturbidimetry without sample pre-dilutionLSHIT7130AR1:2 x 16ml R2: 1 x model2.8°CNoAd with DetergentFor HITACHI™912, & 917LSHIT7130AR1:2 x 16ml R2: 1 x model2.8°CNaOH with DetergentFor HITACHI™912, & 917LSHIT771AR1: 2 x 20ml R2: 1 x model2.8°CRFUrbidimetricLSHIT731AR1: 6 x 63ml R2: 6 x model2.8°CTBG (Direct)ColorimetricLSHIT731AR1: 6 x 63ml R2: 6 x model2.8°CTasferrinSimple pre-dilutionLSHIT713AR1: 6 x 60ml R2: 6 x model2.8°CTasferrinGlorimetricLSHIT731AR1: 6 x 20ml R2: 6 x model2.8°CTasferrinSimple pre-dilutionLSHIT713AR1: 6 x 60ml R2: 6 x model2.8°CTasferrinSimple pre-dilutionLSHIT713AR1: 6 x 60ml R2: 6 x model2.8	Lactate	Enzymatic	Lyo	HIT7621A	R1: 2 x 60ml R2: 6 x 20ml	2-8°C
Lackate Dehydrogenase (LD)Pyrivate > LackateLSH17/22AIomitCell2.8°CLackate Dehydrogenase (LD)Lackate > PyruvateLSH177243ARine (A some R2:6 som	Lactate	Enzymatic	Mono LS	HIT7622A	R1: 6 x 20ml	2-8°C
Lactate Dehydrogenase (LD)Lactate > PyruvateLSH17/2434HomHom24°CLambda Light ChainsIsmmunoturbidimetry without sample pre-dilutionLSH17731AR1:2 x 10ml R2:1 x S1ml R2:1 x S1ml R2:1 x S1ml R2:1 x S2ml R2:6 x S2ml R2:6 x S2ml R2:6 xS2ml R2:2 x S2ml R2:6 x S2ml R2:	Lactate Dehydrogenase (LD)	Pyruvate -> Lactate	LS	HIT7242A		2-8°C
Lambda Light Chainssample pre-dilutionLSHIT731AJomt2-8 CLDL CholesterolDirectLSHIT7431AR1:6 x 20ml R2:6 x 8ml2-8°CLipaseColorimetricLSHIT7511AR1:6 x 20ml R2:6 x2-8°CMagnesiumSample pre-dilutionLSHIT7130AR1:6 x 20ml R2:6 x2-8°CMicroalbuminSample pre-dilutionLSHIT7130AR1:6 x 20ml R2:6 x2-8°CNaGroshburinSample pre-dilutionLSHIT7130AR1:6 x 20ml R2:6 x2-8°CNaGroshburinSample pre-dilutionLSHIT7130AR1:6 x 20ml R2:6 x2-8°CNaGroshburinImmunoturbidimetry without sample pre-dilutionLSHIT741AR1:2 x 16ml R2:1 x 10ml s1:1 x 16ml R2:1 x 10ml2-8°CNaGh with DetergentMolydateLSHIT7130AR1:6 x 63ml R2:6 x2-8°CRFColorimetricLSHIT7130AR1:6 x 20ml R2:1 x 10ml s1ml2-8°CTIBC (Direct)ColorimetricLSHIT713AR1:6 x 20ml R2:1 x 10ml s1ml2-8°CTransferrinSinveLSHIT713AR1:6 x 60ml R2:6 x2-8°CTiglyceridesGPO-PAPMono LSHIT713AR1:1 2 x 50ml s1ml2-8°CUrea Liquid StableGLD K Interic & End PointLSHIT713AR1:1 2 x 50ml s1ml2-8°CUric AcidUricse PAPKonLSHIT713AR1:2 x 50ml R2:6 x3-8°CLipus Lipus Li	Lactate Dehydrogenase (LD)	Lactate -> Pyruvate	LS	HIT7243A		2-8°C
LipaseColorimetricLSHIT7511AR1:6 x 20m R2: 6 x 3ml2.8°CLipoprotein (a)Immunoturbidimetry without sample predilutionLSHIT7130AR1: 2 x 16m R2: 1 x 0ml2.8°CMagnesiumXylidy BlueLSHIT7352AR1: 6 x 20ml2.8°CMicroalbuminSmunnoturbidimetry without sample predilutionLSHIT731AR1: 6 x 20ml2.8°CNaOH with DetergentFor HITACH™912, 8.917LSHIT701AR1: 2 x 16m R2: 1 x10m2.9°CNaOH with DetergentFor HITACH™912, 8.917LSHIT701AR1: 6 x 63ml R2: 6 x 3ml2.9°CRFUrbidimetry without sample pre-dilutionLSHIT731AR1: 6 x 63ml R2: 6 x 3ml2.9°CTBC (Direct)ColorimetricLSHIT731AR1: 6 x 63ml R2: 6 x 3ml2.9°CTatal Protein RTUGolorimetricLSHIT731AR1: 6 x 63ml R2: 6 x 3ml2.9°CTatal Protein RTUGolorimetricLSHIT731AR1: 6 x 63ml R2: 6 x 3ml2.9°CTatal Protein RTUBiuretLSHIT731AR1: 6 x 60ml R2: 6 x 3ml2.9°CTatal Protein RTUGolorimetric Without sample pre-dilutionLSHIT731AR1: 2 x 16ml R2: 1 x 3ml2.9°CTatal Protein RTUGolorimetric Kend PointLSHIT731AR1: 2 x 16ml R2: 1 x 3ml2.9°CTatal Protein RTUGolorimetric Kend PointLSHIT731AR1: 2 x 50ml R2: 6 x 3ml2.9°CTatal Quigit StableGOLP AMono LSHIT73	Lambda Light Chains		LS	HIT7381A		2-8°C
LipaseColonmetricLSHIT/511A13ml2.8°CLipoprotein (a)Immunoturbidimetry without sample pre-dilutionLSHIT7130AR1: 2 × 16ml R2: 1 × 10ml2.8°CMagnesiumXylidy BleLSHIT7352AR1: 6 × 20ml2.8°CMicroalbuminImmunoturbidimetry without sample pre-dilutionLSHIT741AR1: 2 × 16ml R2: 1 × 10ml2.8°CNaOH with DetergentFor HITACH™ 912, & 917LSHIT741AR1: 2 × 16ml R2: 1 × 10ml2.8°CPhosphorus InorganicMolydateLSHIT7611AR1: 2 × 20ml R2: 1 × 10ml2.8°CTIBC (Direct)ColorimetricLSHIT7611AR1: 2 × 20ml R2: 1 × 10ml2.8°CTotal Protein RTUBiuretLSHIT7730AR1: 6 × 60ml R2: 6 × 032.8°CTransferrinSimule pre-dilutionLSHIT771AR1: 2 × 10ml R2: 1 × 10ml2.8°CTriglyceridesGDP APMono LSHIT771AR1: 2 × 10ml R2: 1 × 10ml2.8°CUric AcidUricase PAPLSHIT7612AR1: 6 × 60ml R2: 6 × 32.8°CUric AcidUricase PAPKono LSHIT7602AR1: 6 × 60ml R2: 6 × 32.8°C	LDL Cholesterol	Direct	LS	HIT7431A	R1: 6 x 20ml R2: 6 x 8ml	2-8°C
Lipoprotein (a)sample pre-dilutionLSH17130A10ml28 CMagnesiumXylidy BlueLSH17352AR1: 6 x 20ml2.8°CMicroalbuminImmunoturbidimetry without sample pre-dilutionLSH17747AR1: 2 x 16ml R2: 1 x 10ml2.8°CNaOH with DetergentFor H1TACH™912, & 917LSH17747AZLTRES2.8°CPhosphorus InorganicMolybdateLSH17730AR1: 6 x 63ml R2: 6 x 31ml2.8°CRFTurbidimetricLSH17737AR1: 6 x 20ml R2: 1 x 10ml2.8°CTotal Protein RTUColorimetricLSH17737AR1: 6 x 20ml R2: 6 x 8ml2.8°CTansferrinBineuLSH17717AR1: 6 x 60ml R2: 6 x 8ml2.8°CTiglyceridesGD-PAPMono LSH1771AAR1: 2 x 50ml R2: 1 x 8ml2.8°CUrca Liquid StableGD-PAPKono LSH1771ABR1: 2 x 50ml R2: 6 x 8ml2.8°CUrcackidUrcase PAPLSH1771ABR1: 2 x 50ml R2: 6 x 8ml2.8°CUrcackidUrcase PAPLSH1771ABR1: 6 x 66ml R2: 6 x 8ml2.8°C	Lipase	Colorimetric	LS	HIT7511A		2-8°C
MicroalbuminImmunoturbidimetry without sample pre-dilutionLSHIT7471AR1: 2 x 16ml R2: 1 x 10ml2-8°CNaOH with DetergentFor HITACHI™ 912, & 917LSHIT7NAOA2 LITRES2-8°CPhosphorus InorganicMolybdateLSHIT7301AR1: 6 x 63ml R2: 6 x 31ml2-8°CRFTurbidimetricLSHIT7611AR1: 2 x 20ml R2: 1 x 10ml R4: 1 x 1ml2-8°CTIBC (Direct)ColorimetricLSHIT7611AR1: 6 x 20ml R2: 6 x 8ml 2-8°C2-8°CTotal Protein RTUBiuretLSHIT7173AR1: 6 x 66ml2-8°CTransferrinImmunoturbidimetry without sample pre-dilutionLSHIT771AR1: 2 x 16ml R2: 1 x 2-8°C2-8°CUrea Liquid StableGDD-PAPMone LSHIT713AR1: 2 x 16ml R2: 6 x R1: 6 x 66ml R2: 6 x 3ml2-8°CUric AcidUricase PAPKone LSHIT762AR1: 6 x 66ml R2: 6 x 16ml2-8°C	Lipoprotein (a)		LS	HIT7130A		2-8°C
Microatburninsample pre-dilutionLSHIT/4/TAR1: 2 × 16mt R2: 1 × 10mt2-8°CNaOH with DetergentFor HITACHI™ 912, & 917LSHIT7NAOA2 LITRES2-8°CPhosphorus InorganicMolybdateLSHIT7301AR1: 6 × 63ml R2: 6 × 31ml2-8°CRFTurbidimetricLSHIT7611AR1: 2 × 20ml R2: 1 × 10ml2-8°CTIBC (Direct)ColorimetricLSHIT713AR1: 6 × 20ml R2: 6 × 81ml2-8°CTotal Protein RTUBiuretLSHIT713AR1: 6 × 60ml R2: 6 × 81ml2-8°CTransferrinImmunoturbidimetry without sample pre-dilutionLSHIT771AR1: 2 × 16ml R2: 1 × 10ml 10ml2-8°CUrea Liquid StableGDO-PAPMono LSHIT721AR1: 1 × 65ml2-8°CUric AcidUricase PAPLSHIT7602AR1: 6 × 66ml R2: 6 × 3ml 6fml R2: 6 × 6fml2-8°CUric AcidUricase PAPMono LSHIT7603AR1: 6 × 66ml R2: 6 × 3ml 6fml R2: 6 × 6fml2-8°C	Magnesium	Xylidyl Blue	LS	HIT7352A	R1: 6 x 20ml	2-8°C
Phosphorus InorganicMolybateLSHIT7301AR1: 6 × 63ml R2: 6 x 31ml2.8°CRFTurbidimetricLSHIT7611AR1: 2 × 20ml R2: 1 × 10ml 4: 1 × 1ml2.8°CTIBC (Direct)ColorimetricLSHIT7237AR1: 6 × 20ml R2: 6 × 8ml 4: 1 × 1ml2.8°CTotal Protein RTUBiuretLSHIT713AR1: 6 × 60ml R2: 6 × 8ml 4: 1 × 1ml2.8°CTransferrinMonuoturbidimetry without sample pre-dilutionLSHIT771AR1: 6 × 60ml R2: 6 × 8ml 4: 1 × 10ml2.8°CUrea Liquid StableGDP-PAPMonu LSHIT721AR1: 2 × 10ml R2: 1 × 8ml 43ml2.8°CUric AcidUricase PAPLSHIT701AR1: 6 × 66ml R2: 6 × 3ml 43ml2.8°CUric AcidUricase PAPLSHIT701AR1: 6 × 66ml R2: 6 × 3ml 43ml2.8°C	Microalbumin	5	LS	HIT7471A	R1: 2 x 16ml R2: 1 x 10ml	2-8°C
Phosphorus horganicMolybateLSHI 7301AJaml24°CRFTurbidmetricLSHIT711AR1:2×20mR2:1×10ml2-8°CTIBC (Direct)ColorimetricLSHIT7237AR1:6×20mR2:6×8ml2-8°CTotal Protein RTUBiuretLSHIT713AR1:6×66ml R2:6×8ml2-8°CTransferrinSample pre-dilutionLSHIT71AAR1:2×16ml R2:1×x2-8°CTriglyceridesOPO-PAPMono LSHIT721AR1:1×65ml2-8°CUrea Liquid StableGLDH Kinetic & End PointLSHIT712AAR1:6×66ml R2:6×32-8°CUricse PAPLSK117602AR1:6×66ml R2:6×32-8°CUricse PAPLSKI17602AR1:6×66ml R2:6×32-8°CUricse PAPLSHIT701AR1:6×66ml R2:6×32-8°CUricse PAPKinetKinetKinet2-8°CMono LSHIT701AR1:6×66ml R2:6×32-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinet2-8°CLick AcidKinetKinetKinetKinetLick AcidKinetKinetKinetKinetLick AcidKinetKinetKinetKinetLick Acid <th>NaOH with Detergent</th> <th>For HITACHI™ 912, & 917</th> <th>LS</th> <th>HIT7NAOA</th> <th>2 LITRES</th> <th>2-8°C</th>	NaOH with Detergent	For HITACHI™ 912, & 917	LS	HIT7NAOA	2 LITRES	2-8°C
RFInfodimetricLSH1761AAR4:1 x 1ml2-8°CTIBC (Direct)ColorimetricLSH17237AR1:6 x 20ml R2:6 x 8ml2-8°CTotal Protein RTUBiuretLSH17713AR1:6 x 66ml2-8°CTransferrinImmunoturbidimetry without sample pre-dilutionLSH1771AAR1:2 x 16ml R2:1 x 10ml2-8°CTriglyceridesGPO-PAPMono LSH1771AAR1:1 x 65ml2-8°CUrea Liquid StableGLDH Kinetic & End PointLSH17712AAR1:6 x 66ml R2:6 x 43ml2-8°CUric AcidUricase PAPLSH17602AR1:6 x 66ml R2:6 x 6ml2-8°C	Phosphorus Inorganic	Molybdate	LS	HIT7301A		2-8°C
Total Protein RTUBiuretLSHTT713AR1: 6 x 66ml2-8°CTransferrinImmunoturbidimetry without sample pre-dilutionLSH1T771AR1: 2 x 16ml R2: 1 x 10ml2-8°CTriglyceridesGPO-PAPMono LSHIT721AR1: 1 x 65ml2-8°CUrea Liquid StableGLDH Kinetic & End PointLSHIT7123AR1: 6 x 66ml R2: 6 x 4ml2-8°CUric AcidUricase PAPLSHIT7602AR1: 6 x 66ml R2: 6 x 6ml2-8°CUric AcidUricase PAPLSHIT7603AR1: 6 x 66ml R2: 6 x 6ml2-8°C	RF	Turbidimetric	LS	HIT7611A		2-8°C
TransferinImmunoturbidimetry without sample pre-dilutionLSHT7741AR1: 2 x 16ml R2: 1 x 10ml2-8°CTriglyceridesGPO-PAPMono LSHT7271AR1: 1 2 x 65ml2-8°CUrea Liquid StableGLDH Kinetic & End PointLSHT7123AR1: 6 x 66ml R2: 6 x 3ml2-8°CUric AcidUricase PAPLSHT7602AR1: 6 x 66ml R2: 6 x 6ml2-8°CUric AcidUricase PAPKono LSHT7603AR1: 6 x 66ml R2: 6 x 6ml2-8°C	TIBC (Direct)	Colorimetric	LS	HIT7237A	R1: 6 x 20ml R2: 6 x 8ml	2-8°C
Transferrinsample pre-dilutionLSH1774TA10ml2-8 CTriglyceridesGPO-PAPMono LSH17271AR1:12 x 65ml2-8°CUrea Liquid StableGLDH Kinetic & End PointLSH177123AR1:6 x 66ml R2:6 x 43ml2-8°CUric AcidUricase PAPLSH177602AR1:6 x 66ml R2:6 x 16ml2-8°CUricase PAPMono LSH17603AR1:6 x 66ml R2:6 x 16ml2-8°C	Total Protein RTU	Biuret	LS	HIT7173A	R1: 6 x 66ml	2-8°C
Urea Liquid Stable GLDH Kinetic & End Point LS HIT7123A R1: 6 × 66ml R2: 6 × 43ml 2-8°C Uric Acid Uricase PAP LS HIT7602A R1: 6 × 66ml R2: 6 × 16ml 2-8°C Uric Acid Uricase PAP LS HIT7602A R1: 6 × 66ml R2: 6 × 16ml 2-8°C Uric Acid Uricase PAP Mono LS HIT7603A R1: 6 × 66ml 2-8°C	Transferrin	5	LS	HIT7741A		2-8°C
Uric Acid Uricase PAP LS HIT /123A 43ml 2-8°C Uric Acid Uricase PAP LS HIT 7602A R1: 6 x 66ml R2: 6 x 16ml 2-8°C Uric Acid Uricase PAP Mono LS HIT 7603A R1: 6 x 66ml 2-8°C	Triglycerides	GPO-PAP	Mono LS	HIT7271A	R1: 12 x 65ml	2-8°C
Uric Acid Uricase PAP LS HIT7602A 16ml 2-8°C Uric Acid Uricase PAP Mono LS HIT7603A R1: 6 x 66ml 2-8°C	Urea Liquid Stable	GLDH Kinetic & End Point	LS	HIT7123A		2-8°C
	Uric Acid	Uricase PAP	LS	HIT7602A		2-8°C
Urinary Protein Pyrogallol Red LS HIT7172A R1: 6 x 66ml 2-8°C	Uric Acid	Uricase PAP	Mono LS	HIT7603A	R1: 6 x 66ml	2-8°C
	Urinary Protein	Pyrogallol Red	LS	HIT7172A	R1: 6 x 66ml	2-8°C

Monarch System Reagents

Suitable for Monarch 240+, 400, 480, 680, 1200 & 1600

Description	Method	Format	Cat No.	Size	Storage
Anti-bacterial Phosphate Free Wash	For MONARCH Series	LS	MONAPFW1	4 x 70 ml	2-8°C
Albumin RTU	BCG	LS	MON0222A	R1: 2 x 60ml	2-8°C
Albumin RTU	BCG	LS	MON0222B	R1: 6 x 60ml	2-8°C
Albumin RTU	BCG	LS	MON0222C	R1: 5 x 50ml	2-8°C
Albumin RTU	BCG	LS	MON0222D	R1: 12 x 50ml	2-8°C
Aldolase	GDH/TIM (UV)	Lyo	MON0391A	R1: 5 x 20ml R2: 2 x 1ml R3:1 x 0.5ml	2-8°C
Alkaline Detergent	For MONARCH Systems	LS	MONDET1L	1 x 1000 ml	2-8°C
Alkaline Detergent	For MONARCH Systems	LS	MONDET6L	6 x 1000 ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	MON0184A	R1: 5 x 25ml R2: 1 x 25ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	MON0184B	R1: 5 x 50ml R2: 1 x 50ml	2-8°C
Alkaline Phosphatase	AMP (IFCC)	LS	MON0184C	R1: 10 x 50ml R2: 2 x 50ml	2-8°C

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
Alkaline Phosphatase	AMP (IFCC)	LS	MON0184D	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	MON0185A	R1: 5 x 25ml R2: 1 x 25ml	2-8°C
Alkaline Phosphatase	DEA (DGKC)	LS	MON0185B	R1: 5 x 50ml R2: 1 x 50ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	STOOL	MON0189A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	LS	MON0190A	R1: 1 x 20ml R2: 1 x 4ml	2-8°C
Alpha-1-Antitrypsin	Immunoturbidimetry without sample pre-dilution	LS	MON0190B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Alpha-1-Microglobulin	Immunoturbidimetry without sample pre-dilution	LS	MON0241A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
ALT	IFCC	LS	MON0213A	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
ALT	IFCC	LS	MON0213B	R1: 4 x 50ml R2: 2 x 20ml	2-8°C
ALT	IFCC	LS	MON0213C	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Ammonia	GLDH, Enzymatic, UV	LS	MON0376A	R1: 5 x 20ml	2-8°C
Amylase	EPSG7	LS	MON0563A	R1: 2 x 25 ml R2: 1 x 10 ml	2-8°C
Amylase	EPSG7	LS	MON0563B	R1: 2 x 50 ml R2: 1 x 20 ml	2-8°C
Anti-bacterial Phosphate Free Wash	For MONARCH Series	LS	MONAPFW2	2 x 500 ml	2-8°C
Anti-CCP	Turbilatex	LS	MON0379A	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	MON0379B	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml	2-8°C
Apolipoprotein A1	Immunoturbidimetry without sample pre-dilution	LS	MON0411A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Apolipoprotein B	Immunoturbidimetry without sample pre-dilution	LS	MON0412A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
ASO	Turbidimetric	LS	MON0501A	R1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
ASO	Turbidimetric	LS	MON0501B	R1: 2 x 40ml R2: 2 x 10ml R4: 1 x 1ml	2-8°C
ASO	Turbidimetric	LS	MON0501D	R1: 2 x 20ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
AST	IFCC	LS	MON0203A	R1: 1 x 50ml R2: 1 x 10ml	2-8°C
AST	IFCC	LS	MON0203D	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
AST	IFCC	LS	MON0203F	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Beta-2-Microglobulin	Immunoturbidimetric	LS	MON0090A	R1: 2 x 12.5ml R2: 1 x 6ml R4: 1x 0.5ml	2-8°C
Bilirubin (Direct)	Jendrassik Grof	LS	MON0191A	R1: 2 x 50ml R2: 1 x 25ml	2-8°C
Bilirubin (Direct)	Jendrassik Grof	LS	MON0191B	R1: 4 x 50ml R2: 1 x 50ml	2-8°C
Bilirubin (Direct)	Jendrassik Grof	LS	MON0191C	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Bilirubin (Total)	Jendrassik Grof	LS	MON0192A	R1: 2 x 50ml R2: 1 x 25ml	2-8°C
Bilirubin (Total)	Jendrassik Grof	LS	MON0192B	R1: 4 x 50ml R2: 1 x 50ml	2-8°C
Bilirubin (Total)	Jendrassik Grof	LS	MON0192C	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Bilirubin (Total)	Vanadate Oxidation	LS	MON0105A	R1: 2 x 70ml R2: 2 x 18ml	2-8°C
Bilirubin Direct	Vanadate Oxidation	LS	MON0104A	R1: 2 x 70ml R2: 2 x 18ml	2-8°C
С3	Immunoturbidimetry without sample pre-dilution	LS	MON0851A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
C4	Immunoturbidimetry without sample pre-dilution	LS	MON0861A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Calcium	Arsenazo	LS	MON0292A	R1: 2 x 60ml	2-8°C
Calcium	Arsenazo	LS	MON0292B	R1: 4 x 60ml	2-8°C
Calcium	Arsenazo	LS	MON0292C	R1: 3 x 50ml	2-8°C
Calcium	CPC	LS	MON0291B	R1: 6 x 50ml R2: 6 x 50ml	2-8°C

Description	Method	Format	Cat No.	Size	Storage
Chloride	Thiocyanate	LS	MON0281B	R1: 3 x 50ml	2-8°C
Chloride Electrode - PN:1110799019		LS	MON00010	Single Electrode	2-8°C
Cholesterol	CHOD-PAP	Mono LS	MON0261A	R1: 2 x 60ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	MON0261B	R1: 6 x 60ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	MON0261C	R1: 12 x 50ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	MON0801C	R1: 3 x 20ml R2: 1 x 13ml	2-8°C
ск	NAC-Activated	LS	MON0252A	R1: 2 x 25ml R2: 1 x 10ml	2-8°C
СК	NAC-Activated	LS	MON0252B	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
ск	NAC-Activated	LS	MON0252D	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
СК-МВ	Immunoinhibition	LS	MON0452C	R1: 1 x 50ml R2: 1 x 10ml	2-8°C
СК-МВ	Immunoinhibition	LS	MON0452D	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
CO2	Enzymatic	LS	MON0152C	R1: 6 x 20ml	2-8°C
Copper Urinary	DBP (Colorimetric)	Mono LS	MON0343A	R1: 2 x 25ml R4: 1 x 5ml	2-8°C
Creatinine	Jaffe	LS	MON0111A	R1: 6 x 50ml R2: 6 x 50ml	2-8°C
Creatinine	Jaffe	LS	MON0111B	R1: 3 x 50ml R2: 3 x 50ml	2-8°C
Creatinine	Enzymatic, SOX	LS	MON0113B	R1: 2 x 50ml R2: 1 x 50ml R4: 2 x 1ml	2-8°C
CRP	Immunoturbidimetric, (Linearity 280mg/L)	LS	MON0384A	R1: 2 x 20ml R2: 1 x 8ml R4: 1 x 1ml(L6)	2-8°C
CRP (Ultra Sensitive)	Immunoturbidimetric, Multipoint	LS	MON0383A	R1: 1 x 16ml R2: 1 x 4ml R4: 1 x 200ul	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	MON0382D	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	MON0382E	R1: 2 x 40ml R2: 2 x 10ml	2-8°C
CRP Full Range	Turbilatex Full Range (Linearity 150mg/L)	LS	MON0382F	R1: 4 x 40ml R2: 4 x 10ml	2-8°C
Cystatin-C	Turbidimetric	LS	MON0777A	R1: 2 x 20ml R2: 1 x 10ml	2-8°C
Fructosamine	NBT	LS	MON0591B	R1: 2 x 6ml R2: 2 x 14ml	2-8°C
G6PDH	Digitonin (UV)	LS	MON0571A	R1: 2 x 50ml R2: 1 x 2ml R3:1 x 20ml R4: 2 x 105ml	2-8°C
бст	Carboxy	LS	MON0362D	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
GGT	Carboxy	LS	MON0362E	R1: 4 x 50ml R2: 2 x 20ml	2-8°C
Glucose	GOD-PAP	Mono LS	MON0101A	R1: 6 x 50ml	2-8°C
Glucose	GOD-PAP	Mono LS	MON0101B	R1: 10 x 50ml	2-8°C
Glucose	GOD-PAP	Mono LS	MON0101C	R1: 12 x 50ml	2-8°C
Glucose	Hexokinase	LS	MON0103A	R1: 2 x 50 R2: 1 x 20ml	2-8°C
Glucose	Hexokinase	LS	MON0103B	R1: 5 x 50 R2: 1 x 50ml	2-8°C
HbA1c	Modified Enzymatic	LS	MON0670A	R1: 1 x 30ml R2: 1 x 10ml R3: 2 x 120ml; 2x0.5ml	2-8°C
HbA1c	Modified Enzymatic	LS	MON0670B	R1: 2 x 30ml R2: 1 x 20ml R3: 4 x 120ml; 2x0.5ml	2-8°C
HDL Cholesterol	Direct	LS	MON0421A	R1: 1 x 30ml R2: 1 x 10ml	2-8°C
HDL Cholesterol	Direct	LS	MON0421B	R1: 2 x 60ml R2: 2 x 20ml	2-8°C
HDL Cholesterol	Direct	LS	MON0421C	R1: 6 x 40ml R2: 4 x 20ml	2-8°C
Homocysteine	Turbidimetric	LS	MON0690A	R1: 2 x 16ml R2: 1 x 8ml	2-8°C
IgA	Immunoturbidimetry without sample pre-dilution	LS	MON0701A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
lgE	Immunoturbidimetric (Without Sample Pre-dilution)	LS	MON0751A	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
IgG	Immunoturbidimetry without sample pre-dilution	LS	MON0721A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
IgM	Immunoturbidimetry without sample pre-dilution	LS	MON0731A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Iron	Ferrozine	LS	MON0235A	R1: 2 x 40ml R2: 2 x 10ml	2-8°C
Iron	Ferrozine	LS	MON0235B	R1: 4 x 40ml R2: 2 x 20ml	2-8°C
Iron	Ferrozine	LS	MON0235C	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Iron	Ferrene	LS	MON0232A	R1 4 x 50ml R2 1 x 65ml R4: 1 x 5ml	2-8°C
ISE Blood Control High		Lyo	MON00014	1 x 4 ml Diluent; 1 x vial of Lyo	2-8°C
ISE Blood Control Low		Lyo	MON00013	1 x 4 ml Diluent; 1 x vial of Lyo	2-8°C
ISE Blood Sample Calibrator	For MONARCH Series	Lyo	MON00005	5 x 4 ml Diluent; 5 x vial of Lyo	2-8°C
ISE Detergent	For MONARCH Series	LS	MON00004	1 x 100ml	2-8°C
ISE Diluent	For MONARCH Series	LS	MON00001	1 x 2L	2-8°C
ISE Internal Standard Solution	For MONARCH Series	LS	MON00003	1 x 2L	2-8°C
ISE Reference Solution	For MONARCH Series	LS	MON00002	1 x 2L	2-8°C
ISE Standard Solution	For MONARCH Series	LS	MON00008	1 x 10ml Std Low; 1 x 10ml Std High	2-8°C
ISE Standard Solution High	For MONARCH Series	LS	MON00006	1 x 10ml	2-8°C
ISE Standard Solution Low	For MONARCH Series	LS	MON00007	1 x 10ml	2-8°C
Kappa Light Chains	Immunoturbidimetry without sample pre-dilution	LS	MON0331A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate	Enzymatic	LS	MON0621B	R1: 2 x 50ml R2: 16 x 6ml	2-8°C
Lactate	Enzymatic	Mono LS	MON0622A	R1: 5 x 20ml	2-8°C
Lactate	Enzymatic	Mono LS	MON0622B	R1: 2 x 50ml	2-8°C
Lactate Dehydrogenase (LD)	Pyruvate -> Lactate	LS	MON0242C	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate Dehydrogenase (LD)	Lactate -> Pyruvate	LS	MON0243C	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lambda Light Chains	Immunoturbidimetry without sample pre-dilution	LS	MON0381A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
LDL Cholesterol	Direct	LS	MON0431A	R1: 1 x 30ml R2: 1 x 10ml	2-8°C
LDL Cholesterol	Direct	LS	MON0431B	R1: 2 x 60ml R2: 2 x 20ml	2-8°C
LDL Cholesterol	Direct	LS	MON0431G	R1: 3 x 20ml R2: 2 x 10ml	2-8°C
Lipase	Colorimetric	LS	MON0511A	R1: 2 x 20ml R2: 1 x 20ml	2-8°C
Lipase	Colorimetric	LS	MON0511B	R1: 2 x 10ml R2: 1 x 10ml	2-8°C
Lipoprotein (a)	Immunoturbidimetry without sample pre-dilution	LS	MON0130A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Magnesium	Xylidyl Blue	LS	MON0352A	R1: 2 x 60ml	2-8°C
Magnesium	Xylidyl Blue	LS	MON0352B	R1: 6 x 60ml	2-8°C
Microalbumin	Immunoturbidimetry without sample pre-dilution	LS	MON0471A	R1: 1 x 50ml R2: 1 x 10ml R4: 5x1ml	2-8°C
Microalbumin	Immunoturbidimetry without sample pre-dilution	LS	MON0471B	R1: 2 x 20ml R2: 1 x 8ml R4: 5x1ml	2-8°C
Phosphorus Inorganic	Molybdate	LS	MON0302A	R1: 5 x 20ml	2-8°C
Phosphorus Inorganic	Molybdate	LS	MON0302B	R1: 4 x 50ml	2-8°C
		LS	MON0135A	R1: 3 x 20ml R2: 1 x 15ml	2-8°C
Potassium	Enzymatic	LJ		ISIII	
Potassium Potassium Electrode Body - PN:1111999039	Enzymatic	LS	MON00012	Single	2-8°C
Potassium Electrode Body -	Enzymatic				2-8°C 2-8°C

Description	Method	Format	Cat No.	Size	Storage
RF	Turbidimetric	LS	MON0611B	R1: 2 x 40ml R2: 1 x 20ml R4: 1 x 2ml	2-8°C
RF	Turbidimetric	LS	MON0611D	R1: 2 x 20ml R2: 1 x 10ml R4: 1 x 2ml	2-8°C
Sodium	Enzymatic	LS	MON0142A	R1: 2 x 20ml R2: 2 x 10ml R4: 2 x 5ml	2-8°C
Sodium Electrode – PN:1110799018		LS	MON00009	Single Electrode	2-8°C
TIBC (Direct)	Colorimetric	LS	MON0237B	R1: 3 x 20ml R2: 2x 10ml	2-8°C
Total Protein RTU	Biuret	LS	MON0173A	R1: 2 x 60ml	2-8°C
Total Protein RTU	Biuret	LS	MON0173B	R1: 6 x 60ml	2-8°C
Transferrin	Immunoturbidimetry without sample pre-dilution	LS	MON0741A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	MON0271A	R1: 2 x 60ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	MON0271B	R1: 4 x 60ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	MON0271C	R1: 6 x 50ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	MON0271D	R1: 12 x 50ml	2-8°C
Troponin-I	Turbidimetric	LS	MON0469A	R1: 1 x 40ml R2: 1 x 10ml	2-8°C
Urea	GLDH Kinetic & End Point	LS	MON0123A	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
Urea	GLDH Kinetic & End Point	LS	MON0123B	R1: 6 x 50ml R2: 3 x 20ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	MON0603B	R1: 2 x 50ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	MON0603C	R1: 10 x 50ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	MON0603D	R1: 6 x 50ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	MON0603A	R1: 5 x 20ml	2-8°C
Urinary Protein	Pyrogallol Red	LS	MON0172A	R1: 2 x 60ml	2-8°C
Urinary Protein	Pyrogallol Red	LS	MON0172B	R1: 4 x 60ml	2-8°C
Urinary Protein	Pyrogallol Red	LS	MON0172C	R1: 3 x 50ml	2-8°C
Zinc	Colorimetric	Mono LS	MON0462B	R1: 6 x 20ml	2-8°C

Olympus & Beckman Coulter System Reagents

Suitable for Olympus AU400, AU600, AU640, AU680 & AU2700 / Beckman Coulter DxC 700

Description	Method	Format	Cat No.	Size	Storage
Albumin RTU	BCG	LS	OLY0222A	R1: 2 x 60ml	2-8°C
Albumin RTU	BCG	LS	OLY0222B	R1: 6 x 60ml	2-8°C
Albumin RTU	BCG	LS	OLY0222D	R1: 4 x 60ml	2-8°C
ALP AMP IFCC	AMP	LS	OLY0184A	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
ALP AMP IFCC	AMP	LS	OLY0184B	R1: 5 x 60ml R2: 1 x 60ml	2-8°C
ALP DEA DGKC	DEA	LS	OLY0185A	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
ALP DEA DGKC	DEA	LS	OLY0185B	R1: 5 x 60ml R2: 1 x 60ml	2-8°C
ALT IFCC	IFCC	LS	OLY0213A	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
ALT IFCC	IFCC	LS	OLY0213B	R1: 5 x 60ml R2: 1 x 60ml	2-8°C
Ammonia	GLDH, Enzymatic, UV	LS	OLY0372B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Amylase EPSG7	EPSG7	LS	OLY0563A	R1: 3 x 30ml R2: 1 x 18ml	2-8°C
Amylase EPSG7	EPSG7	LS	OLY0563B	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
Amylase Pancreatic	EPSG7	LS	OLY0564B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Anti-CCP	Turbilatex	LS	OLY0379A	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	OLY0379B	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5ml	2-8°C
Anti-CCP **USA ONLY**	Turbilatex	LS	OLY0379A	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 1ml	2-8°C

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
Apolipoprotein A1	Immunoturbidimetry without sample pre-dilution	LS	OLY0411A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Apolipoprotein B	Immunoturbidimetry without sample pre-dilution	LS	OLY0412A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
ASO Immunoturbidimetry	Turbilatex	LS	OLY0501A	R1: 1 x 40ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
ASO Immunoturbidimetry	Turbilatex	LS	OLY0501B	R1: 2 x 40ml R2: 1 x 20ml R4: 1 x 1ml	2-8°C
ASO Immunoturbidimetry	Turbilatex	LS	OLY0501C	R1: 4 x 40ml R2: 2 x 20ml R4: 1 x 1ml	2-8°C
AST IFCC	IFCC	LS	OLY0203A	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
AST IFCC	IFCC	LS	OLY0203B	R1: 5 x 60ml R2: 1 x 60ml	2-8°C
Beta-2-Microglobulin	Turbilatex	LS	OLY0090A	R1: 2 x 12.5ml R2: 1 x 6ml	2-8°C
Bilirubin (Direct)	Vanadate Oxidation	LS	OLY0104A	R1: 2 x 70ml R2: 2 x 18ml	2-8°C
Bilirubin (Total)	Vanadate Oxidation	LS	OLY0105A	R1: 2 x 70ml R2: 2 x 18ml	2-8°C
Bilirubin (Direct)	Jendrassik GROF	LS	OLY0191A	R1: 2 x 60ml R2: 1 x 24ml	2-8°C
Bilirubin (Direct)	Jendrassik GROF	LS	OLY0191B	R1: 4 x 60ml R2: 2 x 24ml	2-8°C
Bilirubin (Total)	Jendrassik GROF	LS	OLY0192A	R1: 2 x 60ml R2: 1 x 24ml	2-8°C
Bilirubin (Total)	Jendrassik GROF	LS	OLY0192B	R1: 4 x 60ml R2: 2 x 24ml	2-8°C
Blood Alcohol	UV	LS	OLY0491A	R1: 3 x 20ml R2: 1 x 10ml R4: 1 x 2ml	2-8°C
Calcium Arsenazo	Arsenazo	LS	OLY0292A	R1: 4 x 60ml	2-8°C
Chloride Thiocyanate	Thiocyanate	LS	OLY0281C	R1: 4 x 30ml	2-8°C
Cholesterol CHOD PAP	CHOD-PAP	LS	OLY0261A	R1: 2 x 60ml	2-8°C
Cholesterol CHOD PAP	CHOD-PAP	LS	OLY0261B	R1: 4 x 60ml	2-8°C
Cholesterol CHOD PAP	CHOD-PAP	LS	OLY0261C	R1: 8 x 60ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	OLY0801A	R1: 5 x 20ml R2: 2 x 10ml	2-8°C
Cholinesterase	Butyryl (DGKC)	LS	OLY0801D	R1: 3 x 20ml R2: 1 x 13ml	2-8°C
СК МВ	Immunoinhibition	LS	OLY0452A	R1: 2 x 25ml R2: 1 x 10ml	2-8°C
СК МВ	Immunoinhibition	LS	OLY0452B	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
CK NAC	NAC-Activated	LS	OLY0252A	R1: 2 x 25ml R2: 1 x 10ml	2-8°C
CK NAC	NAC-Activated	LS	OLY0252B	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
Co2	Enzymatic	Mono LS	OLY0152A	R1: 4 x 20ml	2-8°C
Complement C3	Immunoturbidimetry without sample pre-dilution	LS	OLY0851B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Complement C4	Immunoturbidimetry without sample pre-dilution	LS	OLY0861A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Copper Serum/Urine	Colorimetric End Point	LS	OLY0343A	R1: 2 x 20ml	2-8°C
Creatinine Enzymatic	Enzymatic SOX	LS	OLY0113A	R1: 1 x 50ml R2: 1 x 25ml CAL: 1x1ml	2-8°C
Creatinine Enzymatic	Enzymatic SOX	LS	OLY0113B	R1: 2 x 50ml R2: 2 x 25ml CAL: 1x1ml	2-8°C
Creatinine Jaffe	Jaffe	LS	OLY0111D	R1: 2 x 30ml R2: 2 x 30ml	2-8°C
Creatinine Jaffe	Jaffe	LS	OLY0111E	R1: 3 x 50ml R2: 3 x 50ml	2-8°C
Creatinine Jaffe	Jaffe	LS	OLY0115B	R1: 4 x 60ml R2: 2 x 24ml	2-8°C
Creatinine Jaffe	Jaffe	LS	OLY0117A	R1: 3 x 50ml R2: 3 x 50ml	2-8°C
CRP Turbidimetry	Turbilatex FUll Range (Linearity 150mg/L)	LS	OLY0382A	R1: 2 x 40ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
CRP Turbidimetry	Turbilatex FUll Range (Linearity 150mg/L)	LS	OLY0382B	R1: 2 x 20ml R2: 1 x 20ml R4: 1 x 1ml	2-8°C
CRP Turbidimetry	Turbilatex FUll Range (Linearity 150mg/L)	LS	OLY0382C	R1: 4 x 40ml R2: 2 x 20ml R4: 1 x 1ml	2-8°C
Cystatin-C	Turbilatex	LS	OLY0777A	R1: 1 x 20 ml; R2: 1 x 4 ml	2-8°C
Ferritin	Turbilatex	LS	OLY0441A	R1: 1 x 20ml R2: 1 x 5ml R4: 1 x 1ml	2-8°C
Fructosamine		LS	OLY0591C	R1: 2 x 9ml R2: 2 x 21ml	2-8°C
GGT	Carboxy	LS	OLY0362A	R1: 3 x 30ml R2: 1 x 18ml	2-8°C

Description	Method	Format	Cat No.	Size	Storage
GGT	Carboxy	LS	OLY0362B	R1: 6 x 30ml R2: 2 x 18ml	2-8°C
GLDH	Enzymatic	LS	OLY0899A	R1: 2 x 20ml R2: 1 x 10ml	2-8°C
Glucose GOD PAP	GOD-PAP	LS	OLY0101A	R1: 6 x 60ml	2-8°C
Glucose GOD PAP	GOD-PAP	LS	OLY0101B	R1: 8 x 60ml	2-8°C
Glucose Hexokinase	Hexokinase	LS	OLY0103C	R1: 4 x 60ml R2: 2 x 24ml	2-8°C
HBA1C	Modified Enzymatic	LS	OLY0670A	R1: 1 x 30ml R2: 1 x 10ml	2-8°C
HBA1C	Modified Enzymatic	LS	OLY0670B	R1: 1 x 60ml R2: 1 x 20ml	2-8°C
HBA1C Direct Enzymatic	Enzymatic	LS	OLY0672A	R1: 1 x 21ml R2: 1 x 9ml	2-8°C
HDL Cholesterol (Direct)	Direct	LS	OLY0421A	R1: 3 x 30ml R2: 1 x 30ml	2-8°C
HDL Cholesterol (Direct)	Direct	LS	OLY0421B	R1: 6 x 30ml R2: 2 x 30ml	2-8°C
HDL Cholesterol (Direct)	Direct	LS	OLY0421C	R1: 4 x 70ml R2: 2 x 50ml	2-8°C
Homocysteine	Enzymatic	LS	OLY0690A	R1: 2 x 16ml R2: 1 x 8ml	2-8°C
IgE	Turbilatex	LS	OLY0751A	R1: 1 x 30ml R2: 1 x 10ml; R4: 1 x 1 ml	2-8°C
Immunoglobulin A	Immunoturbidimetry without sample pre-dilution	LS	OLY0701B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Immunoglobulin G	Immunoturbidimetry without sample pre-dilution Immunoturbidimetry without	LS	OLY0721B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Immunoglobulin M	sample pre-dilution	LS	OLY0731A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Inorganic Phosphorous	Molybdate	LS	OLY0301C	R1: 4 x 30ml R2: 2 x 15ml	2-8°C
Inorganic Phosphorous	Molybdate	LS	OLY0302A	R1: 5 x 20ml	2-8°C
Iron Ferrene	Ferrene	LS	OLY0232A	R1: 2 x 40ml R2: 1 x 20ml	2-8°C
Iron Ferrozine	Ferrozine	LS	OLY0235A	R1: 2 x 40ml R2: 2 x 10ml	2-8°C
Iron Ferrozine	Ferrozine	LS	OLY0235B	R1: 4 x 40ml R2: 2 x 20ml	2-8°C
Iron Ferrozine	Ferrozine	LS	OLY0235C	R1: 6 x 40ml R2: 3 x 20ml	2-8°C
Kappa Light Chain	Immunoturbidimetry without sample pre-dilution	LS	OLY0331B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Lactate	Enzymatic	Mono LS	OLY0622B	R1: 2 x 30ml	2-8°C
Lambda Light Chain	Immunoturbidimetry without sample pre-dilution	LS	OLY0381B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
LDH L-P	Lactate -> Pyruvate	LS	OLY0243D	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
LDH P-L	Pyruvate -> Lactate	LS	OLY0242A	R1: 5 x 20ml R2: 1 x 20ml	2-8°C
LDL Cholesterol (Direct)	Direct	LS	OLY0431A	R1: 3 x 30ml R2: 1 x 30ml	2-8°C
LDL Cholesterol (Direct)	Direct	LS	OLY0431B	R1: 6 x 30ml R2: 2 x 30ml	
LDL Cholesterol (Direct)	Direct	LS	OLY0431C	R1: 4 x 70ml R2: 2 x 50ml	
Lipase Colorimetric	Colorimetric	LS	OLY0511A	R1: 1 x 20ml R2: 1 x 10ml	2-8°C
Lipoprotein (a)	Immunoturbidimetry without sample pre-dilution	LS	OLY0130A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Magnesium Calmagite	Calmagite	LS	OLY0351C	R1: 2 x 20ml R2: 2 x 20ml	2-8°C
Magnesium Xylidyl Blue	Xylidyl Blue	LS	OLY0352A	R1: 2 x 60 ml	2-8°C
Magnesium Xylidyl Blue Microalbumin	Xylidyl Blue Immunoturbidimetry without	LS	OLY0352B OLY0471C	R2: 6 x 60 ml R1: 2 x 20ml R2: 1 x 8ml	2-8°C 2-8°C
	sample pre-dilution				
Olympus Cleaning Solution	Factor a bia	LS	OLY0001A	R1: 2000ml	2-8°C
Potassium Enzymatic	Enzymatic	LS	OLY0133B	R1: 1 x 20ml R2: 1 x 10ml	2-8°C
RF Turbidimetry	Turbilatex	LS	OLY0611A	R1: 1 x 40ml R2: 1 x 10ml, R4: 1x2ml R1: 2 x 40ml R2: 1 x	2-8°C
RF Turbidimetry	Turbilatex	LS	OLY0611B	R1: 4 x 40ml R2: 1 x 20ml, R4: 1x2ml R1: 4 x 40ml R2: 2 x	2-8°C
RF Turbidimetry	Turbilatex	LS	OLY0611C	20ml, R4: 1x2ml	2-8°C
Sodium Enzymatic	Enzymatic	LS	OLY0142B	R1: 2 x 20ml R2: 2 x 10ml	2-8°C
TIBC (Direct)	Colorimetric	LS	OLY0237A	R1: 1 x 50ml R2: 1 x 15ml	2-8°C

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
Total Protein RTU	Biuret	LS	OLY0173B	R1: 6 x 60ml	2-8°C
Transferrin	Immunoturbidimetry without sample pre-dilution	LS	OLY0741B	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
Triglycerides	GPO-PAP	LS	OLY0271A	R1: 4 x 60ml	2-8°C
Triglycerides	GPO-PAP	LS	OLY0271B	R1: 12 x 60ml	2-8°C
Urea	GLDH Kinetic & End Point	LS	OLY0123A	R1: 2 x 50ml R2: 1 x 20ml	2-8°C
Urea	GLDH Kinetic & End Point	LS	OLY0123B	R1: 4 x 60ml R2: 2 x 24ml	2-8°C
Urea	GLDH Kinetic & End Point	Lyo	OLY0121B	R1: 4 x 50ml R2: 10 x 20ml	2-8°C
Uric Acid	Uricase PAP	Lyo	OLY0601A	R1: 2 x 60ml R2: 2 x 60ml	2-8°C
Uric Acid	Uricase PAP	Lyo	OLY0601B	R1: 4 x 60ml R2: 4 x 60ml	2-8°C
Uric Acid	Uricase PAP	LS	OLY0602A	R1: 4 x 50ml R2: 1 x 50ml	2-8°C
Uric Acid	Uricase PAP	LS	OLY0602B	R1: 4 x 20ml R2: 1 x 20ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	OLY0603A	R1: 2 x 60ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	OLY0603B	R1: 4 x 60ml	2-8°C
Uric Acid	Uricase PAP	Mono LS	OLY0603C	R1: 8 x 60ml	2-8°C
Urinary CSF Protein	Pyrogallol Red	LS	OLY0172C	R1: 4 x 30ml	2-8°C
Zinc	Colorimetric	Mono LS	OLY0462A	R1: 2x 40ml	2-8°C

Furuno System Reagents

Suitable for Furuno CA180, CA270 & CA400

Description	Method	Format	Cat No.	Size	Storage
Acid Phosphatase	Fast Red	Lyo	BXCF401A	R1: 20 x 3ml	2-8°C
Acid Phosphatase	Fast Red	Lyo	BXCF401B	R1: 10 x 10ml	2-8°C
Albumin	BCG	Mono LS	BXCF222A	R1: 9 x 51ml	2-8°C
ALP (AMP)	AMP	LS	BXCF184A	R1: 6 x 51ml R2: 6 x 14ml	2-8°C
ALP (AMP)	AMP	Mono LS	BXCF184B	6 x 21ml	2-8°C
ALP (DEA)	DEA	LS	BXCF185A	R1: 6 x 51ml R2: 6 x 14ml	2-8°C
ALP (DEA)	DEA	Mono LS	BXCF185B	6 x 21ml	2-8°C
Alpha-1 Acid Glycoprotein	Immunoturbidimetric	LS	BXCF890A	R1: 3 x 16ml R2: 3 x 5ml	2-8°C
Alpha-1 Antitrypsin	Immunoturbidimetric	LS	BXCF190A	R1: 3 x 16ml R2: 3 x 5ml	2-8°C
ALT (GPT)	UV (IFCC)	LS	BXCF213A	R1: 6 x 51 ml R2: 6 x 14 ml	2-8°C
ALT (GPT)	UV (IFCC)	Mono LS	BXCF213B	R1: 6 x 17.5ml R1: 6 x 3.5ml	2-8°C
Ammonia	UV	LS	BXCF376A	R1: 4 x 20ml R2: 2 x 10ml	2-8°C
Amylase	Ethylidene PNPG7	LS	BXCF563A	R1: 4 x 16ml R2: 4 x 5ml	2-8°C
Amylase Pancreatic	Ethylidene PNPG7	LS	BXCF564A	R1: 4 x 16ml R2: 4 x 5ml	2-8°C
Anti-CCP	Turbilatex	LS	BXCF379A	R1: 1 x 30ml R2: 1 x 10ml R4: 6 x 1 x 0.5 ml	2-8°C
Anti-CCP USA ONLY	Turbilatex	LS	BXCF379B	R1: 1 x 30ml R2: 1 x 10ml R4: 6x 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	BXCF379C	R1: 1 x 30ml R2: 1 x 10ml R4: 1 x 1ml	2-8°C
Anti-CCP	Turbilatex	LS	BXCF379D	R1: 1 x 30ml R2: 1 x 10ml	2-8°C
APO A1	Immunoturbidimetric	LS	BXCF411A	R1: 4 x 30ml R2: 4 x 12ml	2-8°C
ΑΡΟ Α2	Immunoturbidimetric	LS	BXCF401A	R1: 2 x 11ml R2: 2 x 5ml	2-8°C
APO B	Immunoturbidimetric	LS	BXCF412A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
APO C2	Immunoturbidimetric	LS	BXCF413A	R1: 2 x 11ml R2: 2 x 5ml	2-8°C
APO C3	Immunoturbidimetric	LS	BXCF414A	R1: 2 x 11ml R2: 2 x 5ml	2-8°C
ΑΡΟ Ε	Immunoturbidimetric	LS	BXCF415A	R1: 2 x 11ml R2: 2 x 5ml	2-8°C

Description	Method	Format	Cat No.	Size	Storage
ASO	Turbilatex	LS	BXCF501A	R1: 4 x 20ml R2: 4 x 7ml R4: 1 x 1ml	2-8°C
ASO	Turbilatex	LS	BXCF501B	R1: 6 x 17.2ml R2: 6 x 28.4ml	2-8°C
AST	UV IFCC	LS	BXCF203A	R1: 6 x 51ml R2: 6 x 14ml	2-8°C
Barbiturates	Homogeneous Enzyme Immunoassay	LS	BXCF050A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Benzodiazepines	Homogeneous Enzyme Immunoassay	LS	BXCF051A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Beta-2-Microglobulin	Immunoturbidimetric	LS	BXCF090A	R1: 2 x 11ml R2: 2 x 4.3ml	2-8°C
Bile Acids	Colorimetric	LS	BXCF581A	R1: 2 x 18ml R2: 2 x 8ml	2-8°C
Bilirubin (Direct)	Jendrassik	LS	BXCF191A	R1: 2 x 30ml R2: 8 x 4ml	2-8°C
Bilirubin (Total)	Jendrassik	LS	BXCF192A	R1: 2 x 50ml R2: 8 x 4ml	2-8°C
C1 Acid Wash Solution		LS	BXCF912A	1 x 5L	2-8°C
Calcium	Arsenazo	Mono LS	BXCF292A	R1: 9 x 51ml	2-8°C
Calcium	CPC	LS	BXCF291A	R1: 6 x 51ml R2: 6 x 14ml	2-8°C
Calibration Serum (Level 2)		LS	BXCF321M	20 x 3ml	2-8°C
Calibration Serum (Level 3)		LS	BXCF321N	20 x 3ml	2-8°C
Cannabinoids	Homogeneous Enzyme Immunoassay	LS	BXCF052A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Carbamazepine	Latex Enhanced Immunoturbidimetric	LS	BXCF053A	R1: 2 x 12ml R2: 2 x 5ml	2-8°C
Ceruloplasmin	Immunoturbidimetric	LS	BXCF765A	R1: 3 x 20ml R2: 3 x 6.2ml	2-8°C
Cholesterol	CHOD-PAP	Mono LS	BXCF261A	R1: 9 x 51ml	2-8°C
СК-МВ	Immunoinhibition	LS	BXCF452A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
СК-МВ	Enzymatic	LS	BXCF458A	R1: 4 x 16.5ml R2: 4 x 6.2ml	2-8°C
CK-NAC	NAC-Activated	LS	BXCF252A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
CK-NAC	UV (IFCC)	Mono LS	BXCF256A	6 x 21ml	2-8°C
CO2 Total	Enzymatic	LS	BXCF152A	R1: 4 x 21.7ml	2-8°C
Cocaine Metabolite	Homogeneous Enzyme Immunoassay	LS	BXCF055A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Complement C3	Immunoturbidimetry without sample pre-dilution	LS	BXCF851A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
Complement C4	Immunoturbidimetry without sample pre-dilution	LS	BXCF861A	R1:4 x 20ml R2:4 x 6ml	2-8°C
Соррег	Colorimetric	LS	BXCF342A	R1: 5 x 20ml R2: 1 x 30ml	2-8°C
Creatinine	Jaffe	LS	BXCF117A	R1: 6 x 51ml R2: 3 x 51ml	2-8°C
Creatinine Enzymatic	Enzymatic SOX	LS	BXCF113B	R1: 2 x 50ml R2: 1 x 50ml CAL: 2x1ml	2-8°C
CRP	Immunoturbidimetric	LS	BXCF384A	R1: 6 x 20ml R2: 3 x 9ml	2-8°C
CRP (Full Range)	Turbidimetric	LS	BXCF382A	R1: 2 x 22ml R2: 1 x 11ml	2-8°C
CRP (Full Range)	Turbidimetric	LS	BXCF382B	R1: 4 x 50ml R2: 1 x 50ml	2-8°C
CRP (Full Range)	Turbidimetric	LS	BXCF382C	R1: 5 x 100ml R2: 1 x 100ml	2-8°C
CRP (High Sensitivity)	Immunoturbidimetric	LS	BXCF383A	R1: 2 x 11ml R2: 2 x 11ml	2-8°C
Cystatin-C	Immunoturbidimetric	LS	BXCF777A	R1: 2 x 17.6ml R2: 2 x 6.1ml	2-8°C
D-3-Hydroxybutyrate	UV	LS	BXCF541A	R1: 5x10ml R2: 5x10ml R4: 1x5ml	2-8°C
D-3-Hydroxybutyrate	UV	LS	BXCF541B	R1: 10x5ml R2: 10x5ml R4: 1x5ml	2-8°C
Digoxin	Immunoturbidimetric	LS	BXCF056A	R1: 2 x 8ml R2: 2 x 6ml	2-8°C
Direct HDL	Direct	LS	BXCF421A	R1: 3 x 51ml R2: 3 x 20ml	2-8°C

SECTION 1 | Analyser Specific Reagents

Description	Method	Format	Cat No.	Size	Storage
Direct LDL	Direct	LS	BXCF431A	R1: 3 x 51ml R2: 3 x 20ml	2-8°C
Ecstasy	Homogeneous Enzyme Immunoassay	LS	BXCF057A	R1: 2x16.9ml R2: 2x8ml	2-8°C
EDDP	Homogeneous Enzyme Immunoassay	LS	BXCF058A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Ethanol	Enzymatic	LS	BXCF059A	R1: 2 x 16.9ml R2: 2 x 8ml	2-8°C
Ferritin	Turbilatex	LS	BXCF441A	R1: 3 x 20ml R2: 3 x 11ml	2-8°C
Fructosamine	Enzymatic	LS	BXCF591A	R1: 4 x 19.8ml R2: 4 x 6.9ml	2-8°C
G6PDH		LS	BXCF571A	R1a: 2 x 50ml R1b: 1 x 2ml R2: 1 x 20ml R3: 1 x 20ml	2-8°C
Gamma GT (GGT)	Jaffe	LS	BXCF362A	R1: 6 x 51ml R2: 6 x 14ml	2-8°C
Gamma GT (GGT)	Colorimetric	Mono LS	BXCF360A	R1a: 6 x 17.5ml R1b: 6 x 3.5ml	2-8°C
Gentamicin	Immunoturbidimetric		BXCF060A	R1: 2 x 15ml R2: 2 x 6ml	2-8°C
GLDH	Enzymatic	LS	BXCF899A	R1: 2 x 20ml R2: 2 x 5ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXCF101A	R1: 4 x 20ml	2-8°C
Glucose	GOD-PAP	Mono LS	BXCF101B	R1: 9 x 51ml	2-8°C
Glucose	Hexokinase	LS	BXCF103A	R1: 4 x 51ml R2: 3 x 20ml	2-8°C
Glucose	Hexokinase	Mono LS	BXCF103B	R1: 4 x 50ml	2-8°C
Glutamate	Colorimetric	LS	BXCF061A	R1: 4 x 20ml	2-8°C
Glutamine	Colorimetric	LS	BXCF062A	R1: 4 x 16ml R2: 4 x 6.7ml	2-8°C
Glutathione Peroxidase	Enzymatic	LS	BXCF551A	R1: 8 x 10ml	2-8°C
Haemoglobin Denaturant Reagent	Immunoturbidimetric	LS	BXCF099A	R1: 2 x 50ml	2-8°C
Haptoglobin	Immunoturbidimetric	LS	BXCF496A	R1: 2 x 20ml R2: 1 x 8ml	2-8°C
HbA1c	Enzymatic	LS	BXCF670X	R1: 4 x 20ml R2: 4 x 8.6ml R3: 2 x 100ml	2-8°C
Homocysteine	Enzymatic	LS	BXCF690A	R1: 2 x 21.7ml R2: 2 x 4.6ml	2-8°C
IGA	Immunoturbidimetric	LS	BXCF701A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
IGA	Immunoturbidimetric	LS	BXCF701B	R1: 3 x 20ml R2: 3 x 6.2ml	2-8°C
IGE	Immunoturbidimetric	LS	BXCF751A	R1: 2 x 10ml R2: 2 x 5ml R4: 1 x 1ml	2-8°C
IGG	Immunoturbidimetric	LS	BXCF721A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
IGG	Immunoturbidimetric	LS	BXCF721B	R1: 3 x 20ml R2: 3 x 6.2ml	2-8°C
IGM	Immunoturbidimetric	Mono LS	BXCF731A	R1: 4 x 20ml R2: 4 x 6ml	2-8°C
IGM	Immunoturbidimetric	LS	BXCF731B	R1: 3 x 20ml R2: 3 x 6.2ml	2-8°C
Inorganic Phosphorus	Molybdate	LS	BXCF301A	R1: 6 x 20ml R2: 3 x 20ml	2-8°C
Inorganic Phosphorus	Molybdate	Mono LS	BXCF302A	R1: 6 x 20ml	2-8°C
Ion Selective Electrodes (ISE-CAL B)		LS	BXCF908A	R1: 1 x 125ml	2-8°C
Iron	Ferrene	LS	BXCF232A	R1:6x20ml R2:3x 11ml	2-8°C
ISE Calibrator A		LS	BXCF909A	R1: 1 x 500ml	2-8°C
ISE Urine Diluent		LS	BXCF910A	R1: 2 x 50ml	2-8°C
ISE Wash Solution		LS	BXCF911A	R1: 1 x 90ml R2: 6 x 15ml	2-8°C
Lactate	Colorimetric	Mono LS	BXCF622A	R1: 4 x 20.5ml	2-8°C
LDH L-P	Lactate -> Pyruvate	LS	BXCF243A	R1:6x20ml R2:3x 18ml	2-8°C
LDH P-L	Pyruvate -> Lactate	LS	BXCF242A	R1:6x20ml R2:3x 11ml	2-8°C
Lipase	Colorimetric	LS	BXCF511A	R1: 3 x 9ml R2: 3 x 6ml	2-8°C

Description	Method	Format	Cat No.	Size	Storage
Lipoprotein (a)	Immunoturbidimetric	LS	BXCF130A	R1: 2 x 10ml R2: 2 x 6ml R4: 1 x 1ml	2-8°C
Lithium	Colorimetric	LS	BXCF125A	R1: 2 x 18.3ml R2: 2 x 6.5ml	2-8°C
Magnesium (Xylidl Blue)	Xylidyl Blue	Mono LS	BXCF352A	R1: 6 x 20ml	2-8°C
Methadone	Homogeneous Enzyme Immunoassay	LS	BXCF063A	R1: 2x16.9ml R2: 2 x 8ml	2-8°C
Microalbumin	Immunoturbidimetric	LS	BXCF471A	R1: 6 x 20ml R2: 3 x 8ml	2-8°C
NEFA	UV Colorimetric	Mono LS	BXCF477A	R1: 1 x 40ml R2: 1 x 10ml CAL: 1 x 1ml	2-8°C
Opiates	Homogeneous Enzyme Immunoassay	LS	BXCF065A	R1: 2 x 16.9ml R2 2 x 8ml	2-8°C
Phenobarbital	Latex Enhanced Immunoturbidimetric	LS	BXCF067A	R1: 2 x 17ml R2: 2 x 6ml	2-8°C
Phosphorus Inorganic	Molybdate	Mono LS	BXCF302A	R1: 6 x 20ml	2-8°C
Potassium	ТРВ	LS	BXCF135A	R1: 3 x 20ml R2: 3 x 9ml	2-8°C
Reagent Precision Test Solution (RABS)		LS	BXCF902A	R1: 1 x 20ml	2-8°C
RF	Immunoturbidimetric Turbilatex	LS	BXCF611A	R1: 2 x 20ml R2: 2 x 6ml R4: 1 x 1ml	2-8°C
Saline Diluent		LS	BXCF903A	R1: 10 x 100ml	2-8°C
Sample Precision Test Solution (SABS)		LS	BXCF901A	R1: 2 x 10ml	2-8°C
SMS Acid Wash Solution (HCL)		LS	BXCF907A	R1: 9 x 50ml (200 mmol/l)	2-8°C
Sodium	Enzymatic	LS	BXCF142A	R1: 3 x 20ml R2: 3 x 10ml	2-8°C
Superoxide Dismutase	Colorimetric	LS	BXCF531A	R1: 5x20ml	2-8°C
тівс	GPO-PAP	LS	BXCF237A	R1: 4 x 9ml R2: 4 x 4ml	2-8°C
Total Protein	Biuret	LS	BXCF171A	R1: 4 x 51ml R2: 4 x 44ml	2-8°C
Total Protein	Biuret	Mono LS	BXCF173A	R1: 9 x 51ml	2-8°C
Transferrin	Immunoturbidimetric	LS	BXCF741A	R1: 6 x 20ml R2: 3 x 14ml	2-8°C
Transthyrethin (PreAlbumin)	Immunoturbidimetric	LS	BXCF070A	R1: 6 x 20ml R2: 3 x 11ml	2-8°C
Triglycerides	GPO-PAP	Mono LS	BXCF271A	R1: 6 x 51ml	2-8°C
Urea	GLDH Kinetic & Endpoint	LS	BXCF123A	R1: 1 x 100ml R2: 1 x 20ml	2-8°C
Urea	Enzymatic Kinetic	LS	BXCF124A	R1: 6 x 51ml R2: 4 x 20ml	2-8°C
Urea	Enzymatic Kinetic	Mono LS	BXCF126A	6 x 21ml	2-8°C
Uric Acid	Uricase PAP	LS	BXCF602A	R1: 6 x 51ml R2: 4 x 20ml	2-8°C
Uric Acid	Enzymatic Colorimetric	Mono LS	BXCF603A	R1: 9 x 51ml	2-8°C
Urinary Protein	Pyrogallol Red	Mono LS	BXCF172A	R1: 3 x 100ml	2-8°C
Valproic Acid	Immunoturbidimetric	LS	BXCF071A	R1: 2 x 15ml R2: 2 x 6ml	2-8°C
Wash Solution No. 1 (Acid)		LS	BXCF904A	6 x 25ml	2-8°C
Wash Solution No. 2 (Alkali)		LS	BXCF905A	6 x 25ml	2-8°C
Wash Solution No. 3 (Neutral)		LS	BXCF906A	6 x 25ml	2-8°C
Zinc	Colorimetric	LS	BXCF462A	R1: 2 x 40ml R2: 1 x 5ml	2-8°C

SECTION 2 QUALITY CONTROLS & CALIBRATORS

AL REAL

Solution Solution Solution Solution

10

50

Quality Controls & Calibrators

Quality controls and calibrators play a vital role in clinical diagnostics by ensuring the accuracy and reliability of test results. Quality controls are designed to monitor the performance of diagnostic assays and reagents, helping to detect any variations or issues that may arise during testing. This ongoing assessment ensures that laboratories maintain high standards and can identify problems before they affect patient care.

Calibrators, on the other hand, are used to establish the measurement accuracy of diagnostic instruments by providing a known reference point against which test results can be compared. Together, these elements contribute to standardisation and consistency in laboratory testing, fostering confidence in the results provided to healthcare professionals. Ultimately, robust quality control and calibration processes are essential for delivering precise diagnoses, guiding effective treatment decisions, and ensuring patient safety in clinical practice.

Fortress Quality Controls (**Seraqual**) are supplied with instrument and method specific assigned values. We can also provide custom made quality control solutions, tailor-made to customer specification and requirements, enabling laboratories to be more efficient and reduce costs. We also work with many universities, research institutions and laboratories to develop quality control and calibration solutions for unique and novel biomarkers.

The Seraqual Portfolio

Fortress Diagnostics Seraqual Controls and Calibrators are available for:

- Antioxidants Aqueous Standards Blood Gas & Electrolytes Cardiac Clinical Chemistry Coagulation/Haemostasis Diabetes
- Drugs of Abuse Haematology Immunoassay/Infectious Immunology/Proteins Lipids Maternal Screening Neonatal Screening Paediatric
- Speciality Controls Thalassaemia Therapeutic Drugs Torch Paediatric Urine Xanthochromia





Why Choose Fortress Quality Controls & Calibrators?



Customer Support

Fortress Diagnostics employ a highly-skilled team of technical support staff to assist customers with all their needs and technical questions. We are always on-hand to help.





Cost Saving

Our Quality Controls Range have a wide-ranging analyte content, enabling laboratories to consolidate their quality control purchases and therefore benefit from cost savings.





Accurate

Manufactured by Fortress Diagnostics in our ISO 13485:2016 accredited labs, Seraqual products are designed to provide highly accurate target values, mimicking human samples, with minimal matrix effects being observed.





Reliable

Our fully in-house logistics team enables us to ship orders globally and our customer support team are available to assist with any labs' special requirements or custom orders to ensure customer satisfaction is always exceeded.





Third Party Controls Suitable For Global Market

We manufacture a wide range of third party controls which are valueassigned, without bias, towards any methods or instrumentation, enabling labs to meet their ISO accreditations through accurate and robust training.





High Quality

Seraqual products are manufactured without the use of preservatives or stabilisers, utilising human-based material, which have undergone stringent quality checks for HIV, HBsAg, HCV, HTLV, and Syphilis for FDA approved methods. Products of animal origin are not added in the Fortress controls, therefore eliminating the lack of specificity in antibody-based tests.





Flexible

As well as offering an extensive range of controls, Fortress Diagnostics has the ability to manufacture custom quality control products to meet any customer requirements providing a fully flexible solution.





Long Shelf Life

Fortress Diagnostics offer quality controls in two different formats - lyopholised (stable for up to 4 years) and liquid (stable with a shelf life of up to 18 months) - enabling labs to utilise lot supply and save costs.





Traceability

Fortress Controls and Calibrators are CE marked and fully traceable to the reference material according to ISO 17511 and ISO 18153 directives.





Consolidation

Fortress Quality Controls offer a wide parameter range allowing more analyte coverage with a single control.





Value Assignment

The Fortress value assignment involves six European Reference Laboratories, (chosen on the basis of the analytes performed in these laboratories), each of which performs five independant runs.





Compatibility

Fortress Controls are made from human derived material, with minimal matrix effects being observed.



Fortress Quality Control Inserts Portal

Download e-inserts for the Fortress Diagnostics Quality Controls & Calibrators range through our online portal.

Fortress Diagnostics are excited to launch the new Fortress QC Inserts Portal. The portal allows customers to easily search our database of inserts by product name, catalogue number or batch number in our easy-to-use filter tool. Inserts can be viewed online or downloaded in PDF format.

insert.fortressdiagnostics.com



Description	Details	Format	Cat No.	Size	Storage
AB0-Rhesus Blood Grouping Control Set		LS	BXC0070A	4 x 6ml	2-8°C
ADA Calibrator		LS	BXC0209A	1 x 1ml	2-8°C
ADA Calibrator		Lyo	BXC0217A	1 x 1ml	2-8°C
ADA Control (Level 1)		Lyo	BXC0218A	1 x 1ml	2-8°C
ADA Control (Level 1)		LS	BXC0210A	1 x 1ml	2-8°C
ADA Control (Level 2)		Lyo	BXC0219A	1 x 1ml	2-8°C
ADA Control (Level 2)		LS	BXC0216A	1 x 1ml	2-8°C
Alcohol, Ammonia, Carbonate Calibrator		LS	BXC0492A	1 x 2ml	2-8°C
Alcohol, Ammonia, Carbonate Control (Level 1)		LS	BXC0493A	1 x 2ml	2-8°C
Alcohol, Ammonia, Carbonate Control (Level 2)		LS	BXC0494A	1 x 2ml	2-8°C
Aldolase Calibrator		LS	BXC0394A	3 x 1ml	2-8°C
Aldolase Control Elevated		LS	BXC0393A	3 x 1ml	2-8°C
Aldolase Control Normal		LS	BXC0392A	3 x 1ml	2-8°C
Alpha-1-Acid Glycoprotein Calibrator		LS	BXC0892A	2 x 1ml	2-8°C
Alpha-1-Acid Glycoprotein Control		LS	BXC0891A	2 x 1ml	2-8°C
Alpha-1-Antitrypsin Calibrator	Immunoturbidimetric	LS	BXC0711A	1 x 2ml	2-8°C
Alpha-1-Antitrypsin Control (Level 1 & Level 2)	Immunoturbidimetric	LS	BXC0712A	2 x 1 x 2ml	2-8°C
Alpha-1-Microglobulin Calibrator		LS	BXC0887A	1 x 1ml	2-8°C
Alpha-1-Microglobulin Control (Level 1 & Level 2)		LS	BXC0889A	2 x 1 x 1ml	2-8°C
Aluminium Control (Level 1)		LS	BXC0696A	1 x 1ml	2-8°C
Aluminium Control (Level 2)		LS	BXC0697A	1 x 1ml	2-8°C
Aluminium Control (Level 3)		LS	BXC0698A	1 x 1ml	2-8°C
Ammonia Calibrator		LS	BXC0373A	3 x 2ml	2-8°C
Ammonia Control High	High	LS	BXC0375A	3 x 2ml	2-8°C
Ammonia Control Low	Low	LS	BXC0374A	3 x 2ml	2-8°C
Angiotensin Converting Enzyme (ACE) Control (Level 1)		LS	BXC0178A	3 x 1ml	2-8°C
Angiotensin Converting Enzyme (ACE) Control (Level 2)		LS	BXC0179A	3 x 1ml	2-8°C
Angiotensin Converting Enzyme (ACE) Calibrator		LS	BXC0177A	3 x 1ml	2-8°C
Anti-CCP Calibrator		LS	BXC0379D	1 x 1ml	2-8°C
Anti-CCP Calibrator Series		LS	BXC0379B	6 x 1 x 1ml	2-8°C
Anti-CCP Control Set		LS	BXC0379C	2 x 1 x 1ml	2-8°C
Anti-Mullerian Hormone (AMH) Calibrators & Control Set		LS	BXC0999A	Cal A: 1 x 0.5ml; Cal B-G: 6 x 0.5ml; Ctl L1: 1 x 0.5ml; Ctl L2: 1 x 0.5ml	2-8°C
APO A1/B Calibrator Series		LS	BXC0413A	6 x 1 x 1ml	2-8°C
ASO & RF Control		LS	BXC0640A	3 x 1ml	2-8°C
ASO (Single Point) Calibrator	Immunoturbidimetric	LS	BXC0323A	1 x 1ml	2-8°C
ASO (Single Point) Calibrator	Immunoturbidimetric	LS	BXC0323B	3 x 1ml	2-8°C
ASO, CRP, RF Control	Immunoturbidimetric	LS	BXC0645A	3 x 1ml	2-8°C
Beta-2-Microglobulin Calibrator		LS	BXC0647A	1 x 1 ml	2-8°C
Beta-2-Microglobulin Control (Level 1 &		LS	BXC0648A	1 x 2 x 1 ml	2-8°C
Level 2)					
Level 2) Bile Acid Calibrator		LS	BXC0582A	1 x 1ml	2-8°C
•		LS LS	BXC0582A BXC0583A	1 x 1ml 2 x 1 x 1ml	2-8°C 2-8°C
Bile Acid Calibrator					

Description	Details	Format	Cat No.	Size	Storage
Bilirubin Control	Normal	LS	BXC0306A	1 x 1ml	2-8°C
Bilirubin Control	Normal	LS	BXC0306B	3 x 1ml	2-8°C
Bilirubin Control	Elevated	LS	BXC0307A	1 x 1ml	2-8°C
Bilirubin Control	Elevated	LS	BXC0307B	3 x 1ml	2-8°C
Bilirubin Control	Elevated	Lyo	BXC0318A	1 x 1ml	2-8°C
Bilirubin Control	Elevated	Lyo	BXC0318B	5 x 1ml	2-8°C
Bilirubin Control Set		LS	BXC0304A	2 x 1 x 1ml	2-8°C
Bilirubin Control Set, Paediatric		LS	BXC0305A	2 x 1 x 1ml	2-8°C
Blood Alcohol Calibrator		LS	BXC0488A	1 x 2ml	2-8°C
Blood Alcohol Control (Level 1)		LS	BXC0489A	1 x 2ml	2-8°C
Blood Alcohol Control (Level 2)		LS	BXC0490A	1 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 1)		LS	BXC0108A	1 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 1)		LS	BXC0108D	10 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 2)		LS	BXC0108B	1 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 2)		LS	BXC0108E	10 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 3)		LS	BXC0108C	1 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Level 3)		LS	BXC0108F	10 x 2ml	2-8°C
Blood Gas & Electrolyte Control (Tri-					
Level)		LS	BXC0108G	30 x 2ml	2-8°C
Calibration Serum		LS	BXC0321K	1 x 3ml	2-8°C
Calibration Serum		LS	BXC0321L	5 x 3ml	2-8°C
Calibration Serum		LS	BXC0321M	10 x 3ml	2-8°C
Cardiac Control (Level 1)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0455A	5 x 2ml	2-8°C
Cardiac Control (Level 1)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0455B	5 x 1ml	2-8°C
Cardiac Control (Level 1)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0455C	5 x 3ml	2-8°C
Cardiac Control (Level 2)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0456A	5 x 2ml	2-8°C
Cardiac Control (Level 2)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0456B	5 x 1ml	2-8°C
Cardiac Control (Level 2)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0456C	5 x 3ml	2-8°C
Cardiac Control (Level 3)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0464A	5 x 2ml	2-8°C
Cardiac Control (Level 3)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0464B	5 x 1ml	2-8°C
Cardiac Control (Level 3)	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0464C	5 x 3ml	2-8°C
Cardiac Control Set	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0450A	3 x 1 x 2ml	2-8°C
Cardiac Control Set	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0450B	3 x 1 x 1ml	2-8°C
Cardiac Control Set	CK, CKMB, Tni,TnT, Digoxin, Homocysteine, hsCRP, BNP, NT- ProBNP	LS	BXC0450C	3 x 1 x 3ml	2-8°C
Cardiac Tri Level Control		LS	BXC0881A	3 x 1 x 0.5ml	2-8°C
Cardiac Tri Level Control Citrate (Urinary) Calibrator	200mg/L	LS LS	BXC0881A BXC0136A	3 x 1 x 0.5ml 2 x 1ml	2-8°C 2-8°C

Description	Details	Format	Cat No.	Size	Storage
Citrate (Urinary) Control (Level 2)	600mg/L	LS	BXC0139A	2 x 1ml	2-8°C
CK / CKMB Calibrator		LS	BXC0454A	1 x 2ml	2-8°C
CK / CKMB Calibrator		LS	BXC0454B	5 x 2ml	2-8°C
CK / CKMB Control (Level 1)		LS	BXC0453A	1 x 2ml	2-8°C
CK / CKMB Control (Level 1)		LS	BXC0453B	5 x 2ml	2-8°C
CK / CKMB Control (Level 2)		LS	BXC0459A	1 x 2ml	2-8°C
CK / CKMB Control (Level 2)		LS	BXC0459B	5 x 2ml	2-8°C
CO2 (Bicarbonate) Calibrator		LS	BXC0155A	3 x 2ml	2-8°C
CO2 (Bicarbonate) Control High	High	LS	BXC0157A	3 x 2ml	2-8°C
CO2 (Bicarbonate) Control Low	Low	LS	BXC0156A	3 x 2ml	2-8°C
COVID-19 Anitgen Control Pair	Positive and Negative Control	LS	COVIDNPC	2 x 1 x 0.5ml	2-8°C
CRP (Multi Point) Calibrator	Immunoturbidimetric	LS	BXC0324B	6 x 1 x 1ml	2-8°C
CRP (Single Point) Calibrator	Immunoturbidimetric	LS	BXC0324A	3 x 1ml	2-8°C
CRP (Ultra Sensitive) Low Control	Immunoturbidimetric	LS	BXC0325A	3 x 1ml	2-8°C
CRP (Ultra Sensitive) Standard Set	Immunoturbidimetric	LS	BXC0327A	5 x 1 x 0.5ml	2-8°C
CRP Control (Level 1)		LS	BXC0326A	5 x 1ml	2-8°C
CRP Control (Level 2)		LS	BXC0326B	5 x 1ml	2-8°C
CRP Control Set		LS	BXC0326C	2 x 1 x1 ml	2-8°C
CSF Control (Level 1)	Albumin, Chloride, Glucose, IgA, IgG,IgM, Lactate, Sodium, Total Protein	LS	BXC0673A	1 x 1ml	2-8°C
CSF Control (Level 2)	Albumin, Chloride, Glucose, IgA, IgG,IgM, Lactate, Sodium, Total Protein	LS	BXC0673B	1 x 1ml	2-8°C
CSF Diff Control (Level 1)		LS	BXC0080A	5 x 2ml	2-8°C
CSF Diff Control (Level 2)		LS	BXC0081A	5 x 2ml	2-8°C
CSF Diff Control (Level 3)		LS	BXC0082A	5 x 2ml	2-8°C
Cyanmethaemoglobin Standard Set	8,10,12,15,18 g/dL	LS	BXC0483A	5 x 1 x 10ml	2-8°C
Cystatin C Calibrator Set (6 Levels)		LS	BXC0334A	6 x 1ml	2-8°C
Cystatin C Control (Level 1)		LS	BXC0333A	2 x 1ml	2-8°C
Cystatin C Control (Level 2)		LS	BXC0333B	2 x 1ml	2-8°C
D3-Hydroxybutyrate Calibrator		Lyo	BXC0544A	1 x 3ml	2-8°C
D3-Hydroxybutyrate Control (Level 1)		Lyo	BXC0543A	1 x 5ml	2-8°C
D3-Hydroxybutyrate Control (Level 2)		Lyo	BXC0543B	1 x 5ml	2-8°C
D-Dimer Turbilatex Calibrator		LS	BXC0788A	1 x 1ml	2-8°C
D-Dimer Turbilatex Control Set (Level 1 & Level 2)		LS	BXC0789A	2 x 1 x 1ml	2-8°C
Drugs of Abuse Control (Level 1)		LS	BXC0784A	10 x 10ml	2-8°C
Drugs of Abuse Control (Level 1)		LS	BXC0784B	10 x 3ml	2-8°C
Drugs of Abuse Control (Level 2)		LS	BXC0785A	10 x 10ml	2-8°C
Drugs of Abuse Control (Level 2)		LS	BXC0785B	10 x 3ml	2-8°C
Drugs of Abuse Control (Level 3)		LS	BXC0786A	10 x 10ml	2-8°C
Drugs of Abuse Control (Level 3)		LS	BXC0786B	10 x 3ml	2-8°C
Electrolytes Calibrator		LS	BXC0140A	5 x 5ml	2-8°C
Electrolytes Control (Level 1)	Low (Na, K, Cl)	LS	BXC0143A	5 x 5ml	2-8°C
Electrolytes Control (Level 2)	Normal (Na, K, Cl)	LS	BXC0144A	5 x 5ml	2-8°C
Electrolytes Control (Level 3)	High (Na, K, Cl)	LS	BXC0145A	5 x 5ml	2-8°C
ESR Control (Level 1)	To be used with all routine ESR instruments	LS	BXC0631A	2 x 10ml	2-8°C
ESR Control (Level 2)	To be used with all routine ESR instruments	LS	BXC0632A	2 x 10ml	2-8°C
Ferritin Calibrator		LS	BXC0445A	1 x 1ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
Ferritin Calibrator Set		LS	BXC0445B	6 x 1 x 1ml	2-8°C
Ferritin Control (Level 1)		LS	BXC0443A	1 x 1ml	2-8°C
Ferritin Control (Level 2)		LS	BXC0444A	1 x 1ml	2-8°C
Fructosamine Calibrator		LS	BXC0592A	1 x 1ml	2-8°C
Fructosamine Control (Level 1)	Low	LS	BXC0593A	3 x 1ml	2-8°C
Fructosamine Control (Level 2)	High	LS	BXC0594A	3 x 1ml	2-8°C
G-6-PDH Control	Deficient	LS	BXC0572B	3 x 1.0ml	2-8°C
G-6-PDH Control	Normal	LS	BXC0573B	3 x 1.0ml	2-8°C
Glucose-Fructose Control (Level 1)		LS	BXC0099A	ТВС	2-8°C
Glucose-Fructose Control (Level 2)		LS	BXC0099B	ТВС	2-8°C
Glutathione Peroxidase Control		LS	BXC0556A	5 x 1ml	2-8°C
Glutathione Reductase Control		LS	BXC0460A	5 x 1ml	2-8°C
Glycerol Control		LS	BXC0278A	2 x 5ml	2-8°C
Haematology Calibrator	Haematology calibrator for 3-part differential Analysers	LS	HAEMC005	3 x 3ml	2-8°C
Haematology Control (3 Levels)	Haematology control for 3-part differential Analysers	LS	HAEMC004	3 x 3 x 3ml	2-8°C
Haematology Control (Level 1)	Haematology control for 3-part differential Analysers	LS	HAEMC001	3 x 3ml	2-8°C
Haematology Control (Level 2)	Haematology control for 3-part differential Analysers	LS	HAEMC002	3 x 3ml	2-8°C
Haematology Control (Level 3)	Haematology control for 3-part differential Analysers	LS	HAEMC003	3 x 3ml	2-8°C
HbA1c (HPLC) Calibrator Set (2 Levels)	For HPLC Method Only	LS	BXC0780A	2 x 1 x 0.25ml	2-8°C
HbA1c (HPLC) Control Set (2 Levels)	For HPLC Method Only	LS	BXC0779A	2 x 1 x 0.5ml	2-8°C
HbA1c Calibrator Set	For use with BXC0670	LS	BXC0668A	2 x 1 x 0.5ml	2-8°C
HbA1c Calibrator Set (2 Levels)	Most Common Methods including HPLC	LS	BXC0678A	2 x 1 x 0.5ml	2-8°C
HbA1c Control High	Most Common Methods including HPLC	LS	BXC0677A	2 x 0.5ml	2-8°C
HbA1c Control Low	Most Common Methods including HPLC	LS	BXC0676A	2 x 0.5ml	2-8°C
HbA1c Control Set	Most Common Methods including HPLC	LS	BXC0675A	2 x 2 x 0.5ml	2-8°C
HbA1c Control Set (2 Levels)	For use with BXC0670	LS	BXC0669A	2 x 1 x 0.5ml	2-8°C
HbA2 Control Set	HPLC	LS	BXC0778A	2 x 2 x 0.5ml	2-8°C
HBsAb Control Panel (2 Levels)		LS	BXC0805A	2 x 1 x 0.5ml	2-8°C
HBsAg Control Panel (5 Levels)		LS	BXC0800A	5 x 1 x 0.5ml	2-8°C
hCG Serum Control	Positive	LS	BXC0681A	1 x 1ml	2-8°C
hCG Serum Control	Negative	LS	BXC0682A	1 x 1ml	2-8°C
hCG Serum Control	Positive	LS	BXC0683A	1 x 5ml	2-8°C
HCV Antibody Control Panel (4 Levels)		LS	BXC0804A	4 x 1 x 0.5ml	2-8°C
HDL/LDL Calibrator		LS	BXC0315B	1 x 3ml	2-8°C
HDL/LDL Calibrator		LS	BXC0315C	5 x 3ml	2-8°C
HDL/LDL Calibrator		LS	BXC0315D	5 x 1ml	2-8°C
HDL/LDL Calibrator		LS	BXC0315E	1 x 1ml	2-8°C
HDL/LDL Calibrator		LS	BXC0315F	3 x 1ml	2-8°C
HE4 (Level 1)		LS	BXC0425A	1 x 0.25ml	2-8°C
HE4 (Level 2)		LS	BXC0425B	1 x 0.25ml	2-8°C
HIV-I Antibody Control Panel (4 Levels)		LS	BXC0802A	4 x 1 x 1ml	2-8°C
HIV-I Antigen (p24) Control Panel (4 Levels)		LS	BXC0803A	4 x 1 x 0.5ml	2-8°C
Homocysteine Calibrator (2 Levels)		LS	BXC0691B	2 x 1 x 1ml	2-8°C
Homocysteine Control (2 Levels)			BXC0692A	2 x 1 x 1ml	2-8°C
H-Pylori Control (2 Levels)			BXC0684A	2 x 1 x 1ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
Human Assayed Control (Level 1)		Lyo	BXC0312A	10 x 5ml	2-8°C
Human Assayed Control (Level 1)		Lyo	BXC0312C	5 x 5ml	2-8°C
Human Assayed Control (Level 1)		Lyo	BXC0312E	1 x 5ml	2-8°C
Human Assayed Control (Level 1)		Lyo	BXC0314A	10 x 5ml	2-8°C
Human Assayed Control (Level 1)		Lyo	BXC0314C	5 x 5ml	2-8°C
Human Assayed Control (Level 1)		Lyo	BXC0314E	1 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0312B	10 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0312D	5 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0312F	1 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0314B	10 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0314D	5 x 5ml	2-8°C
Human Assayed Control (Level 2)		Lyo	BXC0314F	1 x 5ml	2-8°C
IgE Calibrator		Lyo	BXC0752A	1 x 1ml	2-8°C
IgE Control Set (Level 1 & Level 2)		Lyo	BXC0753A	2 x 2 x 3ml	2-8°C
Immunoassay Control (Level 1)		LS	BXC0363A	4 x 3ml	2-8°C
Immunoassay Control (Level 2)		LS	BXC0363B	4 x 3ml	2-8°C
Immunoassay Control (Level 3)		LS	BXC0363C	4 x 3ml	2-8°C
Immunoassay Speciality Control (Level 1)	Vitamin-D, Anti TG, Anti- TPO, C-Peptide, IGF-1, Insulin,Osteocalcin, PTH, Procalcitonin	LS	BXC0364A	1x2 ml	2-8°C
Immunoassay Speciality Control (Level 2)	Vitamin-D, Anti TG, Anti- TPO, C-Peptide, IGF-1, Insulin,Osteocalcin, PTH, Procalcitonin	LS	BXC0364B	1x2 ml	2-8°C
Immunoassay Speciality Control (Level 3)	Vitamin-D, Anti TG, Anti- TPO, C-Peptide, IGF-1, Insulin,Osteocalcin, PTH, Procalcitonin	LS	BXC0364C	1x2 ml	2-8°C
Immunoassay Tri-Level Control		LS	BXC0363D	3 x 4 x 3ml	2-8°C
Lipase, Calibrator		LS	BXC0510A	3 x 1ml	2-8°C
Lipid Calibrator	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0317A	3 x 1ml	2-8°C
Lipid Control	Elevated, Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0316A	3 x 1ml	2-8°C
Lipid Control	Normal, Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0330A	3 x 1ml	2-8°C
Lipid Control (Level 1)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0347A	1 x 3 ml	2-8°C
Lipid Control (Level 1)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0347B	5 x 3 ml	2-8°C
Lipid Control (Level 2)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0348A	1 x 3 ml	2-8°C
Lipid Control (Level 2)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL Chalasterol, Triglycerides, APO A1	LS	BXC0348B	5 x 3 ml	2-8°C
Lipid Control (Level 3)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0349A	1 x 3 ml	2-8°C
Lipid Control (Level 3)	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0349B	5 x 3 ml	2-8°C
Lipid Control Set	Cholesterol, Triglycerides, APO A1, APO B, HDL, LDL	LS	BXC0350A	3 x 1 x 3 ml	2-8°C
Lipoprotein (a) Calibrator	High	LS	BXC0134A	1 x 1ml	2-8°C
Lipoprotein (a) Control	Low	LS	BXC0131A	1 x 1ml	2-8°C
Lipoprotein (a) Control	High	LS	BXC0133A	1 x 1ml	2-8°C
Lithium Calibrator	Only for Enzymatic Lithium	LS	BXC0141A	1 x2x1ml	2-8°C
Lithium Calibrator	3000 mmol/L	LS	BXC0106A	2 x 1ml	2-8°C

Maternal Screening Control (Level 1)For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems LSLSBXC0695A2 x 1mlMaternal Screening Control (Level 2)For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems LSLSBXC0695A2 x 1mlMaternal Screening Control (Level 2)For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems AFP, F-beta-hCG, Total hCG, UnConjugated Estriol, Inhibin-A, PAPP-A, AMHSBXC0695B2 x 1mlMaternal Screening Control (Level 3)For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems AFP, F-beta-hCG, Total hCG, UnConjugated Estriol, Inhibin-A, PAPP-A, AMHSBXC0695B2 x 1mlMaternal Screening Control (Level 3)For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems LSLSBXC0695C2 x 1ml	2-8°C 2-8°C
Maternal Screening Control (Level 2) & Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems LS BXC0695B 2 x 1ml Maternal Screening Control (Level 2) For Screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems LS BXC0695B 2 x 1ml Maternal Screening Control (Level 3) For screening of Down's syndrome including Immunoassay systems LS BXC0695C 2 x 1ml	2-8°C
& Spina Bifida compatible for use on most clinical chemistry systems including Immunoassay systems BXC0695C 2 x 1ml	
UnConjugated Estriol, Inhibin-A, PAPP-A, AMH	2-8°C
For screening of Down's syndrome & Spina Bifida compatible for use on most clinical chemistry systems Maternal Screening Control Tri-Level AFP, F-beta-hCG, Total hCG, UnConjugated Estriol, Inhibin-A, PAPP-A, AMH	2-8°C
Methanol Calibrator LS BXC0497A 1 x 2ml	2-8°C
Methanol Control (Level 1)LSBXC0498A1 x 2ml	2-8°C
Methanol Control (Level 2)LSBXC0499A1 x 2ml	2-8°C
Microalbumin Calibrator Series LS BXC0329A 5 x 1ml	2-8°C
Microalbumin Control LS BXC0328A 2 x 1ml	2-8°C
Microalbumin Control LS BXC0328B 5 x 1ml	2-8°C
Myoglobin Calibrator Series LS BXC0486A 5 x 1 x 0.5ml	2-8°C
Myoglobin Control LS BXC0487A 1 x 0.5ml	2-8°C
NTProBNP Calibrator LS BXC0335A 1 x 1ml	2-8°C
NTProBNP Control Series (6 Levels) LS BXC0336A 6 x 1 x 1ml	2-8°C
Oxalate (Urinary) Calibrator 0.5 mmol/L LS BXC0147A 1 x 5ml	2-8°C
Oxalate (Urinary) Control (Level 1) 0.2 mmol/L LS BXC0148A 1 x 5ml	2-8°C
Oxalate (Urinary) Control (Level 2)0.7 mmol/LLSBXC0149A1 x 5ml	2-8°C
Paediatric Control LS BXC0807A 3 x 1ml	2-8°C
PAPP-A & f-beta-hCG Controls (Level 1) LS BXC0337A 1 x 0.5ml	2-8°C
PAPP-A & f-beta-hCG Controls (Level 2) LS BXC0338A 1 x 0.5ml	2-8°C
PAPP-A & f-beta-hCG Controls (Level 3) LS BXC0339A 1 x 0.5ml	2-8°C
PAPP-A Control LS BXC0240A 1 x 0.5ml	2-8°C
Plasma Calibrator For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG125A 5 x 1ml	2-8°C
Plasma Control (Level 1 & Level 2) For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG112A 2 x 5 x 1ml	2-8°C
Plasma Control (Level 1) For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG108A 1 x 1ml	2-8°C
Plasma Control (Level 1) For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG108B 5 x 1ml	2-8°C
Plasma Control (Level 2) For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG109A 1 x 1ml	2-8°C
Plasma Control (Level 2) For Coagulation Series only, PT, aPTT, Fibrinogen, Thrombin Time LS COAG109B 5 x 1ml	2-8°C
Procalcitonin Control Set (2 Levels)LSBXC0590B2x1x1ml	2-8°C
Protein Control (Level 1)Liquid Ready to UseLSBXC0641A1 x 2ml	2-8°C
Protein Control (Level 2) Liquid Ready to Use LS BXC0642A 1 x 2ml	2-8°C
Protein Control Set (Level 1 & Level 2) Liquid Ready to Use LS BXC0643A 2 x 1 x 2ml	2-8°C
Retinol Binding Protein Control LS BXC0995A 1 x 1ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
RF (Multi-Point) Calibrator	Immunoturbidimetric	LS	BXC0612A	1 x 2ml	2-8°C
RF (Multi-Point) Calibrator	Immunoturbidimetric	LS	BXC0612B	3 x 2ml	2-8°C
Serum Indices (LIH)	Lyophilised for Beckman Coulter Series	Lyo	BXC0599E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Beckman Coulter Series	Lyo	BXC0599D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Beckman Coulter Series	Lyo	BXC0599C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Beckman Coulter Series	Lyo	BXC0599A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Beckman Coulter Series	LF	BXC0600E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Beckman Coulter Series	LF	BXC0600D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Beckman Coulter Series	LF	BXC0600C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Beckman Coulter Series	LF	BXC0600B	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyopholised for Roche Series	Lyo	BXC0604E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyopholised for Roche Series	Lyo	BXC0604D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyopholised for Roche Series	Lyo	BXC0604C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyopholised for Roche Series	Lyo	BXC0604B	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Roche Series	LF	BXC0605E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Roche Series	LF	BXC0605D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Roche Series	LF	BXC0605C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Roche Series	LF	BXC0605A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Abbott Series	Lyo	BXC0606E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Abbott Series	Lyo	BXC0606D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Abbott Series	Lyo	BXC0606C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Abbott Series	Lyo	BXC0606B	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Abbott Series	LF	BXC0607E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Abbott Series	LF	BXC0607D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Abbott Series	LF	BXC0607C	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Abbott Series	LF	BXC0607A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Vitros Series	Lyo	BXC0608F	3 x 4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Vitros Series	Lyo	BXC0608E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Lyophilised for Vitros Series	Lyo	BXC0608D	4 x 5 ml	2-8°C
Serum Indices (LIH)	5 1	-			2-8°C
	Lyophilised for Vitros Series	Lyo	BXC0608C	4 x 5 ml	
Serum Indices (LIH) Serum Indices (LIH)	Liquid Frozen for Vitros Series	LF	BXC0609F	3 x 4 x 5 ml	2-8°C
	Liquid Frozen for Vitros Series	LF	BXC0609E	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Vitros Series	LF	BXC0609D	4 x 5 ml	2-8°C
Serum Indices (LIH)	Liquid Frozen for Vitros Series	LF	BXC0609B	4 x 5 ml	2-8°C
Serum Indices (LIH) Serum Indices (LIHN)	Liquid Frozen for Vitros Series Lyophilised for Beckman Coulter	LF Lyo	BXC0609A BXC0599F	4 x 1 x 5 ml 3 x 4 x 5 ml	2-8°C 2-8°C
Serum Indices (LIHN)	Series Lyophilised for Beckman Coulter	Lyo	BXC0599F	4 x 5 ml	2-8°C
Serum Indices (LIHN)	Series Liquid Frozen for Beckman Coulter	LF	BXC0600F	4 x 5 ml	2-8°C
Serum Indices (LIHN)	Series Liquid Frozen for Beckman Coulter	LF	BXC0600P	4 x 1 x 5 ml	2-8°C
	Series				
Serum Indices (LIHN)	Lyopholised for Roche Series	Lyo	BXC0604F	3 x 4 x 5 ml	2-8°C
Serum Indices (LIHN)	Lyopholised for Roche Series	Lyo	BXC0604A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIHN)	Liquid Frozen for Roche Series	LF	BXC0605F	3 x 4 x 5 ml	2-8°C
Serum Indices (LIHN)	Liquid Frozen for Roche Series	LF	BXC0605B	4 x 5 ml	2-8°C
Serum Indices (LIHN)	Lyophilised for Abbott Series	Lyo	BXC0606F	3 x 4 x 5 ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
Serum Indices (LIHN)	Lyophilised for Abbott Series	Lyo F	BXC0606A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIHN)	Liquid Frozen for Abbott Series	LF	BXC0607F	3 x 4 x 5 ml	2-8°C
Serum Indices (LIHN)	Liquid Frozen for Abbott Series	LF	BXC0607B	4 x 5 ml	2-8°C
Serum Indices (LIHN)	Lyophilised for Vitros Series	Lyo	BXC0608B	4 x 5 ml	2-8°C
Serum Indices (LIHN)	Lyophilised for Vitros Series	Lyo	BXC0608A	4 x 1 x 5 ml	2-8°C
Serum Indices (LIHN)	Liquid Frozen for Vitros Series	LF	BXC0609C	4 x 5 ml	2-8°C
Specific Protein Calibrator	Liquid Ready to Use	LS	BXC0644A	1 x 2ml	2-8°C
Specific Protein Calibrator Set	Liquid Ready to Use	LS	BXC0646A	6 x 1 x 1ml	2-8°C
Superoxide Dismutase Control		LS	BXC0433A	5 x 1ml	2-8°C
Syphilis Control Panel (6 Levels)		LS	BXC0806A	6 x 1 x 0.5ml	2-8°C
Thalassaemia (Alpha & Beta) Control (Level 2)	For use on all major systems & Methods including HPLC & Immunoassay , HbA2, HbF, HbS	LS	BXC0665B	2 x 0.5ml	2-8°C
Thalassaemia (Alpha & Beta) Control (Level 1)	For use on all major systems & Methods including HPLC & Immunoassay , HbA2, HbF, HbS	LS	BXC0665A	2 x 0.5ml	2-8°C
Therapeutic Drug Monitoring (Level 1)		LS	BXC0781A	5 x 5ml	2-8°C
Therapeutic Drug Monitoring (Level 2)		LS	BXC0782A	5 x 5ml	2-8°C
Therapeutic Drug Monitoring (Level 3)		LS	BXC0783A	5 x 5ml	2-8°C
TIBC Calibrator (Single Level)		Lyo	BXC0238A	2 x 1ml	2-8°C
Torch Control Negative	For use on all major systems	LS	BXC0799A	3 x 1ml	2-8°C
Torch IgG Control Positive	For use on all major systems	LS	BXC0799B	3 x 1ml	2-8°C
Torch IgM Control Positive	For use on all major systems	LS	BXC0799C	5 x 1 x 1ml	2-8°C
Total Antioxidant Status (TAS) Calibrator		LS	BXC0555A	1 x 1ml	2-8°C
Total Antioxidant Status (TAS) Control		LS	BXC0554A	10 x 5ml	2-8°C
Total Galactose (Control Set) (Level 1 & Level 2)		LS	BXC0651A	2 x 1 x 1ml	2-8°C
Total Galactose (Level 1)		LS	BXC0652A	1 x 1ml	2-8°C
Total Galactose (Level 2)		LS	BXC0653A	1 x 1ml	2-8°C
Total Galactose Calibrator		LS	BXC0654A	1 x 1ml	2-8°C
Triage Control (2 Levels)		LS	BXC0880A	2 x 1 x 0.5ml	2-8°C
Troponin-I Calibrator Set	For use on all major systems	LS	BXC0473A	6 x 1 x 1ml	2-8°C
Troponin-I Control Set	For use on all major systems	LS	BXC0470A	2 x 1 x 1ml	2-8°C
Troponin-T High Sensitivity Control	For use on all major systems	LS	BXC0457A	3 x 1ml	2-8°C
Tumour Marker Control (Level 1)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0792A	5 x 2ml	2-8°C
Tumour Marker Control (Level 1)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0794A	5 x 2ml	2-8°C
Tumour Marker Control (Level 2)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0792B	5 x 2ml	2-8°C
Tumour Marker Control (Level 2)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0795A	5 x 2ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
Tumour Marker Control (Level 3)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0792C	5 x 2ml	2-8°C
Tumour Marker Control (Level 3)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0796A	5 x 2ml	2-8°C
Tumour Marker Control (Tri-Level)	For use on all major systems & Methods including Immunoassay, AFP, CEA, CA-125, CA-19.9, CA-15- 3, Ferritin, Beta-2-Microglobulin, FPSA, PSA, F-beta-hCG, Thyroglobulin.	LS	BXC0793A	3 x 1 x 2ml	2-8°C
Urine Assayed Control (Level 1)		LS	BXC0661A	10 x 10ml	2-8°C
Urine Assayed Control (Level 2)		LS	BXC0661B	10 x 10ml	2-8°C
Urine Precision Control (Level 1)		LS	BXC0662A	10 x 10ml	2-8°C
Urine Precision Control (Level 2)		LS	BXC0662B	10 x 10ml	2-8°C
Urine Strip Control (Level 1 & Level 2)	For Urine and hCG Strips Only	LS	BXC0663A	2 x 3 x 12ml	2-8°C
Vitamin-D Calibrator Set		LS	BXC0475A	5 x 1ml	2-8°C
Vitamin-D Control Set		LS	BXC0474A	2 x 1ml	2-8°C
Zinc & Copper Calibrator		Lyo	BXC0463A	2 x 1ml	2-8°C

Aqueous Standards

Aqueous standards play a critical role in diagnostics by serving as reference solutions with known concentrations of analytes, which are essential for calibrating diagnostic instruments and validating test accuracy. These standards ensure consistency and reliability in diagnostic results by providing a benchmark for comparison, enabling laboratories to detect and quantify substances in biological samples accurately.

Their use minimises variability, ensures compliance with regulatory requirements, and enhances the reproducibility of diagnostic tests, leading to more reliable patient outcomes and improved clinical decision-making. Fortress Diagnostics Aqueous Standards are available in a liquid stable format for convenience and ease-of-use, with a long shelf life, and are traceable to reference standard material.

Description	Details	Format	Cat No.	Size	Storage
Albumin	45 g/l (4.5g/dL)	LS	BXCSTD01	10 x 5ml	2-8°C
Calcium	Referenced against NIST 909b (2.5mmol/l10mg/dL)	LS	BXCSTD02	10 x 5ml	2-8°C
Chloride	Referenced against NIST 909b (75mmol/l)	LS	BXCSTD03	10 x 5ml	2-8°C
Chloride	Referenced against NIST 909b (100mmol/l)	LS	BXCSTD04	10 x 5ml	2-8°C
Cholesterol	Referenced against NIST 909b (5.17mmol/l200mg/dL)	LS	BXCSTD05	10 x 5ml	2-8°C
Creatinine	Referenced against NIST 909b (177 µmol/l2 mg/dL)	LS	BXCSTD06	10 x 5ml	2-8°C
Creatinine/Glucose/Urea Standard	Referenced against NIST 965a / 909b (5.55 mmol/l100mg/dL/ 177 µmol/l / 8.33mmol/l)	LS	BXCSTD24	10 x 5ml	2-8°C
Glucose	Referenced against NIST 965a (5.55 mmol/l100mg/dL)	LS	BXCSTD07	10 x 5ml	2-8°C
Glucose-Fructose Standard		LS	BXCSTD26	ТВС	2-8°C
Haemoglobin Standard	For Drabkin's Method (18g/dL)	LS	BXCSTD08	5 x 10ml	2-8°C
Haemoglobin Standard Set	For Drabkin's Method (8, 10, 12, 15 & 18g/dL)	LS	BXCSTD09	5 x 1 x 10ml	2-8°C
Iron	35.8 µmol/l (0.2 mg/dL)	LS	BXCSTD10	10 x 5ml	2-8°C
Iron	107.4 µmol/l (0.6 mg/dL)	LS	BXCSTD11	10 x 5ml	2-8°C
Lactate	4.44mmol/l (40mg/dL)	LS	BXCSTD12	10 x 5ml	2-8°C

Description	Details	Format	Cat No.	Size	Storage
Magnesium	Referenced against NIST 909b (1.0 mmol/l2.43mg/dL)	LS	BXCSTD13	10 x 5ml	2-8°C
Phosphorus	1.61 mmol/l (5 mg/dL)	LS	BXCSTD14	10 x 5ml	2-8°C
Potassium	Referenced against NIST 909b (0.5 mmol/l)	LS	BXCSTD15	10 x 5ml	2-8°C
Potassium	Referenced against NIST 909b (5 mmol/l)	LS	BXCSTD16	10 x 5ml	2-8°C
Pyruvate	2 mmol/l	LS	BXCSTD17	10 x 5ml	2-8°C
Sodium	Referenced against NIST 909b (150.0 mmol/l150 mEq/l)	LS	BXCSTD18	10 x 5ml	2-8°C
Total Protein	Referenced against NIST 927d (60g/l (6.0 g/dL)	LS	BXCSTD19	10 x 5ml	2-8°C
Triglycerides	Referenced against NIST 909b (2.28 mmol/l200mg/dL)	LS	BXCSTD20	10 x 5ml	2-8°C
Urea	Referenced against NIST 909b (8.33 mmol/l50mg/dL)	LS	BXCSTD21	10 x 5ml	2-8°C
Uric Acid	Referenced against NIST 909b (595 µmol/l10mg/dL)	LS	BXCSTD22	10 x 5ml	2-8°C
Urinary Copper Standard		LS	BXCSTD25	10 x 5ml	2-8°C
Urinary/CSF Protein Standard	Referenced against NIST 927d (1g/l)	LS	BXCSTD23	10 x 5ml	2-8°C





Immunoassays (ELISA)

Enzyme-Linked Immunosorbent Assay (ELISA) is a widely used technique in diagnostics for detecting and quantifying specific proteins, antibodies, or antigens in a sample. It plays a crucial role in diagnosing infections, monitoring disease progression, and assessing immune responses.

ELISA works by using enzyme-linked antibodies to bind to target molecules, generating a measurable signal, typically color change, that indicates the presence and concentration of the target. The significance of ELISA lies in its high sensitivity, allowing the detection of very low concentrations of biomolecules, and its specificity, which is achieved through the use of highly specific antigen-antibody interactions. Its versatility comes from the ability to adapt the assay to detect various types of molecules, including hormones, peptides, proteins, and pathogens. ELISA is widely used in diagnostic tests for detecting infectious diseases like HIV, Hepatitis, and COVID-19, as well as in immunological studies to quantify cytokines, antibodies, or antigens, making it invaluable in both clinical and research settings.

Key Features & Benefits

ELISA (Enzyme-Linked Immunosorbent Assay) offers several benefits that make it a widely used technique in various fields, including medical diagnostics, research, and biotechnology:



High Sensitivity: ELISA can detect target substances at very low concentrations. This is crucial for detecting trace amounts of analytes.



Quantitative and Qualitative Analysis: By measuring the intensity of the signal produced, ELISA allows for accurate quantification of the target substance in the sample.



High Throughput: ELISA can be performed in multiwell plates, allowing for the simultaneous analysis of multiple samples.



Stability: ELISA reagents, particularly antibodies and enzyme conjugates, are generally stable when stored properly.

High Performance ELISA Kits

Category	Description	Range	Incubation Time	Sensitivity
Allergy	IgE	0-400 IU/ml	30 + 30 + 15 min	1.00 IIU/ml
Anaemia	Ferritin	1 -50,000 ng/ml	30 + 30 + 15 min	1 ng/ml
Autoimmune	Anti-TG	25.2 - 124.7 IU/ml	60 + 30 + 15 min	1.94 IU/ml
	Anti-TPO	10.8 - 39.2 IU/ml	60 + 30 + 15 min	0.92 IU/ml
Cardiac Markers	CRP (High Sensetivity)	0.014 - 5,000 ug/ml	15 + 15 min	0.014 ug/ml
	Digoxin	0-4 ng/ml	30 + 15 min	0.072 ng/ml
	Troponin-I	0.05-10,000 ng/ml	15 + 15 min	0.05 ng/ml
Diabetes	C-Peptide	0 - 0.82 ng/ml	120 + 15 min	0.02 ng/ml
	Insulin	0-300 µlU/ml	120 + 15 min	0.75 uIU/ml
Fertility & Prenatal	LH	0.0 - 200 mIU/ml	60 + 15 min	0.8 mIU/ml
	FSH	0.0 - 100 mg/dL	60 + 15 min	0.8 mIU/ml
	Prolactin	0.0 - 100 ng/ml	60 + 15 min	0.8 ng/ml
Growth Deficiency	HGH	0-150 U/ml	60 + 15 min	0.025 uIU/ml
Maternal Screening	Down's Syndrome	Screening (during first trim	ester of pregnancy 9th - 13th	week)
	PAPP-A	0.120- 320 ug/ml	30 + 30 + 20 min	0.12 ug/ml
	Free Beta-hCG	0-200 ng/dl	30 + 30 + 20 min	0.05 ng/ml
	Total hCG	0-250 mlU/ml	60 + 15 min	0.02 mIU/ml
	Down's Syndrome Sc	reening (during second trin	nester of pregnancy 14th - 24	:h week).
	AFP	0.025 -10,000 ng/ml	60 + 15 min	0.025 ng/ml
	Unconjugated Estriol (uE3)	0-30 ng/ml	60 + 15 min	2.9 pg/T
	Free Beta-hCG	0-200 ng/dl	30 + 30+ 20 min	0.05 ng/ml
	Inhibin-A	0-1000 pg/ml	180 + 60+ 15 min	5.0 pg/ml

Immunoassays (ELISA) | SECTION 3

Category	Description	Range	Incubation Time	Sensitivity
Steroid Panel	Progesterone	0.0 - 60.0 IU/ml	60 + 15 min	0.105 ng/ml
	Testosterone	0.0 - 12.0 mg/ml	60 + 15 min	0.038 ng/ml
	Estradiol	0.0 - 3000 IU/ml	60 + 15 min	6.5 pg/ml
	Free testosterone	0-34 pg/ml	60 + 15 min	0.02 pg/ml
	DHEA-S	0.0 - 8.0 mg/ml	60 + 15 min	0.035 mg/ml
	17Alpha-OH Progesterone	0.0 - 10.0 ng/ml	60 + 15 min	0.03 ng/ml
	Cortisol	0.0 - 50.0 mg/dl	60 + 15 min	0.25 mg/dL
Thyroid & Adrenal	Т3	0.0 - 7.5 ng/ml	60 + 15 min	0.04 ng/ml
	Τ4	0.0 - 25 mg/dl	60 + 15 min	0.4 mg/dL
	тѕн	0.0 - 40 mIU/ml	60 + 15 min	0.078 mIU/ml
	FT3	0.0 - 16 pg/ml	60 + 15 min	0.05 pg/ml
	FT4	0.0 - 2.7 pg/ml	60 + 15 min	0.05 ng/dL
	TG	0.0 - 2000 IU/ml	60 + 15 min	5.0 IU/ml dL
	ТРО	0.0 - 500 IU/ml	60 + 15 min	1.5 IU/ml
	T3-Uptake	18.0 - 47.0%	60 + 15 min	-
	NT4	0.0 - 25 mg/dL	16 - 20 Hours	0.5 mg/dL
	NTSH	0.0 - 250 mIU/ml	16 - 20 Hours	1.0 mIU/ml
Fumour Markers	AFP	0.025 -10,000 ng/ml	60 + 15 min	0.025 ng/ml
	CA 15.3	0-400 U/ml	60 + 60 + 20 min	0.2 U/ml
	CA 125	0-400 IU/ml	60 + 15 min	1 U/ml
	CA 19.9	0-500 U/ml	60 + 15 min	1 U/ml
	CEA	0.025 - 60,000 ng/ml	60 + 15 min	0.025 ng/ml
	fPSA	0 - 10 ng/ml	60 + 15 min	0.052 ng/ml
	PSA	0 - 100 ng/dl	30 + 15 min	0.012 ng/ml
√itamin-D	Vitamin-D	0 - 150 ng/ml	90 + 30 + 30 min	0.6 ng/ml

Description	Details	Method	Cat No.	Size	Storage
17Alpha-OH Progesterone		ELISA	BXE0996A	96T	2-8°C
AFP		ELISA	BXE0801A	96T	2-8°C
AFP		ELISA	BXE0801B	192T	2-8°C
Anti-HBc		ELISA	BXE0761A	96T	2-8°C
Anti-HBe		ELISA	BXE0772A	96T	2-8°C
Anti-HBs (Quantitative)		ELISA	BXE0752A	96T	2-8°C
Anti-Mullerian Hormone (AMH)		ELISA	BXE0999A	96T	2-8°C
Anti-Mullerian Hormone (AMH) BC		ELISA	BXE0989A	96T	2-8°C
Anti-Tg		ELISA	BXE0895A	96T	2-8°C
Anti-TPO		ELISA	BXE0896A	96T	2-8°C
CA125		ELISA	BXE0821A	96T	2-8°C
CA15-3		ELISA	BXE0881A	96T	2-8°C
CA19.9		ELISA	BXE0841A	96T	2-8°C
CEA		ELISA	BXE0811A	96T	2-8°C
Cortisol		ELISA	BXE0831A	96T	2-8°C
COVID-19 IgM Antibody		ELISA	BXE0883A	96T	2-8°C
COVID-19 Neutralising Antibody		ELISA	BXE0885A	96T	2-8°C
COVID-19 Total Antibody		ELISA	BXE0882A	96T	2-8°C
C-Peptide		ELISA	BXE0616A	96T	2-8°C
CRP (High Sensitivity)		ELISA	BXE0617A	96T	2-8°C
Cytomegalo Virus (CMV) IgG		ELISA	BXE0880A	96T	2-8°C

SECTION 3 | Immunoassays (ELISA)

Description	Details	Method	Cat No.	Size	Storage
Cytomegalo Virus (CMV) IgM		ELISA	BXE0890A	96T	2-8°C
Dengue IgG		ELISA	BXE0695A	96T	2-8°C
Dengue IgM		ELISA	BXE0696A	96T	2-8°C
DHEA-S		ELISA	BXE0998A	96T	2-8°C
Digoxin		ELISA	BXE0992A	96T	2-8°C
Estradiol		ELISA	BXE0860A	96T	2-8°C
Estradiol		ELISA	BXE0860B	192T	2-8°C
Ferritin		ELISA	BXE0891A	96T	2-8°C
Folate		ELISA	BXE0892A	96T	2-8°C
Free PSA		ELISA	BXE0861A	96T	2-8°C
Free Testosterone		ELISA	BXE0997A	96T	2-8°C
Free BetahCG		ELISA	BXE0872A	96T	2-8°C
FSH		ELISA	BXE0631A	96T	2-8°C
FSH		ELISA	BXE0631B	192T	2-8°C
FSH		ELISA	BXE0631C	480T	2-8°C
FT3		ELISA	BXE0731A	96T	2-8°C
FT4		ELISA	BXE0721A	96T	2-8°C
HAV.IgG		ELISA	BXE0601A	96T	2-8°C
HAV.IgM		ELISA	BXE0602A	96T	2-8°C
HBc.lgM		ELISA	BXE0603A	96T	2-8°C
- HBeAg		ELISA	BXE0771A	96T	2-8°C
HbsAg		ELISA	BXE0741A	96T	2-8°C
HbsAg		ELISA	BXE0741B	192T	2-8°C
HbsAg		ELISA	BXE0741C	480T	2-8°C
- HbsAg (High Sensitivity)		ELISA	BXE0742A	96T	2-8°C
HbsAg (High Sensitivity)		ELISA	BXE0742B	192T	2-8°C
HbsAg (High Sensitivity)		ELISA	BXE0742C	480T	2-8°C
HbsAg (High Sensitivity) with CE		ELISA	BXE0743A	96T	2-8°C
HbsAg (High Sensitivity) with CE		ELISA	BXE0743C	480T	2-8°C
hCG		ELISA	BXE0871A	96T	2-8°C
hCG		ELISA	BXE0871B	192T	2-8°C
hCG Rapid		ELISA	BXE0873A	96T	2-8°C
hCG Rapid		ELISA	BXE0873B	192T	2-8°C
hCG Rapid Extended Range		ELISA	BXE0874A	96T	2-8°C
hCG Rapid Extended Range		ELISA	BXE0874B	192T	2-8°C
нсу		ELISA	BXE0781A	96T	2-8°C
нсу		ELISA	BXE0781B	192T	2-8°C
нсу		ELISA	BXE0781C	480T	2-8°C
HCV High Sensitivity with CE		ELISA	BXE0783A	96T	2-8°C
HCV High Sensitivity with CE		ELISA	BXE0783C	480T	2-8°C
HDV.Ag		ELISA	BXE0605A	96T	2-8°C
HDV.IgG		ELISA	BXE0606A	96T	2-8°C
HDV.IgM		ELISA	BXE0607A	96T	2-8°C
Helicobacter Pylori, IgA		ELISA	BXE0672A	96T	2-8°C
Helicobacter Pylori, IgG		ELISA	BXE0673A	96T	2-8°C
Helicobacter Pylori, IgM		ELISA	BXE0674A	96T	2-8°C
Herpes Simplex Virus (HSV-I/2), IgG		ELISA	BXE0621A	96T	2-8°C
Herpes Simplex Virus (HSV-I/2), IgM		ELISA	BXE0622A	96T	2-8°C
		LLIJA	DALUUZZA	501	2-0 C

Immunoassays (ELISA) | SECTION 3

Description	Details	Method	Cat No.	Size	Storage
Herpes Simplex Virus-I (HSV-I), IgG		ELISA	BXE0611A	96T	2-8°C
Herpes Simplex Virus-II (HSV-II), IgG		ELISA	BXE0613A	96T	2-8°C
Herpes Simplex Virus-II (HSV-II), IgM		ELISA	BXE0614A	96T	2-8°C
HEV Ab		ELISA	BXE0745A	96T	2-8°C
HEV Ag		ELISA	BXE0903A	96T	2-8°C
HEV Ag Plus		ELISA	BXE0904A	96T	2-8°C
HEV IgG		ELISA	BXE0901A	96T	2-8°C
HEV IgM		ELISA	BXE0902A	96T	2-8°C
HGV.lgG		ELISA	BXE0608A	96T	2-8°C
HIV 1+2 3rd gen		ELISA	BXE0791A	96T	2-8°C
HIV 1+2 3rd gen		ELISA	BXE0791B	192T	2-8°C
HIV 1+2 3rd gen		ELISA	BXE0791C	480T	2-8°C
HIV 1+2 3rd Gen. (with CE)		ELISA	BXE0793A	96T	2-8°C
HIV 1+2 3rd Gen. (with CE)		ELISA	BXE0793C	480T	2-8°C
HIV Ag/Ab 4th gen		ELISA	BXE0792A	96T	2-8°C
HIV Ag/Ab 4th gen		ELISA	BXE0792B	192T	2-8°C
HIV Ag/Ab 4th gen. (with CE)		ELISA	BXE0794A	96T	2-8°C
HIV Ag/Ab 4th gen. (with CE)		ELISA	BXE0794C	480T	2-8°C
HTLV 1+2 ELISA		ELISA	BXE0910A	96T	2-8°C
Human Growth Hormone (HGH)		ELISA	BXE0991A	96T	2-8°C
lgE		ELISA	BXE0641A	96T	2-8°C
IL-6		ELISA	BXE0960A	96T	2-8°C
Inhibin-A		ELISA	BXE0615A	96T	2-8°C
Inhibin-B		ELISA	BXE0618A	96T	2-8°C
Insulin		ELISA	BXE0610A	96T	2-8°C
Leptin		ELISA	BXE0962A	96T	2-8°C
LH		ELISA	BXE0651A	96T	2-8°C
LH		ELISA	BXE0651B	192T	2-8°C
LH		ELISA	BXE0651C	480T	2-8°C
Malaria		ELISA	BXE0691A	96T	2-8°C
МСМ5		ELISA	BXE0950A	96T	2-8°C
Myoglobin		ELISA	BXE0609A	96T	2-8°C
Neonatal T4		ELISA	BXE0863A	96T	2-8°C
Neonatal TSH		ELISA	BXE0682A	96T	2-8°C
PAPP-A		ELISA	BXE0888A	96T	2-8°C
Patient Cards for NTSH & NT4	6 Specimen per Paper Card	ELISA	BXEPC100	SET OF 100	2-8°C
Progesterone		ELISA	BXE0661A	96T	2-8°C
Progesterone		ELISA	BXE0661B	192T	2-8°C
Prolactin		ELISA	BXE0671A	96T	2-8°C
Prolactin		ELISA	BXE0671B	192T	2-8°C
Prolactin		ELISA	BXE0671C	480T	2-8°C
PSA		ELISA	BXE0851A	96T	2-8°C
PSA		ELISA	BXE0851B	192T	2-8°C
PSA (High Sensitivity)		ELISA	BXE0852A	96T	2-8°C
PSA (High Sensitivity)		ELISA	BXE0852B	192T	2-8°C
PTH Intact (Parthormone)		ELISA	BXE0684A	96T	2-8°C
Rubella, IgG		ELISA	BXE0685A	96T	2-8°C
Rubella, IgM		ELISA	BXE0686A	96T	2-8°C

SECTION 3 | Immunoassays (ELISA)

Syphilis ELISAELISABXE0995B192T2-8'CSyphilis ELISAELISABXE0995C480T2-8'CT3ELISABXE0701A96T2-8'CT3ELISABXE0701B192T2-8'CT3StreptavidinELISABXE0703A96T2-8'CT3 StreptavidinELISABXE0703A96T2-8'CT3-UptakeELISABXE0702A96T2-8'CT3-UptakeELISABXE0702A96T2-8'CT4ELISABXE0702A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE071A96T2-8'CT4ELISABXE0864A96T2-8'CT5TestosteroneELISABXE0864A96T2-8'CT5HELISABXE0864A96T2-8'CT5HELISABXE0864A96T2-8'CT5HELISABXE0864A96T2-8'CT5HELISABXE0864A96T2-8'CT5HELISABXE0864A96T2-8'CT5H	Description	Details	Method	Cat No.	Size	Storage
Symplitis ELISAELISABXE 0995C480T2.8°CT3ELISABXE0701A96T2.8°CT3ELISABXE0701B192T2.8°CT3ELISABXE0703A96T2.8°CT3SteptavidinELISABXE0703B96T2.8°CT3 StreptavidinELISABXE0703B96T2.8°CT3-UptakeELISABXE0702B96T2.8°CT3-UptakeELISABXE0702B96T2.8°CT3-UptakeELISABXE0702B96T2.8°CT4ELISABXE0702B96T2.8°CT4ELISABXE071LB96T2.8°CT4ELISABXE071LB96T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT4ELISABXE0712B192T2.8°CT5ELISABXE0682A96T2.8°CT5HELISABXE0684A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681B192T2.8°CT5HELISABXE0681B192T2.8°CT5HELISA <th>Syphilis ELISA</th> <th></th> <th>ELISA</th> <th>BXE0995A</th> <th>96T</th> <th>2-8°C</th>	Syphilis ELISA		ELISA	BXE0995A	96T	2-8°C
TaELISABXE0701A96T2-8°CT3ELISABXE0701B192T2-8°CT3ELISABXE0701C480T2-8°CT3 StreptavidinELISABXE0703A96T2-8°CT3 StreptavidinELISABXE0703A96T2-8°CT3 UptakeELISABXE0703B192T2-8°CT3 UptakeELISABXE0702A96T2-8°CT3 UptakeELISABXE0702B192T2-8°CT4ELISABXE0711B96T2-8°CT4ELISABXE0711B96T2-8°CT4ELISABXE0712B96T2-8°CT4ELISABXE0712B192T2-8°CT4ELISABXE0712B192T2-8°CT4ELISABXE0712B192T2-8°CT4ELISABXE0712B192T2-8°CT4Streptavidin96T2-8°CT5ELISABXE0828192T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T2-8°CT5HELISABXE0834A96T	Syphilis ELISA		ELISA	BXE0995B	192T	2-8°C
T3ElISABKE0701B19272.8°CT3ELISABKE0701C48072.8°CT3 StreptavidinELISABKE0703B19272.8°CT3 StreptavidinELISABKE0703B19272.8°CT3 UptakeELISABKE0702B19272.8°CT3 UptakeELISABKE0702B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4ELISABKE071B19272.8°CT4SteptavidinELISABKE071B19272.8°CT4SteptavidinELISABKE071B19272.8°CT5ELISABKE082A9672.8°C2.8°CT5ELISABKE082B19272.8°C2.8°CT5ELISABKE084B9672.8°C2.8°CT5ELISABKE084B9672.8°C2.8°CT5ELISABKE084B9672.8°C2.8°CT5ELISABKE084B9672.8°C2.8°CT5ELISABKE084B9672.8°C2.8°CT5ELISABKE084B9672.8°C2.8°C <th>Syphilis ELISA</th> <th></th> <th>ELISA</th> <th>BXE0995C</th> <th>480T</th> <th>2-8°C</th>	Syphilis ELISA		ELISA	BXE0995C	480T	2-8°C
T3ELISABKE0701C480T2.8°CT3 StreptavidinELISABKE0703A9CT2.8°CT3 StreptavidinELISABKE0703B192T2.8°CT3-UptakeELISABKE0702B9CT2.8°CT3-UptakeELISABKE0702B192T2.8°CT4ELISABKE0702B192T2.8°CT4ELISABKE0711B9CT2.8°CT4ELISABKE0711B192T2.8°CT4ELISABKE0711B192T2.8°CT4ELISABKE0711B192T2.8°CT4ELISABKE0711B192T2.8°CT4StreptavidinELISABKE0711B192T2.8°CT4 StreptavidinELISABKE0712B192T2.8°CT5 SteptavidinELISABKE0712B192T2.8°CT6 stosteroneELISABKE082A9CT2.8°CToxoplasma gondii, 19GELISABKE0862A9CT2.8°CT5HELISABKE0681A9CT2.8°CT5HELISABKE0681A9CT2.8°CT5HELISABKE0681B192T2.8°CT5H RapidELISABKE0681B192T2.8°CUtconjugated Estriol (UE3)ELISABKE0681B9CT2.8°CVitamin-D [Blood Spots)ELISABKE08649CT2.8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABKE08639CT2.8°CVitamin-D Fast (90 Min)	тз		ELISA	BXE0701A	96T	2-8°C
T3 StreptavidinELISA8XE0703A9cT2.8°CT3 StreptavidinELISA8XE0702B192T2.8°CT3 UptakeELISA8XE0702B192T2.8°CT3 UptakeELISA8XE0702B192T2.8°CT4ELISA8XE071IB96T2.8°CT4ELISA8XE071IB192T2.8°CT4ELISA8XE0712B480T2.8°CT4 StreptavidinELISA8XE0712B96T2.8°CT4 StreptavidinELISA8XE0712B192T2.8°CT4 StreptavidinELISA8XE0712B192T2.8°CT4 StreptavidinELISA8XE0712B192T2.8°CT5 StreptavidinELISA8XE0712B192T2.8°CT6 streptavidinELISA8XE0712B192T2.8°CT5 StreptavidinELISA8XE082B192T2.8°CT5 StreptavidinELISA8XE082B192T2.8°CT6 streptavidinELISA8XE082B192T2.8°CT6 streptavidinELISA8XE082B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°CT6 streptavidinELISA8XE083B192T2.8°C	тз		ELISA	BXE0701B	192T	2-8°C
T3 StreptavidinELISABXE0703B19272.8°CT3-UptakeELISABXE0702A96T2.8°CT3-UptakeELISABXE0702B19272.8°CT4ELISABXE071IA96T2.8°CT4ELISABXE071IB19272.8°CT4ELISABXE071IC480T2.8°CT4ELISABXE0712B19272.8°CT4 StreptavidinELISABXE0712B96T2.8°CT4 StreptavidinELISABXE0712B19272.8°CT4 StreptavidinELISABXE0712B19272.8°CT4 StreptavidinELISABXE062A96T2.8°CT6 StreptavidinELISABXE069A96T2.8°CT6 StreptavidinELISA	тз		ELISA	BXE0701C	480T	2-8°C
Ta-UptakeELISABXE0702A9F124°CTa-UptakeELISABXE0702B192724°CTaELISABXE0711A9F124°CTaELISABXE0711B192724°CTaELISABXE0712B9F124°CTa StreptavidinELISABXE0712B9F124°CTa StreptavidinELISABXE0712B9F124°CTa StreptavidinELISABXE072B9F124°CTa StreptavidinELISABXE062B9F124°CTa StreptavidinELISABXE06BA9F124°CTa StreptavidinELISABXE06BA9F124°CTa StreptavidinELISABXE06BA9F124°CTa StreptavidinELISABXE06BA9F124°CTa StreptavidinELISABXE06BA9F124°CTa StreptavidinELISABXE06BA </th <th>T3 Streptavidin</th> <th></th> <th>ELISA</th> <th>BXE0703A</th> <th>96T</th> <th>2-8°C</th>	T3 Streptavidin		ELISA	BXE0703A	96T	2-8°C
T3-bptakeELISABXE0702B192T2.9°CT4ELISABXE0711A96T2.8°CT4ELISABXE0711B192T2.8°CT4ELISABXE0711C480T2.8°CT4ELISABXE0712A96T2.8°CT4 StreptavidinELISABXE0712B192T2.8°CT4 StreptavidinELISABXE0712B192T2.8°CT4 StreptavidinELISABXE072B192T2.8°CT4 StreptavidinELISABXE082A96T2.8°CT5 StosteroneELISABXE0862B192T2.8°CToxoplasma gondii, IgGELISABXE069A96T2.8°CT5 MapiaELISABXE0683A96T2.8°CT5 HELISABXE0681B192T2.8°CT5 HELISABXE0681B192T2.8°CT5 H RapidELISABXE0681B96T2.8°CVitamin F12ELISABXE0683A96T2.8°CVitamin P5 at (00 Min)Liquid Stable (LS)ELISABXE011A96T2.8°CVitamin P Fast (00 Min)Liquid Stable (Lso)ELISABXE011A96T2.8°CVitamin P Fast (00 Min)Liquid Stable (Lso)	T3 Streptavidin		ELISA	BXE0703B	192T	2-8°C
T4ELISABXE0711A9F12-8°CT4ELISABXE0711B192T2-8°CT4ELISABXE0711C480T2-8°CT4 StreptavidinELISABXE0712B9F12-8°CT4 StreptavidinELISABXE0712B192T2-8°CTestosteroneELISABXE082B96T2-8°CToxoplasma gondii, IgGELISABXE082B192T2-8°CToxoplasma gondii, IgGELISABXE082B96T2-8°CToxoplasma gondii, IgMELISABXE082B96T2-8°CTsHELISABXE081B96T2-8°CT5HELISABXE081B96T2-8°CTsH RapidELISABXE082B96T2-8°CTsH RapidELISABXE082B96T2-8°CVitamin-D (Blood Spots)ELISABXE013A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE011A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE011A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE011A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE012A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE012A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISABXE012A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LSO)ELISA <td< th=""><th>T3-Uptake</th><th></th><th>ELISA</th><th>BXE0702A</th><th>96T</th><th>2-8°C</th></td<>	T3-Uptake		ELISA	BXE0702A	96T	2-8°C
T4ELISABXE0711B19272.8°CT4ELISABXE0711C48072.8°CT4 StreptavidinELISABXE0712B96T2.8°CT4 StreptavidinELISABXE0712B19272.8°CTestosteroneELISABXE082A96T2.8°CToxoplasma gondii, IgGELISABXE082B92T2.8°CToxoplasma gondii, IgMELISABXE082B96T2.8°CToxoplasma gondii, IgMELISABXE080A96T2.8°CToxoplasma gondii, IgMELISABXE081A96T2.8°CTshELISABXE081B92T2.8°CTshELISABXE081B96T2.8°CTshELISABXE081B96T2.8°CTsh RapidELISABXE081B96T2.8°CUnconjugated Estriol (UE3)ELISABLISABXE083A96T2.8°CVitamin-D fest (90 Min)Liquid Stable (Liyo)ELISABXE081A96T2.8°CVitamin-D Fast (90 Min)Liquid Stable (Liyo)ELISABXE011A96T2.8°CVitamin-D Fast (90 Min)Liquid	T3-Uptake		ELISA	BXE0702B	192T	2-8°C
T4ELISABXE0711C400T2.8°CT4 StreptavidinELISABXE0712A96T2.8°CT4 StreptavidinELISABXE0712B192T2.8°CTestosteroneELISABXE082A96T2.8°CTestosteroneELISABXE082B192T2.8°CToxoplasma gondii, IgGELISABXE082B96T2.8°CToxoplasma gondii, IgMELISABXE0680A96T2.8°CToponin-IELISABXE0681A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681B192T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0681A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°CT5HELISABXE0683A96T2.8°C <t< th=""><th>Т4</th><th></th><th>ELISA</th><th>BXE0711A</th><th>96T</th><th>2-8°C</th></t<>	Т4		ELISA	BXE0711A	96T	2-8°C
T4 StreptavidinELISABXE0712A96T2-8°CT4 StreptavidinELISABXE0712B192T2-8°CTestosteroneELISABXE082A96T2-8°CTestosteroneELISABXE082BA192T2-8°CToxoplasma gondii, IgGELISABXE089A96T2-8°CToxoplasma gondii, IgMELISABXE089A96T2-8°CTorponin-IELISABXE081A96T2-8°CT5HELISABXE081A96T2-8°CT5HELISABXE081A96T2-8°CT5HELISABXE081A96T2-8°CT5HELISABXE081A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5HELISABXE085A96T2-8°CT5H	Т4		ELISA	BXE0711B	192T	2-8°C
T4 StreptavidinELISABXE0712B192T2-8°CTestosteroneELISABXE0862A96T2-8°CTestosteroneELISABXE0882B192T2-8°CToxoplasma gondii, IgGELISABXE0689A96T2-8°CToxoplasma gondii, IgMELISABXE0689A96T2-8°CToponin-IELISABXE0681A96T2-8°CT5HELISABXE0681A96T2-8°CT5HELISABXE0681B192T2-8°CT5HELISABXE0681B192T2-8°CT5HELISABXE0681B96T2-8°CT5HELISABXE0681B96T2-8°CT5HELISABXE0681B96T2-8°CT5HELISABXE0681B96T2-8°CT5HELISABXE0681B96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT5HELISABXE0683A96T2-8°CT	Т4		ELISA	BXE0711C	480T	2-8°C
TestosteroneELISABXE0862A96T2-8°CTestosteroneELISABXE0862B192T2-8°CToxoplasma gondii, IgGELISABXE0689A96T2-8°CToxoplasma gondii, IgMELISABXE0690A96T2-8°CToponin-IELISABXE0864A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELISABXE0113A96T2-8°CTSHELI	T4 Streptavidin		ELISA	BXE0712A	96T	2-8°C
TestosteroneELISABXE0862B19272-8°CToxoplasma gondii, IgGELISABXE0689A96T2-8°CToxoplasma gondii, IgMELISABXE0690A96T2-8°CToponin-IELISABXE0861A96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0683A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A96T2-8°CTSHELISABXE085A<	T4 Streptavidin		ELISA	BXE0712B	192T	2-8°C
Toxoplasma gondii, IgGELISABXE0689A96T2-8°CToxoplasma gondii, IgMELISABXE0690A96T2-8°CTroponin-IELISABXE0864A96T2-8°CTSHELISABXE0681B96T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681C480T2-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE085A96T2-8°CVitamin-D (Blood Spots)ELISABXE0960A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°CUnconjugated Lstriol (UE3)Liquid Stable (LS)ELISABXE011A96T2-8°COther Hamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE011A96T2-8°COther Hamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE011A96T2-8°C	Testosterone		ELISA	BXE0862A	96T	2-8°C
Toxoplasma gondii, IgMELISABXE0690A96T2-8°CTroponin-IELISABXE0864A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681C480T2-8°CTSHELISABXE0683A96T2-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0863A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°C	Testosterone		ELISA	BXE0862B	192T	2-8°C
Troponin-IELISABXE0864A96T2-8°CTSHELISABXE0681A96T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681C480T2-8°CTSHELISABXE0683A96T2-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0112A96T2-8°C	Toxoplasma gondii, IgG		ELISA	BXE0689A	96T	2-8°C
TSHELISABXE0681A96T2-8°CTSHELISABXE0681B192T2-8°CTSHELISABXE0681C480T2-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin-D (Blood Spots)ELISABXE0960A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°C	Toxoplasma gondii, IgM		ELISA	BXE0690A	96T	2-8°C
TSHELISABXE0681B19272-8°CTSHELISABXE0681C48072-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0112A96T2-8°C	Troponin-I		ELISA	BXE0864A	96T	2-8°C
TSHELISABXE0681C480T2-8°CTSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)Liquid Stable (LS)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0112A96T2-8°C	TSH		ELISA	BXE0681A	96T	2-8°C
TSH RapidELISABXE0683A96T2-8°CUnconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°CVitamin-D Fast (90 Min)Liopholised (Lyo)ELISABXE0112A96T2-8°C	тѕн		ELISA	BXE0681B	192T	2-8°C
Unconjugated Estriol (UE3)ELISABXE0865A96T2-8°CVitamin B12ELISABXE0960A96T2-8°CVitamin-D (Blood Spots)ELISABXE0113A96T2-8°CVitamin-D Fast (90 Min)Liquid Stable (LS)ELISABXE0111A96T2-8°CVitamin-D Fast (90 Min)Liopholised (Lyo)ELISABXE0112A96T2-8°C	TSH		ELISA	BXE0681C	480T	2-8°C
Vitamin B12 ELISA BXE0960A 96T 2-8°C Vitamin-D (Blood Spots) ELISA BXE0113A 96T 2-8°C Vitamin-D (Blood Spots) ELISA BXE0113A 96T 2-8°C Vitamin-D Fast (90 Min) Liquid Stable (LS) ELISA BXE0111A 96T 2-8°C Vitamin-D Fast (90 Min) Lyopholised (Lyo) ELISA BXE0112A 96T 2-8°C	TSH Rapid		ELISA	BXE0683A	96T	2-8°C
Vitamin-D (Blood Spots) ELISA BXE0113A 96T 2-8°C Vitamin-D Fast (90 Min) Liquid Stable (LS) ELISA BXE0111A 96T 2-8°C Vitamin-D Fast (90 Min) Lyopholised (Lyo) ELISA BXE0112A 96T 2-8°C	Unconjugated Estriol (UE3)		ELISA	BXE0865A	96T	2-8°C
Vitamin-D Fast (90 Min) Liquid Stable (LS) ELISA BXE0111A 96T 2-8°C Vitamin-D Fast (90 Min) Lyopholised (Lyo) ELISA BXE0112A 96T 2-8°C	Vitamin B12		ELISA	BXE0960A	96T	2-8°C
Vitamin-D Fast (90 Min) Lyopholised (Lyo) ELISA BXE0112A 96T 2-8°C	Vitamin-D (Blood Spots)		ELISA	BXE0113A	96T	2-8°C
	Vitamin-D Fast (90 Min)	Liquid Stable (LS)	ELISA	BXE0111A	96T	2-8°C
	Vitamin-D Fast (90 Min)	Lyopholised (Lyo)	ELISA	BXE0112A	96T	2-8°C
	Zinc		ELISA	BXE0462A	96T	2-8°C





Immunoassays (CLIA)

Chemiluminescence Immunoassay (CLIA) is an advanced diagnostic technique used to detect and quantify specific molecules such as proteins, hormones, and antibodies in biological samples. It combines the sensitivity of immunoassays with chemiluminescence, where a chemical reaction produces light as a signal.

In CLIA, antibodies or antigens are labeled with a chemiluminescent substance, and when the target molecule binds to the labeled antibodies, a luminescent reaction is triggered. The emitted light is measured, and its intensity correlates with the concentration of the target molecule.

Chemiluminescence Immunoassay (CLIA) is a widely used detection method which combines highly sensitive chemiluminescence with highly specific immune response and affinity of antibodies. CLIA is widely used in clinical laboratories for various applications, including hormone assays, tumor marker testing, infectious disease diagnostics, and therapeutic drug monitoring. It offers high sensitivity, specificity, and a wide dynamic range, making it a valuable tool in medical diagnostics. Fortress Diagnostics manufactures a broad range of CLIA tests to support the growing demand for this greater quality method of testing.

Key Features & Benefits

Chemiluminescence Immunoassay (CLIA) offers several advantages that makes it the most advanced immunoassay technology on the market and invaluable in clinical diagnostics and research:



High Sensitivity: CLIA is the most sensitive method among immunoassay technologies.



High Linearity: This is crucial for ensuring the accuracy of results. The high linearity of CLIA assays makes them suitable for quantitative analysis.



No Protein Quenching Problem: Minimises the impact of interfering proteins while preserving the specificity and sensitivity of the assay.



Fast Incubation: Hence reduced assay time making CLIA more efficient and suitable for high-throughput testing environments, such as clinical laboratories.

Description	Details	Method	Cat No.	Size	Storage
Anti-HBc QN		CLIA	BXL0110A	100T/Kit	2-8°C
Alpha Feto Protein (AFP)		CLIA	BXL0129A	100T/Kit	2-8°C
Anti-HBc IgM		CLIA	BXL0113A	100T/Kit	2-8°C
Anti-HBe QN		CLIA	BXL0109A	100T/Kit	2-8°C
Anti-Mullerian Hormone (AMH)		CLIA	BXL0185A	100T/Kit	2-8°C
Beta2-MG		CLIA	BXL0144A	100T/Kit	2-8°C
BNP		CLIA	BXL0170A	100T/Kit	2-8°C
CIV		CLIA	BXL0183A	100T/Kit	2-8°C
CA125		CLIA	BXL0134A	100T/Kit	2-8°C
CA15-3		CLIA	BXL0135A	100T/Kit	2-8°C
CA19-9		CLIA	BXL0136A	100T/Kit	2-8°C
CA242		CLIA	BXL0139A	100T/Kit	2-8°C
CA50		CLIA	BXL0138A	100T/Kit	2-8°C
CA72-4		CLIA	BXL0137A	100T/Kit	2-8°C
Ceruloplasmin (CEA)		CLIA	BXL0130A	100T/Kit	2-8°C
СК-МВ		CLIA	BXL0159A	100T/Kit	2-8°C
CK-MB Stat		CLIA	BXL0167A	100T/Kit	2-8°C
Cortisol		CLIA	BXL0197A	100T/Kit	2-8°C
C-Peptide		CLIA	BXL0172A	100T/Kit	2-8°C

Immunoassays (CLIA) | SECTION 4

c-PSA CT cTnl cTnT	CLIA CLIA	BXL0133A	100T/Kit	
cTnl	CLIA		1001/140	2-8°C
	CLIA	BXL0148A	100T/Kit	2-8°C
cTnT	CLIA	BXL0161A	100T/Kit	2-8°C
	CLIA	BXL0160A	100T/Kit	2-8°C
CWS (Concentrated Wash Solution)	CLIA	BXL0200A	500ml/bottle	2-8°C
Cy21-1	CLIA	BXL0140A	100T/Kit	2-8°C
Disposable Tips (Bulk)	CLIA	BXL0204A	1000pcs/bag	2-8°C
Disposable Tips (Fully-Loaded)	CLIA	BXL0206A	96pcs*5levels*8sets/box	2-8°C
DU (Diluent Universal)	CLIA	BXL0201A	200ml/bottle	2-8°C
EBV EBNA1 IgA	CLIA	BXL0122A	100T/Kit	2-8°C
EBV VCA IgA	CLIA	BXL0123A	100T/Kit	2-8°C
EBV Zta IgA	CLIA	BXL0124A	100T/Kit	2-8°C
Estradiol (E2)	CLIA	BXL0191A	100T/Kit	2-8°C
F-Beta-HCG	CLIA	BXL0187A	100T/Kit	2-8°C
Ferritin	CLIA	BXL0174A	100T/Kit	2-8°C
Follicle Stimulating Hormone (FSH)	CLIA	BXL0190A	100T/Kit	2-8°C
Free PSA (f-PSA)	CLIA	BXL0131A	100T/Kit	2-8°C
Free-T3 (FT3)	CLIA	BXL0153A	100T/Kit	2-8°C
Free-T4 (FT4)	CLIA	BXL0154A	100T/Kit	2-8°C
G17	CLIA	BXL0149A	100T/Kit	2-8°C
НА	CLIA	BXL0181A	100T/Kit	2-8°C
HAV Ab	CLIA	BXL0101A	100T/Kit	2-8°C
HAV IgM	CLIA	BXL0102A	100T/Kit	2-8°C
HBcAb	CLIA	BXL0107A	100T/Kit	2-8°C
нвсАЬ нQ	CLIA	BXL0112A	100T/Kit	2-8°C
HBeAb	CLIA	BXL0106A	100T/Kit	2-8°C
HBeAg	CLIA	BXL0105A	100T/Kit	2-8°C
HBeAg QN	CLIA	BXL0108A	100T/Kit	2-8°C
HBsAb	CLIA	BXL0104A	100T/Kit	2-8°C
HBsAg	CLIA	BXL0103A	100T/Kit	2-8°C
HBsAg HQ	CLIA	BXL0111A	100T/Kit	2-8°C
HBV PreS1	CLIA	BXL0114A	100T/Kit	2-8°C
HCG Stat (HCG)	CLIA	BXL0194A	100T/Kit	2-8°C
HCV Ab	CLIA	BXL0115A	100T/Kit	2-8°C
HDV IgG	CLIA	BXL0119A	100T/Kit	2-8°C
HDV IgM	CLIA	BXL0118A	100T/Kit	2-8°C
HE4	CLIA	BXL0143A	100T/Kit	2-8°C
HEV IgG	CLIA	BXL0121A	100T/Kit	2-8°C
HEV IgM	CLIA	BXL0120A	100T/Kit	2-8°C
H-FABP	CLIA	BXL0165A	100T/Kit	2-8°C
ні Ад/АЬ	CLIA	BXL0116A	100T/Kit	2-8°C
Hs-CRP	CLIA	BXL0163A	100T/Kit	2-8°C
Hs-cTnT STAT	CLIA	BXL0171A	100T/Kit	2-8°C
Human Growth Hormone (hGH)	CLIA	BXL0196A	100T/Kit	2-8°C
IL-6	CLIA	BXL0176A	100T/Kit	2-8°C
IL-6 Stat	CLIA	BXL0179A	100T/Kit	2-8°C
	CLIA	BXL0173A	100T/Kit	2-8°C
Insulin			100T/Kit	2-8°C

SECTION 4 | Immunoassays (CLIA)

LNCLIABXL01PHA1007/Nit2.4°CLp-PLA2CLIABXL016A1007/Nit2.4°CMyoCLIABXL016A1007/Nit2.4°CMyo StatCLIABXL016A1007/Nit2.4°CNGALCLIABXL0180A1007/Nit2.4°CNT-proBPCLIABXL0140A1007/Nit2.4°CNT-proBPCLIABXL0141A1007/Nit2.4°CNT-proBPSCLIABXL0140A1007/Nit2.4°CPCTCLIABXL0140A1007/Nit2.4°CPCTCLIABXL0147A1007/Nit2.4°CPCTCLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCICLIABXL0147A1007/Nit2.4°CPCGRPCLIABXL0147A1007/Nit2.4°CProBationCLIABXL0147A1007/Nit2.4°CPCGRPCLIABXL0147A1007/Nit2.4°CPCGRPCLIABXL0147A1007/Nit2.4°CPCGRPCLIABXL0147A1007/Nit2.4°CPCGRPCLIABXL0147A1007/Nit2.4°CPCGRPCLIA<	Description	Details	Method	Cat No.	Size	Storage
Lp-P.A2CILIABXL0166A100T/KI:2.8°CMyoCILIABXL0162A100T/KI:2.4°CMyo StatCILIABXL0163A100T/KI:2.4°CNSALCILIABXL0163A100T/KI:2.4°CNT-proBNPCILIABXL0164A100T/KI:2.4°CNT-proBNPCILIABXL0164A100T/KI:2.4°CPCTCILIABXL0164A100T/KI:2.4°CPCTCILIABXL0164A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCTCILIABXL0174A100T/KI:2.4°CPCOSCIOCILIABXL0174A100T/KI:2.4°CPCIACTIOCILIABXL0174A100T/KI:2.4°CPCIACTIOCILIABXL0174A100T/KI:2.4°CPCIACTIOCILIABXL0174A100T/KI:2.4°CPCIACTIOCILIABXL0174A100T/KI:2.4°CPCIACTIOCILIABXL0174A100T/KI:2.4°C	Leutinising Hormone (LH)		CLIA	BXL0186A	100T/Kit	2-8°C
MyoCLIABXL0162A1007/Kit2.4°CMyo StabCLIABXL0162A1007/Kit2.4°CNGAL.CLIABXL0180A1007/Kit2.4°CNSECLIABXL0180A1007/Kit2.4°CNT-proBNPCLIABXL0164A1007/Kit2.4°CNT-proBNPS StatCLIABXL0164A1007/Kit2.4°CPCTStat1007/Kit2.4°CPCTCLIABXL0176A1007/Kit2.4°CPCTStat1007/Kit2.4°CPCTCLIABXL0176A1007/Kit2.4°CPCTStat1007/Kit2.4°CPCTCLIABXL0176A1007/Kit2.4°CPCIPCIAStat1007/Kit2.4°CPCICLIABXL0176A1007/Kit2.4°CPCIPCIAStat1007/Kit2.4°CPCIPCIABXL0176A1007/Kit2.4°CPCIACLIABXL0174A1007/Kit2.4°CProJectin (PRL)CLIABXL0182A1007/Kit2.4°CProJectin (PRL)CLIABXL017A1007/Kit2.4°CPS (PC-Tigger Solution)CLIABXL017A1007/Kit2.4°CSACLIABXL017A1007/Kit2.4°CSAS-COV-2 AbCLIABXL017A1007/Kit2.4°CSAS-COV-2 IgMCLIABXL017A1007/Kit2.4°CSAS-COV-2 IgMCLIABXL017A1007/Kit2.4°CSAS-COV-2 IgMCLIA <td>LN</td> <td></td> <td>CLIA</td> <td>BXL0184A</td> <td>100T/Kit</td> <td>2-8°C</td>	LN		CLIA	BXL0184A	100T/Kit	2-8°C
Myo Stat:CIAIBXL0169A1007/Kit2-8°CNGALCIAIBXL0160A1007/Kit2-8°CNSECIAIBXL0164A1007/Kit2-8°CNT-proBNPSatCIAIBXL016A1007/Kit2-8°CPCTCIAIBXL017A1007/Kit2-8°CPCTCIAIBXL017A1007/Kit2-8°CPCTCIAIBXL017A1007/Kit2-8°CPCTCIAIBXL017A1007/Kit2-8°CPCTCIAIBXL017A1007/Kit2-8°CPGICIAIBXL017A1007/Kit2-8°CPGICIAIBXL017A1007/Kit2-8°CPGICIAIBXL017A1007/Kit2-8°CPGICIAIBXL017A1007/Kit2-8°CProgesteroneCIAIBXL017A1007/Kit2-8°CProGRPCIAIBXL017A1007/Kit2-8°CProSteringer Solution/CIAIBXL017A1007/Kit2-8°CSAS-Cov2 AbCIAIBXL017A1007/Kit2-8°CSARS-Cov2 AbCIAIBXL017A1007/Kit2-8°CSARS-Cov2 AbCIAIBXL017A1007/Kit2-8°CSARS-Cov2 AbCIAIBXL017A1007/Kit2-8°CSARS-Cov2 AbCIAIBXL017A1007/Kit2-8°CSARS-Cov2 AbCIAIBXL017A1007/Kit2-8°CTebta-HCGCIABXL017A1007/Kit2-8°CTebta-HCGCIABXL017A1007/Kit	Lp-PLA2		CLIA	BXL0166A	100T/Kit	2-8°C
NGALCLIABXL0180A1007/Xit2.4°CNSECLIABXL0141A1007/Xit2.4°CNT-proBNPCLIABXL016A1007/Xit2.4°CNT-proBNPStatCLIABXL015A1007/Xit2.4°CPCTCLIABXL015A1007/Xit2.4°CPCTCLIABXL015A1007/Xit2.4°CPCTCLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CPCICLIABXL015A1007/Xit2.4°CProgstatCLIABXL015A1007/Xit2.4°CProgstatCLIABXL015A1007/Xit2.4°CProdschi (PRL)CLIABXL015A1007/Xit2.4°CProdschi (PRL)CLIABXL012A1007/Xit2.4°CSAS-CoV-2 IgGCLIABXL012A1007/Xit2.4°CSARS-CoV-2 IgGCLIABXL012A1007/Xit2.4°CSAS-CoV-2 IgGCLIABXL012A1007/Xit2.4°CTestsstoreneCLIABXL012A1007/Xit2.4°CTo-AbCLIABXL012A1007/Xit2.4°CTo-AbCLIABXL012A1007/Xit2.4°CTo-AbCLIABXL012A1007/Xit2.4°C </td <td>Муо</td> <td></td> <td>CLIA</td> <td>BXL0162A</td> <td>100T/Kit</td> <td>2-8°C</td>	Муо		CLIA	BXL0162A	100T/Kit	2-8°C
NSECIANSKL014AIOT/KR2.4°CNT-proBNPCIANSKL016AI1007/KR2.4°CNT-proBNP StatCIANSKL016AI1007/KR2.4°CPCTStatSKL017AI1007/KR2.4°CPCTStatSKL017AI1007/KR2.4°CPCTStatSKL017AI1007/KR2.4°CPCTStatSKL017AI1007/KR2.4°CPGICIANSKL017AI1007/KR2.4°CPGICIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CProg StatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStatCIANSKL017AI1007/KR2.4°CStat <t< td=""><td>Myo Stat</td><td></td><td>CLIA</td><td>BXL0169A</td><td>100T/Kit</td><td>2-8°C</td></t<>	Myo Stat		CLIA	BXL0169A	100T/Kit	2-8°C
NT-proBNPCLIABXL0164A100T/Kik2-8°CNT-proBNP StatCLIABXL0168A100T/Kik2-8°CPCTCLIABXL0175A100T/Kik2-8°CPCT StatDCIABXL0176A100T/Kik2-8°CPCIDCIABXL0146A100T/Kik2-8°CPCIDCIABXL0146A100T/Kik2-8°CPCIDCIABXL0146A100T/Kik2-8°CPCIDCIABXL0146A100T/Kik2-8°CPCIDCIABXL0146A100T/Kik2-8°CProg StatDCIABXL018A100T/Kik2-8°CProGsPCLIABXL018A100T/Kik2-8°CProGsteroneCLIABXL018A100T/Kik2-8°CProGsteroneCLIABXL018A100T/Kik2-8°CProGsteroneCLIABXL018A100T/Kik2-8°CProGsteroneCLIABXL018A100T/Kik2-8°CProGsteroneCLIABXL0207A96c+S1evels*4sets/bx2-8°CSADCIABXL017A100T/Kik2-8°CSADCIABXL017A100T/Kik2-8°CSAS-CoV-2 Ig/CLIABXL018A100T/Kik2-8°CSAS-CoV-2 Ig/CLIABXL018A100T/Kik2-8°CTG-DatineCLIABXL017A100T/Kik2-8°CTG-DatineCLIABXL018A100T/Kik2-8°CTG-DatineCLIABXL018A100T/Kik2-8°CTG-DatineCLIABXL017A	NGAL		CLIA	BXL0180A	100T/Kit	2-8°C
NrproBNP StatCLIABXL0168A1007/Kit2.8°CPCTCLIABXL0175A1007/Kit2.8°CPCT StatCLIABXL0175A1007/Kit2.8°CPGICLIABXL0178A1007/Kit2.8°CPGICLIABXL0182A1007/Kit2.8°CPGICLIABXL0182A1007/Kit2.8°CPGICLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL0182A1007/Kit2.8°CProg StatCLIABXL017A1007/Kit2.8°CProg StatCLIABXL027A1007/Kit2.8°CProg StatCLIABXL017A1007/Kit2.8°CStaction Cuvettes (Gulk)CLIABXL017A1007/Kit2.8°CSAR-CoV-2 JgCLIABXL017A1007/Kit2.8°CSARS-CoV-2 JgCLIABXL017A1007/Kit2.8°CTebea-HCGCLIABXL017A1007/Kit2.8°CTebea-HCGCLIABXL0152A1007/Kit2.8°CTotal TACLIABXL0152A1007/Kit2.8°CTotal TACL	NSE		CLIA	BXL0141A	100T/Kit	2-8°C
PcTCLIABXL017SA100T/Kit2.8°CPCT StatCLIABXL017SA100T/Kit2.8°CPCIBXL017SA100T/Kit2.8°CPCICLIABXL014SA100T/Kit2.8°CPCICLIABXL018ZA100T/Kit2.8°CPCICLIABXL018ZA100T/Kit2.8°CProg StatCLIABXL019SA100T/Kit2.8°CProgSteroneCLIABXL019SA100T/Kit2.8°CProgSteroneCLIABXL019SA100T/Kit2.8°CProdRPCLIABXL018SA100T/Kit2.8°CProdRPCLIABXL018SA100T/Kit2.8°CProdRPCLIABXL020SA100T/Kit2.8°CProdRPCLIABXL020SA100T/Kit2.8°CProdRPCLIABXL020SA100T/Kit2.8°CSatation Cuvettes (Bulk)CLIABXL012A100T/Kit2.8°CSAR-Cov-2 AbCLIABXL017A100T/Kit2.8°CSAR-Cov-2 IgGCLIABXL012A100T/Kit2.8°CSCCLIABXL012A100T/Kit2.8°CTestasteroneCLIABXL013ZA100T/Kit2.8°CTotal PSA (L-SA)CLIABXL012A100T/Kit2.8°CSCCLIABXL015ZA100T/Kit2.8°CTotal PSA (L-SA)CLIABXL015ZA100T/Kit2.8°CTotal PSA (L-SA)CLIABXL015ZA100T/Kit2.8°CTotal PSA (L-SA)CLIA	NT-proBNP		CLIA	BXL0164A	100T/Kit	2-8°C
PC StatCLIABXL0178A1007/kit2.8°CPGICLIABXL014SA1007/kit2.8°CPGICLIABXL014SA1007/kit2.8°CPGINCLIABXL014SA1007/kit2.8°CPINPSCLIABXL018ZA1007/kit2.8°CProgesteroneCLIABXL019XA1007/kit2.8°CProgesteroneCLIABXL019XA1007/kit2.8°CProGRPCLIABXL019XA1007/kit2.8°CProGRPCLIABXL019XA1007/kit2.8°CProGRPCLIABXL019XA1007/kit2.8°CProGRPCLIABXL019XA1007/kit2.8°CProGRPCLIABXL017A1007/kit2.8°CProGRPCLIABXL017A1007/kit2.8°CSAACUIABXL012XA1007/kit2.8°CSAACLIABXL017A1007/kit2.8°CSARS-Cov-2 JgCLIABXL017A1007/kit2.8°CSARS-Cov-2 IgMCLIABXL012XA1007/kit2.8°CSCCCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°CTo-AbCLIABXL015XA1007/kit2.8°C <td>NT-proBNP Stat</td> <td></td> <td>CLIA</td> <td>BXL0168A</td> <td>100T/Kit</td> <td>2-8°C</td>	NT-proBNP Stat		CLIA	BXL0168A	100T/Kit	2-8°C
PGICLIABXL014SA100T/Kit2-8°CPGIICLIABXL014GA100T/Kit2-8°CPIINPCLIABXL018ZA100T/Kit2-8°CProgStatCLIABXL019SA100T/Kit2-8°CProgsteroneCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CProGRPCLIABXL019SA100T/Kit2-8°CSAGCLIABXL019SA100T/Kit2-8°CSAACLIABXL017A100T/Kit2-8°CSARS-CoV-2 AbCLIABXL017A100T/Kit2-8°CSARS-CoV-2 IgGCLIABXL017A100T/Kit2-8°CSARS-CoV-2 IgACLIABXL017A100T/Kit2-8°CTebehGCCLIABXL015A100T/Kit2-8°CToT-AbCLIABXL015A100T/Kit2-8°CTotal-TaffYCLIABXL015A100T/Kit2-8°CTotal-TaffYCLIABXL015A100T/Kit2-8°CTotal-TaffYCLIABXL015A100T/Kit2-8°CTotal-TaffYCLIABXL015A100T/Kit2-8°CTotal-TaffYCLIABXL015A<	РСТ		CLIA	BXL0175A	100T/Kit	2-8°C
PGIICI.IABXL0146A1007/Kit2.8°CPIINPCI.IABXL015A1007/Kit2.8°CProg StatCI.IABXL015A1007/Kit2.8°CProgesteroneCI.IABXL013A1007/Kit2.8°CProGRPCI.IABXL013A1007/Kit2.8°CProlactin (PRL)CI.IABXL0202A200m/bottle2.8°CReaction Covettes (Bulk)CI.IABXL027A1007/Kit2.8°CSAACI.IABXL017A1007/Kit2.8°CSAACI.IABXL017A1007/Kit2.8°CSARS-Cov-2 AbCI.IABXL017A1007/Kit2.8°CSARS-Cov-2 IgGCI.IABXL012A1007/Kit2.8°CSARS-Cov-2 IgGCI.IABXL012A1007/Kit2.8°CSCCCI.IABXL012A1007/Kit2.8°CTestesteroneCI.IABXL012A1007/Kit2.8°CTotal TaffCI.IABXL012A1007/Kit2.8°CSCCCI.IABXL012A1007/Kit2.8°CTotal TaffCI.IABXL012A1007/Kit2.8°CTotal TaffCI.IABXL013A1007/Kit2.8°CTotal TaffCI.IABXL013A1007/Kit2.8°CTotal TaffCI.IABXL013A1007/Kit2.8°CTotal TaffCI.IABXL013CA1007/Kit2.8°CTotal TaffCI.IABXL013CA1007/Kit2.8°CTotal TaffCI.IABXL013CA1007/Kit2.8°C <td>PCT Stat</td> <td></td> <td>CLIA</td> <td>BXL0178A</td> <td>100T/Kit</td> <td>2-8°C</td>	PCT Stat		CLIA	BXL0178A	100T/Kit	2-8°C
PHINPCILIABXL0152A100T/Kit2.8°CProg StatCILABXL0193A100T/Kit2.8°CProgesteroneCILABXL0193A100T/Kit2.8°CProGRPCILABXL0147A100T/Kit2.8°CProlactin (PRL)CILABXL0120A100T/Kit2.8°CProfere-Trigger Solution)CILABXL020A200ml/bottle2.8°CReaction Cuvettes (Bulk)CILABXL020A200ml/bottle2.8°CSAACILABXL027A100T/Kit2.8°CSAACILABXL012AA100T/Kit2.8°CSARS-Cov-2 IgGCILABXL012AA100T/Kit2.8°CSARS-Cov-2 IgGCILABXL012AA100T/Kit2.8°CSARS-Cov-2 IgGCILABXL012AA100T/Kit2.8°CSTCCILABXL012AA100T/Kit2.8°CTebeta-HCGCILABXL012AA100T/Kit2.8°CTGCILABXL012AA100T/Kit2.8°CTGCILABXL012AA100T/Kit2.8°CTotal TSCILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°CTotal TACILABXL015AA100T/Kit2.8°C <td>PGI</td> <td></td> <td>CLIA</td> <td>BXL0145A</td> <td>100T/Kit</td> <td>2-8°C</td>	PGI		CLIA	BXL0145A	100T/Kit	2-8°C
Prog StatCLIABXL0195A100T/Kit2.8°CProgesteroneCLIABXL0193A100T/Kit2.8°CProGRPCLIABXL0147A100T/Kit2.8°CProlactin (PRL)CLIABXL022A200ml/bottle2.6°CProfer-Trigger Solution)CLIABXL022A1000r/kit2.6°CReaction Cuvettes (Bulk)CLIABXL022A1000r/kit2.6°CSAACLIABXL027A06pcs/bag2.6°CSARS-Cov-2 AbgCLIABXL017A100T/Kit2.6°CSARS-Cov-2 IgGCLIABXL017A100T/Kit2.6°CSARS-Cov-2 IgGCLIABXL012A100T/Kit2.6°CSARS-Cov-2 IgGCLIABXL012A100T/Kit2.6°CSARS-Cov-2 IgGCLIABXL012A100T/Kit2.6°CSARS-Cov-2 IgGCLIABXL012A100T/Kit2.6°CSCCLIABXL012A100T/Kit2.6°CTogCLIABXL012A100T/Kit2.6°CTGCLIABXL012A100T/Kit2.6°CTGCLIABXL015A100T/Kit2.6°CTogCLIABXL015A100T/Kit2.6°CTogCLIABXL015A100T/Kit2.6°CTGCLIABXL015A100T/Kit2.6°CTogCLIABXL015A100T/Kit2.6°CTogCLIABXL015A100T/Kit2.6°CTogCLIABXL015A100T/Kit2.6°CTogCLIA <t< td=""><td>PGII</td><td></td><td>CLIA</td><td>BXL0146A</td><td>100T/Kit</td><td>2-8°C</td></t<>	PGII		CLIA	BXL0146A	100T/Kit	2-8°C
ProgesteroneCLIABXL0193A100T/Kit2.8°CProGRPCLIABXL0147A100T/Kit2.8°CProlactin (PRL)CLIABXL0202A200ml/bottle2.8°CReaction Cuvettes (Bulk)CLIABXL0205A1000pcs/bag2.8°CReaction Cuvettes (Fully-loaded)CLIABXL0207A96pcs*5levels*4sets/box2.8°CSAACLIABXL0177A100T/Kit2.8°CSARS-CoV-2 AbCLIABXL0127A100T/Kit2.8°CSARS-CoV-2 IgGCLIABXL0127A100T/Kit2.8°CSCCCLIABXL0128A100T/Kit2.8°CTestosteroneCLIABXL0128A100T/Kit2.8°CTGCLIABXL0128A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0192A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL015A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL015A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL015A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL015A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL015A<	PIIINP		CLIA	BXL0182A	100T/Kit	2-8°C
ProGRPCI.IABXL0147A100T/Kit2.8°CProlactin (PR.)CI.IABXL018AA100T/Kit2.8°CPTS (Pre-Trigger Solution)CI.IABXL0202A200m//bottle2.8°CReaction Cuvettes (Bulk)CI.IABXL0207A96pcs*Slevels*4sets/box2.8°CSAACI.IABXL0177A100T/Kit2.8°CSARS-Cov-2 AbCI.IABXL017A100T/Kit2.8°CSARS-Cov-2 IgGCI.IABXL0126A100T/Kit2.8°CSCCCI.IABXL0127A100T/Kit2.8°CTebet-HCGCI.IABXL0128A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTogCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL0157A100T/Kit2.8°CTog AbCI.IABXL015A100T/Kit2.8°C <t< td=""><td>Prog Stat</td><td></td><td>CLIA</td><td>BXL0195A</td><td>100T/Kit</td><td>2-8°C</td></t<>	Prog Stat		CLIA	BXL0195A	100T/Kit	2-8°C
Prolactin (PRL)CLIABXL0188A100T/Kit2.8°CPTS (Pre-Trigger Solution)CLIABXL0202A200m/bottle2.8°CReaction Cuvettes (Bulk)CLIABXL0207A96pcs*Slevels*4sets/bx02.8°CReaction Cuvettes (Fully-loaded)CLIABXL0177A100T/Kit2.8°CSAACLIABXL0126A100T/Kit2.8°CSARS-CoV-2 AbCLIABXL0126A100T/Kit2.8°CSARS-CoV-2 IgGCLIABXL0127A100T/Kit2.8°CSARS-CoV-2 IgMCLIABXL0128A100T/Kit2.8°CSCCCLIABXL0189A100T/Kit2.8°CTebta-HCGCLIABXL0192A100T/Kit2.8°CTGCLIABXL0157A100T/Kit2.8°CTGCLIABXL0157A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0154A100T/Kit2.8°CTotal-T3 (TT3)CLIABXL0154A100T/Kit2.8°CTPAbCLIABXL0154A100T/Kit2.8°CTPA-AbCLIABXL015A100T/Kit2.8°CTFAACLIABXL015A100T/Kit2.8°CTFAACLIABXL015A100T/Kit2.8°CTotal-T4 (TT4)CLIABXL015A100T/Kit2.8°CTFAACLIABXL015A100T/Kit2.8°CTFAACLIABXL015A100T/Kit2.8°CTFAACLIABXL015A </td <td>Progesterone</td> <td></td> <td>CLIA</td> <td>BXL0193A</td> <td>100T/Kit</td> <td>2-8°C</td>	Progesterone		CLIA	BXL0193A	100T/Kit	2-8°C
PTS (Pre-Trigger Solution)CIABXL0202A200ml/bottle2.8°CReaction Cuvettes (Bulk)CIABXL0203A1000pcs/bag2.8°CReaction Cuvettes (Fully-loaded)CIABXL027A96pcs*Slevels*4sets/bx02.8°CSAACIABXL0177A100T/Kit2.8°CSARS-Cov'2 AbCIABXL0127A100T/Kit2.8°CSARS-Cov'2 IgGCIABXL012A100T/Kit2.8°CSARS-Cov'2 IgMCIABXL012A100T/Kit2.8°CTebea-HCGCIABXL012A100T/Kit2.8°CTo Tabeta-HCGCIABXL012A100T/Kit2.8°CTGCIABXL012A100T/Kit2.8°CTG-AbCIABXL012A100T/Kit2.8°CTotal PSA (t-PSA)CIABXL012A100T/Kit2.8°CTotal T3 (TT3)CIABXL013A100T/Kit2.8°CTotal-T4 (TT4)CIABXL013A100T/Kit2.8°CTP-AbCIABXL013A100T/Kit2.8°CTP-AbCIABXL013A100T/Kit2.8°CTFA-BCIABXL013A100T/Kit2.8°CTFA-BCIABXL013A100T/Kit2.8°CTFA-BCIABXL013A100T/Kit2.8°CTotal PSA (t-PSA)CIABXL013A100T/Kit2.8°CTotal PSA (t-PSA)CIABXL013A100T/Kit2.8°CTotal T4 (TT4)CIABXL013A100T/Kit2.8°CTFA-BCIABXL013A <td>ProGRP</td> <td></td> <td>CLIA</td> <td>BXL0147A</td> <td>100T/Kit</td> <td>2-8°C</td>	ProGRP		CLIA	BXL0147A	100T/Kit	2-8°C
Reaction Cuvettes (Bulk)CLIABXL0205A1000pcs/bag2-8°CReaction Cuvettes (Fully-loaded)CLIABXL0207A96pcs*Slevels*4sets/box2-8°CSAACLIABXL0177A1007/Kit2-8°CSARS-CoV-2 AbCLIABXL0126A1007/Kit2-8°CSARS-CoV-2 IgGCLIABXL0127A1007/Kit2-8°CSARS-CoV-2 IgGCLIABXL0128A1007/Kit2-8°CSCCCLIABXL0128A1007/Kit2-8°CSCCCLIABXL0128A1007/Kit2-8°CT-Beta-HCGCLIABXL0192A1007/Kit2-8°CTestosteroneCLIABXL0157A1007/Kit2-8°CTGCLIABXL015A1007/Kit2-8°CTotal PSA (t-PSA)CLIABXL015A1007/Kit2-8°CTotal-T3 (TT3)CLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAbCLIABXL015A1007/Kit2-8°CTPAC <td>Prolactin (PRL)</td> <td></td> <td>CLIA</td> <td>BXL0188A</td> <td>100T/Kit</td> <td>2-8°C</td>	Prolactin (PRL)		CLIA	BXL0188A	100T/Kit	2-8°C
Reaction Cuvettes (Fully-loaded)CLIABXL0207A96pcs*5levels*4sets/box2.8°CSAACLIABXL0177A100T/Kit2.8°CSARS-CoV-2 IgGCLIABXL0126A100T/Kit2.8°CSARS-CoV-2 IgGCLIABXL0127A100T/Kit2.8°CSARS-CoV-2 IgGCLIABXL0128A100T/Kit2.8°CSCCCLIABXL0128A100T/Kit2.8°CSCCCLIABXL0192A100T/Kit2.8°CT-Beta-HCGCLIABXL0192A100T/Kit2.8°CTGCLIABXL0192A100T/Kit2.8°CTG-AbCLIABXL0157A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal-T3 (TT3)CLIABXL0152A100T/Kit2.8°CTotal-T4 (TT4)CLIABXL0154100T/Kit2.8°CTo-AbCLIABXL0154100T/Kit2.8°CTotal-T5 (TT3)CLIABXL0154100T/Kit2.8°CTotal-T4 (TT4)CLIABXL0154100T/Kit2.8°CTo-AbCLIABXL0154100T/Kit2.8°CTAbCLIABXL0154100T/Kit2.8°CTG-AbCLIABXL0154100T/Kit2.8°CTotal-T4 (TT4)CLIABXL0154100T/Kit2.8°CTotal-T5CLIABXL0154100T/Kit2.8°CTAbCLIABXL0154100T/Kit2.8°CTG-AbCLIABXL0154100T/Kit2.8°C<	PTS (Pre-Trigger Solution)		CLIA	BXL0202A	200ml/bottle	2-8°C
SAA CLIA BXL0177A 100T/Kit 2-8°C SARS-CoV-2 Ab CLIA BXL0126A 100T/Kit 2-8°C SARS-CoV-2 IgG CLIA BXL0127A 100T/Kit 2-8°C SARS-CoV-2 IgG CLIA BXL0127A 100T/Kit 2-8°C SARS-CoV-2 IgG CLIA BXL0128A 100T/Kit 2-8°C SARS-CoV-2 IgM CLIA BXL0128A 100T/Kit 2-8°C SCC CLIA BXL0128A 100T/Kit 2-8°C TBeta-HCG CLIA BXL0128A 100T/Kit 2-8°C Testosterone CLIA BXL017A 100T/Kit 2-8°C TG CLIA BXL0157A 100T/Kit 2-8°C Total PSA (t-PSA) CLIA BXL0152A 100T/Kit 2-8°C Total PSA (t-PSA) CLIA BXL0151A 100T/Kit 2-8°C Total-T3 (TT3) CLIA BXL0151A 100T/Kit 2-8°C TP Ab CLIA BXL0151A 100T/Kit 2-8°C TPA-Ab CLIA BXL0156A 100T/Kit 2-8°C <td< td=""><td>Reaction Cuvettes (Bulk)</td><td></td><td>CLIA</td><td>BXL0205A</td><td>1000pcs/bag</td><td>2-8°C</td></td<>	Reaction Cuvettes (Bulk)		CLIA	BXL0205A	1000pcs/bag	2-8°C
SARS-CoV-2 AbCLIABXL0126A100T/Kit2-8°CSARS-CoV-2 IgGCLIABXL0127A100T/Kit2-8°CSARS-CoV-2 IgMCLIABXL0128A100T/Kit2-8°CSCCCLIABXL012A100T/Kit2-8°CT-Beta-HCGCLIABXL0192A100T/Kit2-8°CTestosteroneCLIABXL015A100T/Kit2-8°CTGCLIABXL015A100T/Kit2-8°CTG-AbCLIABXL015A100T/Kit2-8°CTotal FSA (t-PSA)CLIABXL015A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL015A100T/Kit2-8°CTP AbCLIABXL015A100T/Kit2-8°CTPAbCLIABXL015A100T/Kit2-8°CTP AbCLIABXL015A100T/Kit2-8°CTP AbCLIABXL015A100T/Kit2-8°CTRAbCLIABXL015A100T/Kit2-8°CTG ABCLIABXL015A100T/Kit2-8°CTRAbCLIABXL015A100T/Kit2-8°CTS (Trigger Solution)CLIABXL015A100T/Kit2-8°CTuberculosis (TB)CLIABXL015A200m/bottle2-8°CCLIABXL015A100T/Kit2-8°C2-8°CTB AbCLIABXL015A100T/Kit2-8°CTB AbCLIABXL015A100T/Kit2-8°CTB AbCLIABXL015A100T/Kit2-8°CTB AbCLIABXL015A<	Reaction Cuvettes (Fully-loaded)		CLIA	BXL0207A	96pcs*5levels*4sets/box	2-8°C
SARS-CoV-2 IgGCLIABXL0127A100T/Kit2.8°CSARS-CoV-2 IgMCLIABXL0128A100T/Kit2.8°CSCCCLIABXL0142A100T/Kit2.8°CT-Beta-HCGCLIABXL0189A100T/Kit2.8°CTestosteroneCLIABXL0192A100T/Kit2.8°CTGCLIABXL0157A100T/Kit2.8°CTG-AbCLIABXL0157A100T/Kit2.8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2.8°CTotal-T3 (TT3)CLIABXL015A100T/Kit2.8°CTotal-T4 (T4)CLIABXL015A100T/Kit2.8°CTP-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTRAbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/Kit2.8°CTFO-AbCLIABXL015A100T/	SAA		CLIA	BXL0177A	100T/Kit	2-8°C
SARS-Cov-2 IgM CLIA BXL0128A 1007/Kit 2.8°C SCC CLIA BXL0142A 1007/Kit 2.8°C T-Beta-HCG CLIA BXL0142A 1007/Kit 2.8°C Testosterone CLIA BXL0192A 1007/Kit 2.8°C TG CLIA BXL0152A 1007/Kit 2.8°C TG-Ab CLIA BXL0152A 1007/Kit 2.8°C Total PSA (t-PSA) CLIA BXL0152A 1007/Kit 2.8°C Total-T3 (TT3) CLIA BXL0152A 1007/Kit 2.8°C Total-T4 (TT4) CLIA BXL0151A 1007/Kit 2.8°C TPO-Ab CLIA BXL0152A 1007/Kit 2.8°C TRAb CLIA BXL0152A 1007/Kit 2.8°C Total-T4 (TT4) CLIA BXL0152A 1007/Kit 2.8°C TRAb CLIA BXL0154A 1007/Kit 2.8°C TRAb CLIA BXL015A 1007/Kit 2.8°C TRAb CLIA BXL015A 1007/Kit 2.8°C TS (Trigger Solution)	SARS-CoV-2 Ab		CLIA	BXL0126A	100T/Kit	2-8°C
SCC CLIA BXL0142A 100T/Kit 2-8°C T-Beta-HCG CLIA BXL0189A 100T/Kit 2-8°C Testosterone CLIA BXL0192A 100T/Kit 2-8°C TG CLIA BXL0157A 100T/Kit 2-8°C TG-Ab CLIA BXL0152A 100T/Kit 2-8°C Thyroid Stimulating Hormone (TSH) CLIA BXL0152A 100T/Kit 2-8°C Total PSA (t-PSA) CLIA BXL0152A 100T/Kit 2-8°C Total-T3 (TT3) CLIA BXL0151A 100T/Kit 2-8°C TP-Ab CLIA BXL0151A 100T/Kit 2-8°C TP-Ab CLIA BXL0151A 100T/Kit 2-8°C TPA-Ab CLIA BXL0151A 100T/Kit 2-8°C TPA-Ab CLIA BXL0151A 100T/Kit 2-8°C TPA-Ab CLIA BXL015A 100T/Kit 2-8°C TRAb CLIA BXL015A 100T/Kit 2-8°C TS (Trigger Solution) CLIA BXL015A 100T/Kit 2-8°C TS (Trigg	SARS-CoV-2 lgG		CLIA	BXL0127A	100T/Kit	2-8°C
T-Beta-HCGCLIABXL0189A100T/Kit2-8°CTestosteroneCLIABXL0192A100T/Kit2-8°CTGCLIABXL0157A100T/Kit2-8°CTG-AbCLIABXL015SA100T/Kit2-8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP-AbCLIABXL0151A100T/Kit2-8°CTRAbCLIABXL0151A100T/Kit2-8°CTFO-AbCLIABXL0151A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTAbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTAbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0156A100T/Kit2-8°CTFO-AbCLIABXL0158A100T/Kit2-8°CTFO-AbCLIABXL0158A200m/bottle2-8°CTFO-AbCLIABXL0203A200m/bottle2-8°CTFO-AbCLIABXL0125A200m/bottle2-8°CTFO-AbCLIABXL0203A200m/bottle2-8°CTFO-AbCLIABXL0203A <td>SARS-CoV-2 IgM</td> <td></td> <td>CLIA</td> <td>BXL0128A</td> <td>100T/Kit</td> <td>2-8°C</td>	SARS-CoV-2 IgM		CLIA	BXL0128A	100T/Kit	2-8°C
TestosteroneCLIABXL0192A100T/Kit2-8°CTGCLIABXL0157A100T/Kit2-8°CTG-AbCLIABXL0155A100T/Kit2-8°CThyroid Stimulating Hormone (TSH)CLIABXL0152A100T/Kit2-8°CTotal PSA (t-PSA)CLIABXL0152A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP AbCLIABXL0117A100T/Kit2-8°CTFO-AbCLIABXL0158A100T/Kit2-8°CTRAbCLIABXL0158A100T/Kit2-8°CTS (Trigger Solution)CLIABXL0203A200m/bottle2-8°CTuberculosis (TB)CLIABXL0125A200m/bottle2-8°C	scc		CLIA	BXL0142A	100T/Kit	2-8°C
TGCLIABXL0157A100T/Kit2-8°CTG-AbCLIABXL0155A100T/Kit2-8°CThyroid Stimulating Hormone (TSH)CLIABXL0152A100T/Kit2-8°CTotal PSA (t-PSA)CLIABXL0132A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP-AbCLIABXL0117A100T/Kit2-8°CTPO-AbCLIABXL0156A100T/Kit2-8°CTRAbCLIABXL0158A100T/Kit2-8°CTA frigger Solution)CLIABXL0158A100T/Kit2-8°CTuberculosis (TB)CLIABXL0125A20ml/bottle2-8°C	T-Beta-HCG		CLIA	BXL0189A	100T/Kit	2-8°C
TG-AbCLIABXL0155A100T/Kit2-8°CThyroid Stimulating Hormone (TSH)CLIABXL0152A100T/Kit2-8°CTotal PSA (t-PSA)CLIABXL0132A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP AbCLIABXL0156A100T/Kit2-8°CTPO-AbCLIABXL0156A100T/Kit2-8°CTRAbCLIABXL0158A100T/Kit2-8°CTS (Trigger Solution)CLIABXL023A200ml/bottle2-8°CTuberculosis (TB)CLIABXL0125A20T/Kit2-8°C	Testosterone		CLIA	BXL0192A	100T/Kit	2-8°C
Thyroid Stimulating Hormone (TSH)CLIABXL0152A100T/Kit2-8°CTotal PSA (t-PSA)CLIABXL0132A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP AbCLIABXL0156A100T/Kit2-8°CTRAbCLIABXL0156A100T/Kit2-8°CTS (Trigger Solution)CLIABXL0158A100T/Kit2-8°CTuberculosis (TB)CLIABXL0125A200ml/bottle2-8°C	тG		CLIA	BXL0157A	100T/Kit	2-8°C
Total PSA (t-PSA)CLIABXL0132A100T/Kit2-8°CTotal-T3 (TT3)CLIABXL0150A100T/Kit2-8°CTotal-T4 (TT4)CLIABXL0151A100T/Kit2-8°CTP AbCLIABXL0117A100T/Kit2-8°CTPO-AbCLIABXL0156A100T/Kit2-8°CTRAbCLIABXL0158A100T/Kit2-8°CTS (Trigger Solution)CLIABXL0203A200ml/bottle2-8°CTuberculosis (TB)CLIABXL0125A20T/Kit2-8°C	тс-АЬ		CLIA	BXL0155A	100T/Kit	2-8°C
Total-T3 (TT3) CLIA BXL0150A 100T/Kit 2-8°C Total-T4 (TT4) CLIA BXL0151A 100T/Kit 2-8°C TP Ab CLIA BXL017A 100T/Kit 2-8°C TPO-Ab CLIA BXL0156A 100T/Kit 2-8°C TKAb 100T/Kit 2-8°C 2-8°C T5 (Trigger Solution) CLIA BXL0158A 100T/Kit 2-8°C Tberculosis (TB) CLIA BXL0125A 200ml/bottle 2-8°C	Thyroid Stimulating Hormone (TSH)		CLIA	BXL0152A	100T/Kit	2-8°C
Total-T4 (TT4) CLIA BXL0151A 100T/Kit 2-8°C TP Ab CLIA BXL0117A 100T/Kit 2-8°C TPO-Ab CLIA BXL0156A 100T/Kit 2-8°C TRAb CLIA BXL0158A 100T/Kit 2-8°C TS (Trigger Solution) CLIA BXL0158A 100T/Kit 2-8°C Tuberculosis (TB) CLIA BXL0125A 200ml/bottle 2-8°C	Total PSA (t-PSA)		CLIA	BXL0132A	100T/Kit	2-8°C
TP Ab CLIA BXL0117A 100T/Kit 2-8°C TPO-Ab CLIA BXL0156A 100T/Kit 2-8°C TRAb CLIA BXL0158A 100T/Kit 2-8°C TS (Trigger Solution) CLIA BXL0158A 100T/Kit 2-8°C Tuberculosis (TB) CLIA BXL0125A 200ml/bottle 2-8°C	Total-T3 (TT3)		CLIA	BXL0150A	100T/Kit	2-8°C
TPO-Ab CLIA BXL0156A 100T/Kit 2-8°C TRAb CLIA BXL0158A 100T/Kit 2-8°C TS (Trigger Solution) CLIA BXL0203A 200ml/bottle 2-8°C Tuberculosis (TB) CLIA BXL0125A 2017/Kit 2-8°C	Total-T4 (TT4)		CLIA	BXL0151A	100T/Kit	2-8°C
TRAb CLIA BXL0158A 100T/Kit 2-8°C TS (Trigger Solution) CLIA BXL0203A 200ml/bottle 2-8°C Tuberculosis (TB) CLIA BXL0125A 20T/Kit 2-8°C	ТР АЬ		CLIA	BXL0117A	100T/Kit	2-8°C
TS (Trigger Solution) CLIA BXL0203A 200ml/bottle 2-8°C Tuberculosis (TB) CLIA BXL0125A 20T/Kit 2-8°C	ТРО-АЬ		CLIA	BXL0156A	100T/Kit	2-8°C
Tuberculosis (TB) CLIA BXL0125A 20T/Kit 2-8°C	TRAb		CLIA	BXL0158A	100T/Kit	2-8°C
	TS (Trigger Solution)		CLIA	BXL0203A	200ml/bottle	2-8°C
Vitamin-D (25-OH-VD) CLIA BXL0199A 100T/Kit 2-8°C	Tuberculosis (TB)		CLIA	BXL0125A	20T/Kit	2-8°C
	Vitamin-D (25-OH-VD)		CLIA	BXL0199A	100T/Kit	2-8°C





Syphilis Testing

Syphilis testing is crucial in diagnosing and managing syphilis, a sexually transmitted infection (STI) caused by the bacterium *Treponema Pallidum* (TP). Early detection is essential as syphilis progresses through stages (primary, secondary, latent, and tertiary), with potentially severe complications if left untreated, including neurological and cardiovascular damage. Common syphilis tests include non-treponemal tests (such as RPR and VDRL), which detect antibodies produced in response to the infection, and treponemal tests (such as FTA-ABS and TPPA), which detect specific antibodies against T. pallidum.

The importance of syphilis testing in diagnostics lies in its role in early detection, allowing timely treatment with antibiotics (usually penicillin) to prevent long-term complications. Testing is also essential for controlling the spread of the infection, especially in pregnant women, as untreated syphilis can lead to congenital syphilis, causing severe harm to the baby. Routine screening in high risk populations helps in public health efforts to reduce syphilis transmission and related health complications.

According to the Center for Disease Control (CDC), the number of cases of Syphilis infection has markedly increased since 1985. Some studies have reported a substantial epidemiological correlation between the acquisition and transmission of the HIV virus and Syphilis. In response to this high demand, Fortress Diagnostics manufactures, develops and supplies an extensive range of high quality syphilis tests with controls and associated reagents.

Featured Products

RPR (Rapid Plasma Reagin)



The Fortress Diagnostics Rapid Plasma Reagin or RPR Card test is a non-treponemal method for the serological detection of syphilis. The antigen– a particulate carbon suspension coated with lipid complexes – agglutinates in the presence of serum reagins. Reagins are antibodies present in the sera of syphilitic patients. Visible agglutination in the form of black clumps which can be viewed macroscopically, indicate the presence of such antibodies in the sample tested.

Syphilis Control Set



The Fortress Diagnostics Syphilis Control Set is designed for the validation of the Fortress range of Syphilis Test Kits. It is recommended that a positive and negative control be included with each run of tests carried out.

TPHA (Treponema Pallidum Haemagglutination)



The Fortress Diagnostics TPHA Test is a sensitive and specific indirect haemagglutination test for the detection of antibodies to Treponema Pallidum. Preserved avian erythrocytes are coated with antigen components of pathogenic T. Pallidum (Nichol's strain). These Test Cells then agglutinate in the presence of specific antibodies to T. Pallidum resulting in the formation of characteristic patterns in microtitration plates. Antibodies to non-pathogenic treponemes are absorbed by an extract of Reiter's treponeme which has been added to the cell suspensions.

VDRL (Venereal Disease Research Laboratory)



The Fortress Diagnostics VDRL test is a non-treponemal flocculation test for the detection and titration of reagins. Reagins are antibodies raised against syphilis induced tissue damage. VDRL Cardiolipin is an alcohol based solution which comprises of three chemically pure components – cardiolipin, lecithin and cholesterol. When the antigen is added to the VDRL buffered saline solution, it forms an emulsion of fine particles which flocculate in the presence of reagins.



Description	Details	Format	Cat No.	Size	Storage
RPR	Reagent Only	LS	SYCA0002	2ml (100T)	2-8°C
RPR	Reagent Only	LS	SYCA0005	5ml (250T)	2-8°C
RPR	Reagent Only	LS	SYCA0010	10ml (500T)	2-8°C
RPR	Reagent Only	LS	SYCA0100	100ml (5000T)	2-8°C
RPR	Reagent Only	LS	SYCA1000	1000ml (50,000T)	2-8°C
RPR	Positive & Negative Control, Pipette/Stirrers, Test Cards, Dispensing Bottle and Needle	LS	SYRPR025	25T	2-8°C
RPR	Positive & Negative Control, Pipette/Stirrers, Test Cards, Dispensing Bottle and Needle	LS	SYRPR050	50T	2-8°C
RPR	Positive & Negative Control, Pipette/Stirrers, Test Cards, Dispensing Bottle and Needle	LS	SYRPR100	100T	2-8°C
RPR	Positive & Negative Control, Pipette/Stirrers, Test Cards, Dispensing Bottle and Needle	LS	SYRPR500	500T	2-8°C
Syphilis Control	Positive & Negative Controls 5 x1.0ml vials of each	LS	SYPN0010	2 x 5 x 1ml	2-8°C
ТРНА	TPHA with +ve and -ve Controls	LS	SYTP0100	100T	2-8°C
ТРНА	TPHA with +ve and -ve Controls	LS	SYTP0200	200T	2-8°C
ТРНА	TPHA Negative Control	LS	STNC0001	1ML	2-8°C
ТРНА	TPHA Positive Control	LS	STPC0001	1ML	2-8°C
VDRL	VDRL Antigen & Buffered Saline Diluent	LS	SYVD0005	1 x 5ml VDRL, 1 x 60ml Buffer	2-8°C







Febrile Antigens

Febrile antigens play an important role in diagnostics by helping to detect infections that cause febrile (feverrelated) illnesses, often associated with diseases like typhoid, brucellosis, and rickettsial infections. Febrile antigen tests involve detecting antibodies in a patient's blood that react to specific bacterial antigens. These tests are useful in diagnosing conditions where fever is a prominent symptom, especially in regions where these infections are endemic.

The importance of febrile antigens in diagnostics lies in their ability to provide a rapid, preliminary diagnosis, aiding in the early identification of bacterial infections that cause fever. Early detection through febrile antigen testing allows for timely treatment, preventing the progression of serious complications. These tests are especially valuable in resource-limited settings where more advanced diagnostics may not be available, and they contribute to better disease management and public health surveillance. Fortress Diagnostics' range of febrile antigens can be used either as a screening tests or as a confirmatory tests.

Featured Products

Febrile Antigen (Widal)



The Fortress Diagnostics Febrile Antigen (Widal) test is a common agglutination test employed in the serological diagnosis of enteric fever and helps to detect the presence of Salmonella antigens in a patient's serum. Patients infected with Salmonella produce antibodies against the antigens of the organism. Antibodies in serum produced in response to exposure to Salmonella organisms will agglutinate bacterial suspension which carries homologous antigens.

Rose Bengal

The Fortress Diagnostics Rose Bengal test is a rapid slide agglutination test for the qualitative and semiquantitative assessment of Brucellosis antibodies. It is one of the easiest to implement and the most widely used for identifying Brucellosis antibodies in human and animal serum. Specific antibodies to the Brucella species are detectable a few weeks after exposure. The stained bacterial suspension agglutinates when mixed with samples containing IgG or IgM antibodies if present in patient samples in concentration ≥25 IU/ml.

Weil-Felix



The Weil-Felix test was developed by Weil and Felix in 1916. It is a heterophile agglutination test due to the antigenic sharing between species of proteus and rickettsia. Antibodies produced against rickettsial antigen cross reacts with OX19 and OX2 strains of Proteus vulgaris and OXK strains of Proteus mirabilis. Proteus stained febrile antigen suspensions can be used to identify and quantitate specific antibodies in human sera following infection with certain Rickettsiae pathogens. Proteus OX19 reacts strongly with the sera of patients with typhus group rickettsae and rocky mountain spotted fever, Proteus OX2 reacts strongly with sera of patients with spotted fever infections, while the Proteus OXK reacts strongly with the sera of patients infected with scrub typhus.

The Fortress Diagnostics Febrile Antigens Weli-Felix test is prepared for the rapid detection and semiquantitation of serum antibodies developed during the acute stage of disease. The antigens agglutinate in the presence of the homologous antibodies in the sample tested.

Description	Details	Format	Cat No.	Size	Storage
Brucella & Proteus +ve Control		LS	FEPBPPC1	1 x 1ml	2-8°C
Brucella Abortus	5ml Stained Antigen Suspension	LS	FEBBAB05	1 x 5ml	2-8°C
Brucella Abortus	1000ml Stained Antigen Suspension	LS	FEBBAB1L	1000ml	2-8°C
Brucella Melitensis	5ml Stained Antigen Suspension	LS	FEBBME05	1 x 5ml	2-8°C
Brucella Melitensis	1000ml Stained Antigen Suspension	LS	FEBBME1L	1000ml	2-8°C
Febrile Antigen Kit (Widal)	8 x 5ml Stained Salmonella Antigens, 100T per Antigen (Without Controls)	LS	FEBNC100	8 x 5ml	2-8°C
Febrile Antigen Kit (Widal)	8 x 5ml Stained Salmonella Antigens, 100T per Antigen (With Controls)	LS	FEBWC100	8 x 5ml/2 x 1ml	2-8°C
Febrile Negative Control	1ml Polyvalent Negative	LS	FEBNCO01	1 x 1ml	2-8°C
Febrile Positive Control	1ml Polyvalent Positive	LS	FEBPCO01	1 x 1ml	2-8°C
Proteus OX19	5ml Stained Antigen Suspension	LS	FEPOX905	1 x 5ml	2-8°C
Proteus OX2	5ml Stained Antigen Suspension	LS	FEPOX205	1 x 5ml	2-8°C
Proteus OXK	5ml Stained Antigen Suspension	LS	FEPOXK05	1 x 5ml	2-8°C
Rose Bengal	2.5ml Rose Bengal, 1ml +ve and -ve Controls, 8x6 Disposable Slides	LS	FEBRB050	50T	2-8°C
Rose Bengal		LS	FEBRB100	100T	2-8°C
Salmonella paratyphi A-H	5ml Stained Antigen Suspension	LS	FEBSAH05	1 x 5ml	2-8°C
Salmonella paratyphi A-O	5ml Stained Antigen Suspension	LS	FEBSAO05	1 x 5ml	2-8°C
Salmonella paratyphi B-H	5ml Stained Antigen Suspension	LS	FEBSBH05	1 x 5ml	2-8°C
Salmonella paratyphi B-O	5ml Stained Antigen Suspension	LS	FEBSBO05	1 x 5ml	2-8°C
Salmonella paratyphi C-H	5ml Stained Antigen Suspension	LS	FEBSCH05	1 x 5ml	2-8°C
Salmonella paratyphi C-O	5ml Stained Antigen Suspension	LS	FEBSCO05	1 x 5ml	2-8°C
Salmonella typhi H	5ml Stained Antigen Suspension	LS	FEBSTH05	1 x 5ml	2-8°C
Salmonella typhi H	1000ml Stained Antigen Suspension	LS	FEBSTH1L	1000ml	2-8°C
Salmonella typhi O	5ml Stained Antigen Suspension	LS	FEBSTO05	1 x 5ml	2-8°C
Salmonella typhi O	1000ml Stained Antigen Suspension	LS	FEBSTO1L	1000ml	2-8°C
Weil-Felix	5ml vials of Brucella Abortus, Brucella Melitensis, Proteus OX19, Proteus OX2 and Proteus OXK	LS	FEWF0025	5 x 5ml	2-8°C





SECTION 7 LATEX SEROLOGY



Latex Serology

Latex serology plays a significant role in diagnostics by utilising latex particles coated with specific antigens or antibodies to detect the presence of corresponding antibodies or antigens in a patient's sample. When the target antibody or antigen is present, the latex particles agglutinate (clump together), which can be visually observed or measured, indicating a positive reaction. This method is commonly used for diagnosing a range of conditions, including infections (e.g., streptococcal infections), autoimmune diseases, and certain cancers.

The importance of latex serology lies in its speed, simplicity, and cost-effectiveness, making it a valuable tool for rapid, point-of-care testing. It is widely used in clinical laboratories for the diagnosis of conditions such as rheumatoid arthritis and meningitis. The test's ability to provide quick results helps clinicians make timely treatment decisions, especially in acute settings. Its versatility and ease of use make it particularly beneficial in resource-limited settings, contributing to improved patient care and disease management. Fortress Diagnostics manufactures, develops and supplies an extensive range of high quality latex serology reagents. Our Latex Serology test kits employ latex particles coated with specific antigens or antibodies to detect the presence of corresponding antibodies or antigens in a sample.

Key Features & Benefits

Fortress Diagnostics' latex serology tests offers several advantages:



Versatility: Latex serology can be adapted to detect a wide range of antigens or antibodies, making it useful for diagnosing a variety of medical conditions.



Cost-Effective: Compared to more advanced diagnostic methods, latex serology is generally low-cost, which can make it more feasible for widespread use.



ASO



The Fortress Diagnostics ASO-Latex agglutination is the rapid and simple test for the qualitative and semiquantitative measurements of antibodies to Anti-Streptolysin-O (ASO) in human serum.


When the latex reagent is mixed with a serum containing ASO agglutination occurs. In acute streptococcal infection the toxic immunogenic exoenzyme Streptolysin-O (ASO) is produced in response to Streptolsin O antigens liberated by haemolytic streptococci of groups A, C and G.

CRP



The Fortress Diagnostics CRP Latex test is a rapid slide agglutination test for the qualitative and semiquantitative detection of C-Reactive Protein in serum. The reagent containing particles coated with specific anti-human C-Reactive protein antibodies, agglutinates in the presence of CRP in the patient's serum.

CRP is the classic acute-phase protein of human serum, synthesised in the liver. Apart from a significant increase in normal serum due to inflammation or acute injury, CRP helps in the management of inflammatory diseases such as rheumatoid arthritis, many malignant diseases and cardiovascular diseases.





Rapid Results: Latex agglutination tests offer rapid results, often within minutes. This makes them useful in urgent clinical situations where timely diagnosis is essential.

Infectious Mononucleosis



The Fortress Diagnostics Infectious Mononucleosis (IM) test is a rapid slide test for the qualitative detection of heterophile antibodies to IM in human serum. The reagent agglutinates when mixed with serum containing the heterophile antibodies.

Infectious Mononucleosis (glandular fever) is an infection usually caused by the Epstein - Barr ViruS (EPV) and primarily affects lymphoid tissues. Mostly children and young adults are affected by this disease. Though specific treatment is rarely required since the disease is usually asymptomatic, symptoms such as inflammation of the liver, enlargement of the spleen, pericarditis, myocarditis and encephalitis associated with this disease requires a physician's attention.

RF



The Fortress Diagnostics RF-Latex test is a rapid slide agglutination test for the qualitative and semiquantitative detection of Rheumatoid Factors in serum. Latex suspension coated with human gamma-globulin, agglutinates in the presence of rheumatoid factors in the patient serum.

Rheumatoid factors are antibodies directed against antigenic sites in the Fc fragment of human and animal IgG. Their frequent occurrence in rheumatoid arthritis makes them useful for diagnosis and monitoring of the disease, although RF is also found in a number of rheumatic disorders such as Sjögren's syndrome and Lupus erythematosus (SLE).

Description	Details	Format	Cat No.	Size	Storage
ASO	5ml Latex	LS	LXASO005	1 x 5ml	2-8°C
ASO	1000ml Latex	LS	LXASO01L	25,000T	2-8°C
ASO	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXASO025	25T	2-8°C
ASO	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXASO050	50T	2-8°C
ASO	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXASO100	100T	2-8°C
ASO	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXASO150	150T	2-8°C
ASO	ASO Latex Negative Control	LS	LXCN0001	1ML	2-8°C
ASO	ASO Latex Positive Control	LS	LAPC0001	1ML	2-8°C
CRP	5ml Latex	LS	LXCRP005	1 x 5ml	2-8°C
CRP	1000ml Latex	LS	LXCRP01L	25,000T	2-8°C
CRP	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXCRP025	25T	2-8°C
CRP	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXCRP050	50T	2-8°C
CRP	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXCRP100	100T	2-8°C
CRP	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXCRP150	150T	2-8°C
CRP	CRP Latex Negative Control	LS	LXCN0001	1ML	2-8°C
CRP	CRP Latex Positive Control	LS	LCPC0001	1ML	2-8°C
hCG White Latex	1000ml Latex	LS	LXHCW01L	25,000T	2-8°C
hCG White Latex	Sensitivity 200IU/ml - White Latex. Kit contains: Latex reagent, +ve and -ve controls, pipettes/stirrers and test slide	LS	LXHCW025	25T	2-8°C



Description	Details	Format	Cat No.	Size	Storage
hCG White Latex	Sensitivity 200IU/ml - White Latex. Kit contains: Latex reagent, +ve and -ve controls, pipettes/stirrers and test slides	LS	LXHCW050	50T	2-8°C
hCG White Latex	Sensitivity 200IU/ml - White Latex. Kit contains: Latex reagent, +ve and -ve controls, pipettes/stirrers and test slides	LS	LXHCW100	100T	2-8°C
Infectious Mononucleosis		LS	LXIM00020	20T	2-8°C
RF	5ml Latex	LS	LXRF0005	1 x 5ml	2-8°C
RF	1000ml Latex	LS	LXRF001L	25,000T	2-8°C
RF	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXRF0025	25T	2-8°C
RF	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXRF0050	50T	2-8°C
RF	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXRF0100	100T	2-8°C
RF	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXRF0150	150T	2-8°C
RF	RF Latex Negative Control	LS	LXCN0001		2-8°C
RF	RF Latex Positive Control	LS	LRPC0001		2-8°C
SLE Latex	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXSLE050	50T	2-8°C
SLE Latex	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXSLE100	100T	2-8°C
Toxoplasmosis	Latex Reagent, +ve and -ve Controls, Pipette/Stirrers & Re- usable Test Slide	LS	LXTX0100	100T	2-8°C





ABOAL ABOAL DWI- DWI- CU AI Contact DWI- DWI- CU AI

Pas

B

NIMI MI

Blood Grouping

Blood grouping is a critical aspect of diagnostics that involves determining a person's blood type based on the presence or absence of specific antigens (A, B, AB, or O) on the surface of red blood cells, as well as the Rhesus (Rh) factor (positive or negative). Blood grouping is essential for safe blood transfusions, organ transplants, and pregnancy care. Incompatible blood transfusions can cause severe, potentially life-threatening immune reactions, such as hemolysis, where the recipient's immune system attacks the donor's blood cells.

The importance of blood grouping in diagnostics lies in its role in preventing complications from transfusion mismatches, ensuring compatibility between donor and recipient in transfusions or transplants, and managing Rh incompatibility during pregnancy. For example, if an Rh-negative mother carries an Rh-positive fetus, the immune response can harm the baby. Blood grouping helps guide appropriate interventions, such as administering Rh immunoglobulin to prevent these complications. It also plays a vital role in emergency care, surgical procedures, and overall patient safety. There are 43 blood group systems containing 345 antigens recognised by the International Society of Blood Transfusion however the most important blood groups in transfusion are the ABO blood group system and the Rh blood group system.

Fortress Diagnostics offers both traditional methods and new techniques to identify the major and rare blood grouping types. Our high-quality blood grouping reagents and gel cards are accurate, easy to use, competitively priced, and conveniently offered in different sizes in order to meet the needs of our distributors, blood banks, clinics, hospital laboratories and transfusion centres across the world.

Description	Details	Format	Cat No.	Size	Storage
AHG (Coomb's)	Anti Human Globulin. A blend of rabbit Anti-Human IgG and Monoclonal Anti-Human c3d	LS	BGAG0010	1 x 10ml	2-8°C
AHG (Coomb's)	Anti Human Globulin. A blend of rabbit Anti-Human IgG and Monoclonal Anti-Human c3d	LS	BGAG001L	1000ml	2-8°C
AHG (Coomb's)	Anti Human Globulin. A blend of rabbit Anti-Human IgG and Monoclonal Anti-Human c3d	LS	BGAG1010	10 x 10ml	2-8°C
AHG (Coomb's)	Anti-Human Globulin, 221 vials x 10ml, packed in trays	LS	BGAG2210	221 x 10ml	2-8°C
Anti-A	Anti-A Colour Coded to International Convention-Acid Blue	LS	BGA00010	1 x 10ml	2-8°C
Anti-A	Anti-A Colour Coded to International Convention-Acid Blue	LS	BGA0001L	1000ml	2-8°C
Anti-A	Anti-A Colour Coded to International Convention-Acid Blue	LS	BGA01010	10 x 10ml	2-8°C
Anti-A	Anti-A Colour Coded to International Convention-Acid Blue, 221 vials x 10ml, packed in trays	LS	BGA02210	221 x 10ml	2-8°C
Anti-A,B	Anti-A,B	LS	BGAB0010	1 x 10ml	2-8°C
Anti-A,B	Anti-A,B	LS	BGAB001L	1000ml	2-8°C
Anti-A,B	Anti-A,B	LS	BGAB1010	10 x 10ml	2-8°C
Anti-A,B	Anti-A,B 221 vials x 10ml, packed in trays	LS	BGAB2210	221 x 10ml	2-8°C
Anti-A1	Lectin	LS	BGA10005	1 x 5ml	2-8°C
Anti-B	Anti-B Colour Coded to International Convention-Acid Yellow	LS	BGB00010	1 x 10ml	2-8°C
Anti-B	Anti-B Colour Coded to International Convention-Acid Yellow	LS	BGB0001L	1000ml	2-8°C
Anti-B	Anti-B Colour Coded to International Convention-Acid Yellow	LS	BGB01010	10 x 10ml	2-8°C
Anti-B	Anti-B Colour Coded to International Convention-Acid Yellow, 221 vials x 10ml, packed in trays	LS	BGB02210	221 x 10ml	2-8°C
Anti-C	Monoclonal	LS	BGBC0005	1 x 5ml	2-8°C
Anti-c	Monoclonal	LS	BGLC0005	1 x 5ml	2-8°C
Anti-CDE	Monoclonal	LS	BGCDE010	1 x 10ml	2-8°C

Blood Grouping | SECTION 8

Description	Details	Format	Cat No.	Size	Storage
Anti-Cw	Monoclonal	LS	BGCW0002	1 x 2ml	2-8°C
Anti-D Blend	Anti-D IgG/IgM Blend	LS	BGD00010	1 x 10ml	2-8°C
Anti-D Blend	Anti-D IgG/IgM Blend	LS	BGD01000	1000ml	2-8°C
Anti-D Blend	Anti-D IgG/IgM Blend	LS	BGD01010	10 x 10ml	2-8°C
Anti-D Blend	Anti-D IgG/IgM Blend, 221 vials x 10ml, packed in trays	LS	BGD02210	221 x 10ml	2-8°C
Anti-E	Monoclonal	LS	BGBE0005	1 x 5ml	2-8°C
Anti-e	Monoclonal	LS	BGLE0005	1 x 5ml	2-8°C
Anti-FYa	Monoclonal	LS	BGFYA002	1 x 2ml	2-8°C
Anti-FYb	Monoclonal	LS	BGFYB002	1 x 2ml	2-8°C
Anti-H	Lectin	LS	BGBH0002	1 x 2ml	2-8°C
Anti-JKa	Monoclonal	LS	BGJKA002	1 x 2ml	2-8°C
Anti-JKb	Monoclonal	LS	BGJKB002	1 x 2ml	2-8°C
Anti-K	Monoclonal	LS	BGBK0005	1 x 5ml	2-8°C
Anti-k	Monoclonal	LS	BGLK0002	1 x 2ml	2-8°C
Anti-KPa	Monoclonal	LS	BGBKPA02	1 x 2ml	2-8°C
Anti-KPb	Monoclonal	LS	BGBKPB02	1 x 2ml	2-8°C
Anti-Lea	Monoclonal	LS	BGLEA002	1 x 2ml	2-8°C
Anti-Leb	Monoclonal	LS	BGLEB002	1 x 2ml	2-8°C
Anti-Lua	Monoclonal	LS	BGLUA002	1 x 2ml	2-8°C
Anti-Lub	Monoclonal	LS	BGLUB002	1 x 2ml	2-8°C
Anti-M	Monoclonal	LS	BGM00002	1 x 2ml	2-8°C
Anti-N	Monoclonal	LS	BGN00002	1 x 2ml	2-8°C
Anti-P1	Monoclonal	LS	BGP10002	1 x 2ml	2-8°C
Anti-S	Monoclonal	LS	BGBS0002	1 x 2ml	2-8°C
Anti-s	Monoclonal	LS	BGLS0002	1 x 2ml	2-8°C
Low Ionic Strength Solution	1000ml LISS. Ready to Use low ionic strength solution	LS	BGLS001L	1000ml	2-8°C
Low Ionic Strength Solution	10x10ml LISS. Ready to Use low ionic strength solution	LS	BGLS1010	10 x 10ml	2-8°C
Low Ionic Strength Solution	221x10ml LISS. Ready to Use low ionic strength solution	LS	BGLS2210	221 x 10ml	2-8°C
Low Ionic Strength Solution	10ml LISS. Ready to Use low ionic strength solution	LS	BGLSS010	1 x 10ml	2-8°C
Monoclonal Grouping Kit	3x10ml of Anti-A, Anti-B, and Anti-D	LS	BGABD030	3 x 10ml	2-8°C
Monoclonal Grouping Kit	4x10ml of Anti-A, Anti-B, Anti-AB and Anti-D	LS	BGABD040	4 x 10ml	2-8°C
Serological Albumin 22%	1000ml of serological Albumin 22%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BG22001L	1000ml	2-8°C
Serological Albumin 22%	10x10ml of serological Albumin 22%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BG221010	10 x 10ml	2-8°C
Serological Albumin 22%	10ml of serological Albumin 22%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BGAL2210	1 x 10ml	2-8°C
Serological Albumin 30%	1000ml of serological Albumin 30%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BG30001L	1000ml	2-8°C
Serological Albumin 30%	10x10ml of serological Albumin 30%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BG301010	10 x 10ml	2-8°C
Serological Albumin 30%	10ml of serological Albumin 30%. Caprylate Free. Free from stabilizers and non-specific agglutinins	LS	BGAL3010	1 x 10ml	2-8°C



Gel Cards

Gel cards, also known as gel columns or gel centrifugation cards, are used primarily in immunohematology for blood typing and cross-matching in transfusion medicine. This technology provides a method for detecting and identifying blood group antigens and antibodies.

Gel cards contain microtubes filled with a gel matrix that acts as a sieve. The principle behind their operation involves the separation of red blood cells (RBCs) and agglutinates (clumps) based on size during centrifugation. The gel matrix can be infused with specific reagents, such as antibodies, to facilitate the detection of antigens or antibodies in the blood sample. Fortress' high-quality blood gel cards are accurate, easy to use, competitively priced, and conveniently offered in different sizes in order to meet the needs of our distributors, blood banks, clinics, hospital laboratories and transfusion centres across the world.

Key Features & Benefits



Reduced working time: The gel card method vastly reduces preparation time due to multi testing and no requirement for washing steps.



Cost effective: Fortress Diagnostics supply gel cards at a competetive price promoting this method as an innovative solution in preference to conventional methods.



Accurate results: Fortress Diagnostics gel cards are proven to provide accurate results in minimal time.



Complete solutions: One test provides comprehensive results for blood grouping compared to multiple tests in conventional tube methods.



Ordering Information

Forward Group 1

- Identifies ABO and Rh D groups by using gel method.
- One card involves 2 tests.Includes control well.

	inc		Concrot	wett.						
		А	В	D	Ctl	А	В	D	Ctl]
	Descri	ption	I		Cat No.		Si	ze	Stora	ge
	Forwa	rd Gro	oup 1		BGGCFC	01-12	12	:T	2-8ºC	
Forward Group 1					BGGCFC	01-24	24	Т	2-8°C	
	Forwa	rd Gro	oup 1		BGGCFG01-48			т	2-8ºC	

Forward Group 3

- Provides the detection of subgroups of A and variants of D on the same gel card.
- Identifies DVI- weak reactive and partial variants of D but not the VI variant of D directly.
 Identifies DVI+ reactive, variant of DVI includes D variants and weak
- Identifies DVI+ reactive, variant or DVI includes D variants and weak D's.
- Identifies the erythrocytes of antibody Rh DVI directly.

А	A1	В	AB	DVI-	DVI+	CDE	Ctl	
Description			(Cat No.		Size	Sto	rage
Forward Gro	oup 3		E	BGGCFG	03-12	12T	2-8	'C
Forward Gro	Forward Group 3		BGGCFG03-24			3-24 24T		'C
Forward Gro	oup 3		E	BGGCFG	03-48	48T	2-8	C

Neonatal Grouping

- Provides applying of ABO, RH D, Rh D VI and direct Coombs test on newborns on one gel card.
- AHG well involves Polyclonal anti-IgG and monoclonal C3 antibodies, IgG well involves monospecific Anti-IgG antibodies.

А	В	AB	DVI-	DVI+	Ctl	AHG	IgE	
Description			Cat No.		Siz	e	Storag	le
Neonatal G	ouping		BGGCNE	EOG-12	12	т	2-8ºC	
Neonatal G	ouping		BGGCNE	EOG-24	24	т	2-8°C	
Neonatal G	ouping		BGGCNE	EOG-48	48	т	2-8°C	

Neutral Gel

 Provides applying of reverse grouping, antibody screening and determining, compability tests.

	NEUT	NEUT	NEUT	NEUT	NEUT	NEUT	NEUT	NEUT	
Des	cription	1		Cat No.		Si	ze	Storag	je
Neu	itral Gel	L		BGGCN	GEL-12	12	T	2-8ºC	
Neutral Gel			BGGCNGEL-24			Т	2-8°C		
Neu	tral Gel	L		BGGCN	GEL-48	48	т	2-8°C	

Forward Group 2

- Identifies ABO and Rh D groups by using gel method.
- One card involves 2 tests.
 Does not include control well, instead includes AB.

А	В	AB	D	А	В	AB	D	
Description			Cat No.		Si	ze	Storag	je
Forward Group 2			BGGCFC	02-12	12	2T	2-8ºC	
Forward Group 2			BGGCFC	02-24	24T		2-8ºC	
Forward Gro	oup 2		BGGCFC	02-48	48	зт	2-8°C	

Forward & Reverse Grouping

- Provides the detection of ABO, Rh D, Rh D VI ve reverse grouping on one gel card.
- Identifies DVI- weak reactive and partial variants of D but not the VI variant of D directly.
- Identifies DVI+ reactive, variant of DVI includes D variants and weak D's.
- Identifies the erythrocytes of antibody Rh DVI directly.
 At the last two wells A1 and B cells are used for reverse apply.

A	В	AB	DVI-	DVI+	Ctl	A1 _{cell}	B _{cell}
Description			c	at No.		Size	Storage
Forward & R	leverse	Groupir	ng E	GGCFRC	51-12	12T	2-8°C
Forward & R	leverse	Groupir	ng E	GGCFRC	51-24	24T	2-8°C
Forward & R	leverse	Groupin	ng B	GGCFRC	61-48	48T	2-8°C

Rh Phenotype +K

 Provides identifying of Rh and K antibodies which have clinic importance by gel method.

С	С	E	е	К	DVI-	DVI+	Ctl	
Description			Cat No.		Sia	ze	Storag	je
Rh Penotype +K			BGGCR⊦	IPK-12	12	Т	2-8ºC	
Rh Penotyp	Rh Penotype +K		BGGCR⊦	IPK-24	24T		2-8°C	
Rh Penotyp	e +K		BGGCRF	IPK-48	48	Т	2-8°C	

A Subgroup

 This test provides identifying of A1 antibody inside erythrocytes by using gel method.

	A1	A1	A1	A1	A1	A1	A1	A1	
Des	cription			Cat No.		Siz	ze	Storag	e
A Su	ıbgroup			BGGCAS	5UB-12	12	т	2-8ºC	
A Subgroup			BGGCASUB-24		24T		2-8°C		
A Su	ıbgroup			BGGCAS	SUB-48	48	Т	2-8°C	



Comp Tests

- The gel card for forward grouping and Cross-match. Provides detection of ABO, Rh mismatch, the existance of antibodies that are against donators erythrocytes in patients serum or the patients erythrocytes in the donor's defi serum.

	А	В	D	Ctl	Enz Major	Enz Minor	AHG	AHG Ctl	
Des	criptior	n		Cat No.		Si	ze	Storag	ge
Соп	np Tests	5		BGGCF	CT1-12	12	!Τ	2-8°C	
Com	np Tests	5		BGGCF	CT1-24	24	IT	2-8ºC	
Com	np Tests	5		BGGCF	CT1-48	48	зт	2-8°C	

DAT Screen

- Provides determining the erythrocytes that get sensitive as in-vivo with IgG or complement C3d.
- Involves IgG well monospesific Anti-IgG, C3d well Anti human C3, AHG well Polyspecific anti-human globuin.

	IgG	C3d	AHG	Ctl	lgG	C3d	AHG	Ctl	
Des	criptior	1		Cat No.		Si	ze	Storag	je
DAT	Screen			BGGCD	ATS-12	12	:T	2-8°C	
DAT	Screen	I		BGGCD	ATS-24	24	Т	2-8°C	
DAT	Screen	1		BGGCD	ATS-48	48	т	2-8°C	

AHG Gel

- Provides applying of Coombs test (DAT) directly, Coombs test (IAT), compatibility, antibody screening and determining tests indirectly.
- All wells include Polyspecific anti-human globulin.

	AHG	AHG	AHG	AHG	AHG	AHG	AHG	AHG	
Des	cription	1		Cat No.		Si	ze	Storag	je
AHG Gel			BGGCAI	HGG-12	12T		2-8ºC		
AHG Gel			BGGCAI	HGG-24	24T		2-8°C		
АНС	i Gel			BGGCAI	HGG-48	48	т	2-8ºC	

Neutral/AHG Gel

Provides applying of Indirect Coombs test (IAT), compatibility, antibody screening and determining tests indirectly.

	NEUT	NEUT	NEUT	NEUT	AHG	AHG	AHG	AHG	
Doc	cription			Cat No.		Siz		Storag	
Description			Cat No.					e	
Neu	tral/AH	G Gel		BGGCNAHG-12		12T		2-8°C	
Neutral/AHG Gel			BGGCNAHG-24		24	Т	2-8°C		
Neutral/AHG Gel			BGGCNAHG-48		48T		2-8°C		

Other Items

Description	Cat No.	Size	Storage
Antibody Screening (IAT)	BGGCAS01		2-8°C
Bromelain Solution	BGGCBS10	10ml	2-8°C
Bromelain Solution	BGGCBS50	50ml	2-8°C
Cell Reverse A1, A2, B, O	BGGCRBC4		2-8°C
Cell Reverse A1, B	BGGCRBC2		2-8°C
Cell Screen 1-2	BGGCSC12		2-8°C
Cell Screen 1-2 Pap	BGGCSC2P		2-8°C
Cell Screen 1-2-3	BGGCSC23		2-8°C
LISS Diluent	BGGCL100	100ml	2-8°C
LISS Diluent	BGGCL500	500ml	2-8°C
Papain Solution	BGGCPAPS	10ml	2-8°C



SECTION 9 RAPID TESTS



Rapid Tests

Rapid tests are essential in diagnostics for providing quick, reliable, and often point-of-care results, enabling timely clinical decisions and interventions. These tests are designed to detect specific biomarkers, antigens, or antibodies in a matter of minutes to hours, rather than days, making them crucial for diagnosing infectious diseases such as COVID-19, HIV, malaria, and influenza. They often involve simple, user-friendly procedures like lateral flow assays or immunochromatographic tests, which require minimal specialised equipment or training.

The importance of rapid tests lies in their ability to facilitate early diagnosis, which can lead to prompt treatment and improved patient outcomes. They are especially valuable in urgent care settings, outbreak management, and remote or resource-limited areas where access to full laboratory infrastructure is limited. Rapid tests help control the spread of contagious diseases by allowing for immediate detection and isolation of infected individuals. Additionally, they play a key role in screening programs, reducing the burden on healthcare systems, and improving access to diagnostic services globally.

Fortress Diagnostics offers a range of rapid diagnostic tests (RDTs) suitable for preliminary or emergency medical screening for use in medical facilities with limited resources and laboratories with low number of tests per day.

Featured Products

CK-MB



The Fortress CK-MB is intended for quantitative determination of creatine kinase-MB in serum or plasma.

The determination of CK MB is an important element in the diagnosis if myocardial ischemia. CK MB values are detectable in serum 3-8 hours after the onset of cardiac symptoms and can be detected over a lengthy period of time. The working stability and range varies with each CK-MB assay, to request an assay IFU contact our technical support team: technical@fortressdiagnostics.com.

Multidrug



The Fortress Diagnostics One Step Multi-Drug Screen Test Dipcard is a lateral flow chromatographic immunoassay for the qualitative detection of multiple drugs and drug metabolites in urine at the following cut-off concentrations in urine. Urine based tests for multiple drugs of abuse range from simple immunoassay tests to complex analytical procedures. The speed and sensitivity of immunoassays have made them the most widely accepted method to screen urine for multiple drugs of abuse.

The Multi-Drug kits are available in 6, 10 and 12 parameters.

HbsAg / HCV / HIV / Syphilis Combo



The Fortress Diagnostics HIV/HCV/HBsAg/Syphilis Combo Rapid Test is a rapid, immunochromatographic assay for the qualitative detection of HCV antibodies, HIV1/2 virus- antibodies, HBsAg, and/or Treponema Pallidum antibodies respectively, in human, serum or plasma samples.

It is intended for professional use as an aid in diagnosing HCV, HIV, HBV and/or Syphilis infections. This assay provides only a preliminary result and all positive specimens should be confirmed with other qualified assays.

Ordering Information (Rapid Test Devices)

Description	Details	Format	Cat No,	Size	Storage
Anti-Hbe	Whole blood/Serum/Plasma	Device	HBEABC20	20T	2-30°C
Anti-Hbe	Whole blood/Serum/Plasma	Device	HBEABC40	40T	2-30°C
Anti-Hbs	Whole blood/Serum/Plasma	Device	HBSAB020	20T	2-30°C
Anti-Hbs	Whole blood/Serum/Plasma	Device	HBSAB040	40T	2-30°C
CEA	Whole blood/Serum/Plasma	Device	CEAC0040	40T	2-30°C
Chagas Antibody Test	Whole blood/Serum/Plasma	Device	CHAGAS20	20T	2-30°C
Chagas Antibody Test	Whole blood/Serum/Plasma	Device	CHAGAS40	40T	2-30°C

Rapid Tests | SECTION 9

Description	Details	Format	Cat No,	Size	Storage
Chlamydia	Swab/Urine,Including sterile swabs, extraction devices & Buffer	Device	RPCH0020	20T	2-30°C
Chlamydia	Swab/Urine,Including sterile swabs, extraction devices & Buffer	Device	RPCH0040	40T	2-30°C
CK-MB, Troponin-I, Myoglobin	Whole Blood, Cut off CK-MB 5ng/ml, Tni-I 1ng/ml, Myo 80ng/ml	Device	CTMC0020	20T	2-30°C
CK-MB, Troponin-I, Myoglobin	Whole Blood, Cut off CK-MB 5ng/ml, Tni-I 1ng/ml, Myo 80ng/ml	Device	CTMC0040	40T	2-30°C
COVID-19 lgG/lgM Rapid Test	(Whole Blood/Serum/Plasma)	Device	COVID010	10T	2-30°C
COVID-19 IgG/IgM Rapid Test	(Whole Blood/Serum/Plasma)	Device	COVID020	20T	2-30°C
Coronavirus Ag	Nasal swab	Device	COVNS020	20T	2-30°C
CRP		Device	CRPC0020	20T	2-30°C
CRP		Device	CRPC0040	40T	2-30°C
Dengue Fever IgG/IgM Antibody Test	Serum/Plasma/Whole Blood, Cut off 500ng/ml	Device	DNGMC020	20T	2-30°C
Dengue Fever IgG/IgM Antibody Test	Serum/Plasma/Whole Blood, Cut off 500ng/ml	Device	DNGMC040	40T	2-30°C
Dengue Fever NS1 Antigen Test		Device	DMNS1020	20T	2-30°C
Dengue Fever NS1 Antigen Test		Device	DMNS1040	40T	2-30°C
Faecal Occult Blood	Cut off 30 ng/ml - (Faeces)	Device	FOBC0020	20T	2-30°C
Faecal Occult Blood	Cut off 30 ng/ml - (Faeces)	Device	FOBC0040	40T	2-30°C
FOB / Transferrin		Device	TRAF0B20	20T	2-30°C
FSH (Menopause)		Device	FSHC0040	40T	2-30°C
Gonorrhoea		Device	GONC0020	20T	2-30°C
Gonorrhoea		Device	GONC0040	40T	2-30°C
HAV Ag One Step		Device	HAVAGC20	20T	2-30°C
HAV IgM		Device	HAVIGM20	20T	2-30°C
HAV IgM		Device	HAVIGM40	40T	2-30°C
Hbe Antigen		Device	HBEAGC20	20T	2-30°C
Hbe Antigen		Device	HBEAGC40	40T	2-30°C
HBsAg /HCV /HIV /SYPHILIS COMBI	Serum/Plasma/Whole Blood	Device	ICOMBI20	20T	2-30°C
HBsAg	Serum/Plasma/Whole Blood - Cut off 1 ng/ml	Device	HBSCWB20	20T	2-30°C
HBsAg	Serum/Plasma/Whole Blood - Cut off 1 ng/ml	Device	HBSCWB40	40T	2-30°C
hCG	Sensitivity 25IU/ml - Serum/Urine, including disposable pipette	Device	HCGCSU20	20T	2-30°C
hCG	Sensitivity 25IU/ml - Serum/Urine, including disposable pipette	Device	HCGCSU40	40T	2-30°C
hCG	Sensitivity 25IU/ml - Urine Only, including disposable pipette	Device	HCGC0020	20T	2-30°C
hCG	Sensitivity 25IU/ml - Urine Only, including disposable pipette	Device	HCGC0040	40T	2-30°C
нси	Serum/Plasma/Whole Blood	Device	HCVC0020	20T	2-30°C
нсv	Serum/Plasma/Whole Blood	Device	HCVC0040	40T	2-30°C
Helicobacter Pylori Ab	Whole blood/Serum/Plasma	Device	HPYC0020	20T	2-30°C
Helicobacter Pylori Ab	Whole blood/Serum/Plasma	Device	HPYC0040	40T	2-30°C
Helicobacter Pylori Ag	Faeces	Device	HPYFC020	20T	2-30°C
Helicobacter Pylori Ag	Faeces	Device	HPYFC040	40T	2-30°C
HEV IgM Rapid	Serum/Plasma	Device	HEVIGM40	40T	2-30°C
HIV 1&2	Serum/Plasma/Whole Blood	Device	HIV00020	20T	2-30°C
HIV 1&2	Serum/Plasma/Whole Blood	Device	HIV00040	40T	2-30°C
Influenze & COVID-19 AG Combo	Nasal Swab	Device	ICAG1920	20T	2-30°C
Influenza & COVID-19 AG Combo	Nasal Swab	Device	ICAG1940	40T	2-30°C
LH (Ovulation)	Serum/Plasma/Whole Blood	Device	LHOV0020	20T	2-30°C
Malaria P.falciparum	Serum/Plasma/Whole Blood, with Test Tube & Buffer	Device	PFC00020	20T	2-30°C
Malaria P.falciparum	Serum/Plasma/Whole Blood, with Test Tube & Buffer	Device	PFC00040	40T	2-30°C
	ce, sing, asing where blood, where set table a barrer	Device		.01	- 30 C

SECTION 9 | Rapid Tests

Description	Details	Format	Cat No,	Size	Storage
Malaria P.falciparum/P. Vivax (Pan Malaria Device)	Serum/Plasma/Whole Blood, with Test Tube & Buffer	Device	PVC00020	20T	2-30°C
Malaria P.falciparum/P. Vivax (Pan Malaria Device)	Serum/Plasma/Whole Blood, with Test Tube & Buffer	Device	PVC00040	40T	2-30°C
Monkeypox IgG/IgM	Serum/Plasma/Whole Blood	Device	MPOX0020	20T	2-30°C
Monkeypox IgG/IgM	Serum/Plasma/Whole Blood	Device	MPOX0040	40T	2-30°C
Multi-drug (6 Parameter)	Test Device for multi-drug detection- Urine. (Many combinations available)	Device	DOAC0006	20T	2-30°C
Multi-drug (10 Parameter)	Test Device for multi-drug detection- Urine- COC, AMP, MET,THC,MTD,MDMA,OPI,PCP,BAR,BZO.	Device	DOAC0010	20T	2-30°C
Multi-drug (12 Parameter)	Test Device for multi-drug detection- Urine- COC, AMP, TCA, MOP, MET,THC,MTD,MDMA,OPI,PCP,BAR,BZO.	Device	DOAC0012	20T	2-30°C
PSA	Serum/Plasma/Whole Blood Cut off - 3 ng/ml	Device	PSA00020	20T	2-30°C
PSA	Serum/Plasma/Whole Blood Cut off - 3 ng/ml	Device	PSA00040	40T	2-30°C
S.Typhi Antigen		Device	STAG0020	20T	2-30°C
Strep A		Device	STREPA20	20T	2-30°C
Syphilis	Serum/Plasma/Whole Blood	Device	TPC00020	20T	2-30°C
Syphilis	Serum/Plasma/Whole Blood	Device	TPC00040	40T	2-30°C
Troponin-I	Serum/Plasma/Whole Blood, Cut off 1 ng/ml	Device	TNIC0010	10T	2-30°C
Troponin-I	Serum/Plasma/Whole Blood, Cut off 1 ng/ml	Device	TNIC0020	20T	2-30°C
Tuberculosis	Serum/Plasma/Whole Blood	Device	TBC00020	20T	2-30°C
Tuberculosis	Serum/Plasma/Whole Blood	Device	TBC00040	40T	2-30°C
Typhoid Antibody	Serum/Plasma/Whole Blood	Device	TYP00020	20T	2-30°C
Typhoid Antibody	Serum/Plasma/Whole Blood	Device	TYP00040	40T	2-30°C

Ordering Information (Rapid Test Strips)

Description	Details	Format	Cat No,	Size	Storage
Anti-HBs	Serum/Plasma - Cut off 30 mIU/ml	Strips	HBSAB050	50T	2-30°C
Anti-HBs	Serum/Plasma - Cut off 30 mIU/ml	Strips	HBSAB100	100T	2-30°C
HBsAg	Serum/Plasma/Whole Blood - Cut off 1 ng/ml	Strips	HBSWB050	50T	2-30°C
HBsAg	Serum/Plasma/Whole Blood - Cut off 1 ng/ml	Strips	HBSWB100	100T	2-30°C
hCG (Urine & Serum)	Sensitivity 25IU/ml - Serum/Urine, including disposable pipette	Strips	HCGSU050	50T	2-30°C
hCG (Urine & Serum)	Sensitivity 25IU/ml - Serum/Urine, including disposable pipette	Strips	HCGSU100	100T	2-30°C
hCG (Urine Only)	Sensitivity 25IU/ml - Urine -2.5mm	Strips	HCGS0050	50T	2-30°C
hCG (Urine Only)	Sensitivity 25IU/ml - Urine -2.5mm	Strips	HCGS0100	100T	2-30°C
нсv	Serum/Plasma/Whole Blood	Strips	HCVS0050	50T	2-30°C
нси	Serum/Plasma/Whole Blood	Strips	HCVS0100	100T	2-30°C
Helicobacter Pylori	Whole Blood/Serum/Plasma	Strips	HPS00050	50T	2-30°C
Helicobacter Pylori	Serum/Plasma	Strips	HPS00100	100T	2-30°C
HIV 1&2	Serum/Plasma/Whole Blood	Strips	HIVS0050	50T	2-30°C
HIV 1&2	Serum/Plasma/Whole Blood	Strips	HIVS0100	100T	2-30°C
Syphilis	Serum/Plasma/Whole Blood	Strips	TPS00050	50T	2-30°C
Syphilis	Serum/Plasma/Whole Blood	Strips	TPS00100	100T	2-30°C
Urine (11 Parameter)		Strips	URS00100	100T	2-30°C



-

SECTION 10 HAEMATOLOGY REAGENTS

SECTION 10 | Haematology Reagents

Haematology Reagents

Hematology reagents play a vital role in the field of diagnostics, particularly in the analysis of blood and its components. These reagents include a variety of substances, such as stains, anticoagulants, and controls, that facilitate the identification and quantification of different blood cell types, hemoglobin levels, and other hematological parameters. They are essential for conducting tests like complete blood counts (CBC), blood smears, and coagulation studies.

The importance of hematology reagents lies in their contribution to accurate and reliable diagnostic results, which are crucial for diagnosing a wide range of conditions, including anemia, infections, blood disorders, and clotting abnormalities. High-quality reagents ensure the consistency and reproducibility of test results, helping clinicians make informed decisions regarding patient management and treatment strategies. Additionally, these reagents are integral in monitoring the effectiveness of therapies, assessing the progression of diseases, and conducting routine health screenings. Overall, hematology reagents are essential tools that support effective patient care and enhance the overall quality of laboratory diagnostics.

Fortress Diagnostics has developed a range of haematology reagents, controls and calibrators for use in detecting blood related disorders.

Featured Product

Haematology Tri-level Controls



Fortress Diagnostics has developed a range of haematology reagents, controls and calibrators for use in detecting blood related disorders.

The Fortress Hematology Control (low, normal, high) kits are intended for use in monitoring the accuracy and precision of blood cell counts on haematology analysers. They have a shelf life of 6 months.

Ordering Information

Description	Details	Format	Cat No.	Size	Storage
Celldiff+ Clean		LS	HAEM042B	5 Litres	2-8°C
Celldiff+ Lyse		LS	HAEM043A	1 Litre	2-8°C
Celldiff+ Clean		LS	HAEM042A	1 Litre	2-8°C
Celldiff+ Diluent		LS	HAEM034A	20 Litres	2-8°C
Celldiff+ Enhanced Cleaner	Daily & weekly Maintenance Solution	LS	HAEM033A	100mls	2-8°C
Celldiff-3+ Reagent Starter Pack	Reagents for Celldiff-3	LS	FORTSP01	5 Litres Diluent; 1 Litre Cleaner; 100mls Lyse.	2-8°C
Haematology Calibrator	Haematology calibrator for 3-part differential Analysers	LS	HAEMC005	3 x 3ml	2-8°C
Haematology Control (3 Levels)	Haematology control for 3-part differential Analysers	LS	HAEMC004	3 x 3 x 3ml	2-8°C
Haematology Control High	Haematology control for 3-part differential Analysers	LS	HAEMC003	3 x 3ml	2-8°C
Haematology Control Low	Haematology control for 3-part differential Analysers	LS	HAEMC001	3 x 3ml	2-8°C
Haematology Control Normal	Haematology control for 3-part differential Analysers	LS	HAEMC002	3 x 3ml	2-8°C
Haematology Triple Pack Box	Reagents for 1000 tests on Celldiff-3	LS	HAEM1000	20 Litres Diluent; 1 Litre Lyse; 5 Litres Cleaner	2-8°C
Proteolyse	Enhanced Cleaner for Clot Removal & Aperture Cleaning	LS	HAEM035A	50mls	2-8°C



SECTION 11 HAEMOSTASIS

Haemostasis

Haemostasis coagulation reagents are vital tools in the diagnosis and management of bleeding and thrombotic disorders, serving as essential components in various laboratory tests that assess the coagulation system's functionality. These reagents, including anticoagulants, clotting factor assays, and controls, are used in tests such as prothrombin time (PT), activated partial thromboplastin time (aPTT), and fibrinogen assays. They enable healthcare professionals to evaluate how effectively blood clots form, which is critical for diagnosing conditions like hemophilia, von Willebrand disease, and other coagulopathies. Accurate results from these tests guide clinical decision-making, helping to determine the appropriate treatment plans, monitor disease progression, and assess the risk of excessive bleeding or thrombosis in patients.

In addition to their diagnostic importance, haemostasis coagulation reagents play a crucial role in monitoring patients undergoing anticoagulant therapy, such as those receiving warfarin or heparin. Regular monitoring using these reagents ensures that patients maintain therapeutic levels of anticoagulation while minimizing the risk of complications. Furthermore, high-quality coagulation reagents are essential for maintaining laboratory standards and ensuring reliable results, which are critical for patient safety. Their role extends to research and development, where they help advance our understanding of coagulation pathways and contribute to the development of new diagnostic tests and treatments for coagulation disorders. Overall, haemostasis coagulation reagents are indispensable for effective patient care and improving health outcomes in individuals with coagulation-related conditions.

Fortress Diagnostics manufactures, develops and supplies an extensive range of high quality haemostasis coagulation reagents.

Featured Product

PT (Thromboplastin)



The Fortress Diagnostics PT reagent is an in vitro diagnostic assay intended for use in performing the one stage prothrombin time (PT) test and assays which are based on a modified prothrombin time.

Prothrombin time is the method of choice for monitoring oral anticoagulation therapy and is a fundamental screening test for acquired or inherited bleeding disorders. During oral anti-coagulation therapy, the activity of vitamin K-dependent clotting factors (II, VII, IX, X, Protein C and Protein S) is reduced and PT time is increased. The test is used for quantitative determination of blood clotting factors in the extrinsic (VII) and common pathways (II, V and X) of coagulation.

Ordering Information

Description	Details	Format	Cat No.	Size	Storage
APTT Reagent		LS	COAG103A	1 x 3ml	2-8°C
APTT Reagent		LS	COAG103B	5 x 3ml	2-8°C
APTT Reagent		LS	COAG103C	1 x 10ml	2-8°C
APTT Reagent with Calcium Chloride		LS	COAG110A	6 x 3ml; 2 x 10ml	2-8°C
Calcium Chloride (0.02M)	Ready To Use for APTT	LS	COAG104A	1 x 10ml	2-8°C
Calcium Chloride (0.02M)	Ready To Use for APTT	LS	COAG104B	5 x 10ml	2-8°C
D-Dimer Latex	D-Dimer Latex Assay	LS	COAG115C	25T	2-8°C
D-Dimer Latex	D-Dimer Latex Assay	LS	COAG115D	50T	2-8°C
D-Dimer Latex	D-Dimer Latex Assay	LS	COAG115E	100T	2-8°C
Factor II Assay Kit		LS	COAG129A	5 x 1ml; 1 x 60ml	2-8°C
Factor IX Assay Kit		LS	COAG127A	5 x 1ml; 1 x 60ml	2-8°C
Factor V Assay Kit		LS	COAG128A	5 x 1ml; 1 x 60ml	2-8°C
Factor VII Assay Kit		LS	COAG124A	5 x 1ml; 1 x 60ml	2-8°C
Factor VIII Assay Kit		LS	COAG122A	5 x 1ml; 1 x 60ml	2-8°C
Factor X Assay Kit		LS	COAG123A	5 x 1ml; 1 x 60ml	2-8°C
Factor XI Assay Kit		LS	COAG133A	5 x 1ml; 1 x 60ml	2-8°C
Factor XII Assay Kit		LS	COAG126A	5 x 1ml; 1 x 60ml	2-8°C

Haemostasis | SECTION 11

Description	Details	Format	Cat No.	Size	Storage
Fibrinogen	Quantitative Estimation of Fibrinogen	LS	COAG116A	1 x 10ml Buffer; 1 x 2ml Bovine Thrombin; 1 x 1ml Calibrator	2-8°C
Fibrinogen	Quantitative Estimation of Fibrinogen	LS	COAG116B	1 x 20ml Buffer; 2 x 2ml Bovine Thrombin; 1 x 1ml Calibrator	2-8°C
Fibrinogen	Quantitative Estimation of Fibrinogen	Lyo	COAG105A	20T	2-8°C
Fibrinogen		Lyo	COAG105B	60T	2-8°C
Fibrinogen		Lyo	COAG105C	100T	2-8°C
Fibrinogen Degradation Products (FDP)	Latex Assay	LS	COAG114A	60T	2-8°C
Haemostasis Control Normal (Including Factor Assays)		LS	COAG131A	1ml	2-8°C
Haemostasis Control Pathological (Including Factor Assays)		LS	COAG132A	1ml	2-8°C
Kaolin Clotting Time	For the Determination of Kaolin Clotting Time	LS	COAG121A	R1: 1 x 2ml R2: 1 x 2ml	2-8°C
Lupus Anticoagulants (LA)	For screening & Confirmation of Lupus Anticoagulants	LS	COAG106A	2 x 1ml; 1 x 2.5ml	2-8°C
Plasma Control (Level 2)	For Coagulation Series only	LS	COAG109B	5 x 1ml	2-8°C
Plasma Control (Level 1 & Level 2)	For Coagulation Series only	LS	COAG112A	2 x 5 x 1ml	2-8°C
Plasma Control (Level 1 & Level 2)	For Coagulation Series only	LS	COAG112B	20 x 5 x 1ml	2-8°C
Plasma Control (Level 1)	For Coagulation Series only	LS	COAG108A	1 x 1ml	2-8°C
Plasma Control (Level 1)	For Coagulation Series only	LS	COAG108B	5 x 1ml	2-8°C
Plasma Control (Level 2)	For Coagulation Series only	LS	COAG109A	1 x 1ml	2-8°C
Protein C	Quantitative Determination of Protein C in Human Plasma	LS	COAG118A	40T	2-8°C
Protein S	Quantitative Determination of Protein S in Human Plasma	LS	COAG119A	40T	2-8°C
PT High Sensitivity	ISI 1.0	Lyo	COAG111A	5 x 5 ml	2-8°C
PT High Sensitivity	ISI 1.0	Lyo	COAG111B	10 x 5 ml	2-8°C
PT High Sensitivity	ISI 1.15	LS	COAG101A	1 x 5ml	2-8°C
PT High Sensitivity	ISI 1.16	LS	COAG101B	5 x 5ml	2-8°C
PT High Sensitivity	ISI 1.17	LS	COAG101C	10 x 5ml	2-8°C
PT High Sensitivity	ISI 1.18	LS	COAG101D	6 x 8ml	2-8°C
PT High Sensitivity	ISI 1.19	LS	COAG101E	6 x 5ml	2-8°C
PT High Sensitivity	ISI 1.20	LS	COAG101F	20 x 5ml	2-8°C
PT High Sensitivity	ISI 1.21	LS	COAG101G	20 x 8ml	2-8°C
PT Low Sensitivity	ISI 1.5	LS	COAG102A	1 x 5ml	2-8°C
PT Low Sensitivity	ISI 1.6	LS	COAG102B	5 x 5ml	2-8°C
PT Low Sensitivity	ISI 1.8	LS	COAG117A	1 x 5ml	2-8°C
PT Low Sensitivity		LS	COAG117B	5 x 5ml	2-8°C
Russell Viper Venom Kit (dRvvT)		LS	COAG130A	2 x 1ml; 1 x 2.5ml	2-8°C
Thrombin Time Reagent	For Qualitative estimation of Fibrinogen	LS	COAG107A	5 x 1ml	2-8°C



Urinalysis

Urinalysis is a fundamental diagnostic tool that plays a crucial role in the evaluation of a patient's overall health and the diagnosis of various medical conditions. This non-invasive test involves the physical, chemical, and microscopic examination of urine, providing valuable insights into renal function, hydration status, and the presence of diseases. Urinalysis can detect a wide range of abnormalities, including infections, kidney disease, metabolic disorders, and urinary tract disorders. By analysing parameters such as protein, glucose, ketones, blood, and the presence of bacteria or crystals, healthcare providers can identify underlying issues early, allowing for timely intervention and treatment.

In addition to its diagnostic value, urinalysis serves as a vital monitoring tool for patients with known health conditions, such as diabetes or chronic kidney disease. Regular urinalysis helps track disease progression, assess treatment effectiveness, and detect complications before they become more serious. Furthermore, urinalysis is commonly used in routine health screenings, drug testing, and pre-employment evaluations, making it a versatile and widely applicable tool in clinical practice. Its ease of collection, low cost, and rapid turnaround time enhance its importance in healthcare, ultimately contributing to improved patient care and outcomes.

The Fortress Diagnostics 11 Parameter Urine Strips are intended for the screening of detection of diabetes, metabolic abnormalities, liver diseases, billiary and hepatic obstructions, haemolytic diseases and diseases of kidney and urinary tract infections.

Ordering Information

Description	Details	Format	Cat No.	Size	Storage
Urine Strips (11 Parameter)	Urobilinogen, Blood, Ketone, Glucose, Protein, pH, Bilirubin, Nitrite, SG, Leucocytes, Ascorbic Acid	Strips	URS00100	100T	2-8°C





FOOD & FEED SAFETY

Food & Feed Safety

Food and feed safety is crucial in diagnostics as it ensures the integrity and quality of food products and animal feed, ultimately protecting public health and agricultural systems. Effective diagnostic testing plays a vital role in identifying pathogens, contaminants, and chemical residues in food and feed supplies, which can pose significant risks to human health and animal welfare.

By employing various analytical techniques, including microbiological, chemical, and molecular assays, diagnostic laboratories can detect harmful substances such as bacteria (e.g., Salmonella, E. coli), viruses, mycotoxins, and pesticides. This proactive approach not only helps prevent foodborne illnesses and outbreaks but also supports regulatory compliance and industry standards, ensuring that food products are safe for consumption.

Fortress Diagnostics offers a wide range of ELISA and Rapid Lateral Flow testing kits.

Ordering Information

Antibiotics

Fortress Diagnostics have developed a range of ELISA and Rapid Lateral Flows test kits to detect antibiotics in a range of food matrices including: Meat & Poultry, Seafood, Honey, Feed Eggs and Dairy. Testing food for antibiotics helps suppliers comply with regulations and ensures the safe supply of food products to consumers.

Description	Details	Format	Cat No.	Size	Storage
AHD FAST	ELISA		BXEFB43A	96T	2-8 °C
AMOZ FAST	ELISA		BXEFB42A	96T	2-8°C
AOZ FAST	ELISA		BXEFB41A	96T	2-8°C
Chloramphenicol FAST	ELISA		BXEFB03A	96T	2-8°C
Multi-Sulfonamide	ELISA		BXEFT08A	96T	2-8°C
Nitroimidazole (Dimetridazole)	ELISA		BXEFB45A	96T	2-8°C
Nitroimidazole (Metronidazole)	ELISA		BXEFB40A	96T	2-8°C
Oxytetracycline/Tetracycline	ELISA		BXEFB46A	96T	2-8°C
Quinolone FAST	ELISA		BXEFB07A	96T	2-8°C
SEM FAST	ELISA		BXEFB44A	96T	2-8°C
Streptomycin FAST	ELISA		BXEFB37A	96T	2-8°C
Tylosin	ELISA		BXEFT13A	96T	2-8°C

FlowSense

The Flowsense product range offers a cost-effective and user-friendly rapid test solution for contaminant testing within the food industry. Unlike traditional test solutions, Flowsense testing can be completed outside a laboratory setting.

With minimal initial investment and technical expertise required, the Flowsense product range enhances the frequency of testing within food production quality schedules, enabling more consistent monitoring and quicker responses to any issues that may arise. By incorporating regular testing, food producers can maintain higher standards of quality and compliance, ultimately benefiting the overall operational efficiency and product safety.

FlowSense Benefits for End Users:

- Minimal initial investment required.
- Eliminate expenses associated with third-party testing.
- No prior experience or extensive technical knowledge needed.
- Portable reader for convenient use.
- Enhance frequency of testing within your quality schedule.

FlowSense Honey Product Features:

- Innovative Multiplex detection capability (3-in-1 and 4-in-1 test strips).
- Straightforward sample preparation process.
- Ability to detect 49 antibiotic compounds.
- Quick 15-minute incubation for Nitrofuran derivatisation.
- Compliance with MMPRs Council Directive 96/23/EC.

Description	Details	Format	Cat No.	Size	Storage
Streptomycin Honey Flowsense Rapid Test	Rapid Lateral Flow		BXEFB55A	48T	2-8°C
Multi-Sulfonamides Honey Flowsense Rapid Test (High Sensitivity)	Rapid Lateral Flow (Multiplex)		BXEFB56A	48T	2-8°C
Multi-Antibiotic Honey (Quin/Thiam/ Spec) Flowsense Rapid Test	Rapid Lateral Flow (Multiplex)		BXEFB57A	48T	2-8°C
Multi-Antibiotic Honey (Eryt/Tylo/Tetra) Flowsense Rapid Test	Rapid Lateral Flow (Multiplex)		BXEFB58A	48T	2-8°C
Chloramphenicol Honey Flowsense Rapid Test	Rapid Lateral Flow		BXEFB59A	48T	2-8°C
Multi-Nitrofuran Honey Flowsense (AOZ, AMOZ, AHD, SEM) Rapid test	Rapid Lateral Flow (Multiplex)		BXEFB60A	48T	2-8°C

Honey Quality Parameters

To determine honey quality measurements, Fortress Diagnostics has developed a range of enzymatic tests. These tests offer honey producers the ability to determine not only quality, but also shelf life and ability to hold moisture.

Description	Details	Format	Cat No.	Size	Storage
Diastase* Phadebas	Instrumentation		1321	50T	2-8°C
Diastase* Phadebas	Colorimetric		1322	500T	2-8°C
Glucose Controls (Low and Elevated)	Colorimetric		BXEFC04A	96T	2-8°C
Glucose/Fructose	Colorimetric		BXEFC02C	50T	2-8°C
НМ	Colorimetric		BXEFC01A	100T	2-8°C

Growth Hormones

Fortress Diagnostics have developed a range of ELISA's for the detection of growth hormones in food. There are a number of regulations in place globally to restrict the level of growth hormones administered to animals, with some being completely banned. Regular testing for growth hormones ensures suppliers conform to regulations and deliver safe products to end users.

Description	Details	Format	Cat No.	Size	Storage
Clenbuterol FAST	ELISA		BXEFB04A	96T	2-8°C
Ractopamine	ELISA		BXEFB05A	96T	2-8°C
Trenbolone	ELISA		BXEFT15A	96T	2-8°C
Zeranol	ELISA		BXEFT17A	96T	2-8°C

SECTION 13 | Food & Feed Safety

Mycotoxins

Mycotoxins are naturally occurring toxins that are often detected in a variety of feed and crops including; cereal, nuts and spices, most often in warm and humid conditions. Contaminated feed can have serious health effects when consumed, making it important to test and detect mycotoxins before any animal or human consumption.

Fortress Diagnostics have a range of kits to detect Total Aflatoxins, Alfatoxin B1, Aflatoxin M1 and Ochratoxin.

Description	Details	Format	Cat No.	Size	Storage
Aflatoxin B1	ELISA		BXEFT01A	96T	2-8°C
Aflatoxin M1	ELISA		BXEFB15A	96T	2-8°C
Aflatoxin Total (1-20 PPB)	ELISA		BXEFB09A	96T	2-8°C
DON deoxynivalenol (vomitoxin)	ELISA		BXEFT18A	96T	2-8°C
Fumonisin	ELISA		BXEFB38A	96T	2-8°C
Ochratoxin	ELISA		BXEFB52A	96T	2-8°C
Т2/НТ2	ELISA		BXEFT20A	96T	2-8°C
Zearalenone	ELISA		BXEFT19A	96T	2-8°C

Rapid Test (FlowSense) Mycotoxin Tests Coming Soon

Mycotoxins - Standard Solutions

Mycotoxin standard solutions are precisely prepared solutions containing known concentrations of specific mycotoxins. These standard solutions are used in laboratory settings for the calibration and validation of analytical methods, ensuring accurate detection and quantification of mycotoxins in various samples such as food, feed, and biological fluids.

Description	Purity	Solvent	Cat No.	Raw Materials	Storage
15-Acetyl DON Solution 100ug/ml	94.7%	Acetonitrile (HPLC grade Supragradient, 99.97%): water, 1:1	BXEFF13A	15-Acetyl Deoxynivalenol	2-8°C
3-Acetyl DON Solution 100ug/ml	99.66%	Acetonitrile HPLC grade Supragradient, 99.97%	BXEFF12A	3-Acetyl-DON	2-8°C
Aflatoxin B1 Solution 2ug/ml	99.90%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF02A	Aflatoxin B1 From Aspergillus flavus	2-8°C
Aflatoxin B2 Solution 0.5ug/ml	100%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF03A	Aflatoxin B2 From Aspergillus flavus	2-8°C
Aflatoxin G1 Solution 2ug/ml	98.57%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF04A	Aflatoxin G1 From Aspergillus flavus	2-8°C
Aflatoxin G2 Solution 0.5ug/ml	99.80%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF05A	Aflatoxin G2 From Aspergillus flavus	2-8°C
Aflatoxin M1 Solution 0.5ug/ml	99.89%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF06A	Aflatoxin M1 From Aspergillus flavus	2-8°C
DON Solution 100ug/ml	99.72%	Acetonitrile HPLC grade Supra-gradient, 99.97%	BXEFF08A	Deoxynivalenol	2-8°C
Fumonisin B1 Solution 50ug /ml	100%	Acetonitrile (HPLC grade Supra-gradient, 99.97%): water, 1:1	BXEFF09A	Fumonisin B1 from Fusarium moniliforme	2-8°C
Fumonisin B2 Solution 50ug/ml	100%	Acetonitrile (HPLC grade Supra-gradient, 99.97%): water, 1:1	BXEFF10A	Fumonisin B2 from Fusarium moniliforme	2-8°C
Fumonisin B3 Solution 50ug/ml	>99%	Acetonitrile (HPLC grade Supragradient, 99.97%): water, 1:1	BXEFF11A	Fumonisin B3 from Fusarium moniliforme	2-8°C
Nivalenol Solution 100ug/ml	99.66%	Acetonitrile HPLC grade Supragradient, 99.97%	BXEFF14A	Nivalenol from Fusarium SP	2-8°C
Ochratoxin A Solution 10ug/ml	>99.9%	Acetonitrile HPLC grade Supragradient, 99.97%	BXEFF01A	Ochratoxin A, Benzene Free, From Aspergillus Ochraceus	2-8°C
Zearalenone Solution 100ug/ml	98.97%	Acetonitrile HPLC grade Supragradient, 99.97%	BXEFF07A	Zearalenone from Giberella Zeae	2 - 8 °C

Allergens



Fortress Diagnostics have a range of Food Allergen Monoclonal Antibody-Based Enzyme-Linked Immunosorbent assay (ELISA) kits available. The highly sensitive ELISA's deliver quantitative results for the detection of food allergens in a wide range of food matrices. The kits can also be used for qualitative, yes/no results for simple assessments.

Allergen kits are available for the detection of; almond, brazil nut, buckwheat, casein, cashew, coconut, crustacean, egg, gluten, hazelnut, lupin e, macadamia, mustard, peanut, peacan, pine nut, pistachio, sesame seed, soya and walnut.

Group	Description	Method	Cat No.	Size	Storage
Fac	Fac	ELISA	EOM-EK-96	96T	2-8°C
Egg	Egg	ELISA	EOM-EK-48	48T	2-8°C
Cluber	Cluber	ELISA	GLU-EK-96	96T	2-8°C
Gluten	Gluten	ELISA	GLU-EK-48	48T	2-8°C
	Lucia a	ELISA	LU2-EK-96	96T	2-8°C
	Lupin e	ELISA	LU2-EK-48	48T	2-8°C
Logumo	Descut	ELISA	PA3-EK-96	96T	2-8°C
Legume	Peanut	ELISA	PA3-EK-48	48T	2-8°C
	Four	ELISA	SG6-EK-96	96T	2-8°C
	Soya	ELISA	SG6-EK-48	48T	2-8°C
Milk	Casein	ELISA	CAS-EK-96	96T	2-8°C
MICK	Casein	ELISA	CAS-EK-48	48T	2-8°C
	Almond	ELISA	AP1-EK-96	96T	2-8°C
	Autona	ELISA	AP1-EK-48	48T	2-8°C
	Brazil Nut	ELISA	BN-EK-96	96T	2-8°C
	Diazitinut	ELISA	BN-EK-48	48T	2-8°C
	Cashew	ELISA	CA2-EK-96	96T	2-8°C
	Casilew	ELISA	CA2-EK-48	48T	2-8°C
	Coconut	ELISA	CN1-EK-96	96T	2-8°C
	coconac	ELISA	CN1-EK-48	48T	2-8°C
	Hazelnut	ELISA	НС9-ЕК-96	96T	2-8°C
Tree nut	Hazethut	ELISA	НС9-ЕК-48	48T	2-8°C
iiee iide	Macadamia	ELISA	MN1-EK-96	96T	2-8°C
	Macadalina	ELISA	MN1-EK-48	48T	2-8°C
	Pecan	ELISA	PC4-EK-96	96T	2-8°C
	recuir	ELISA	PC4-EK-48	48T	2-8°C
	Pine nut	ELISA	PN1-EK-96	96T	2-8°C
	r nie nie	ELISA	PN1-EK-48	48T	2-8°C
	Pistachio	ELISA	PV1-EK-96	96T	2-8°C
	, istatilio	ELISA	PV1-EK-48	48T	2-8°C
	Walnut	ELISA	WJ4-EK-96	96T	2-8°C
	Wathat	ELISA	WJ4-EK-48	48T	2-8°C



Group	Description	Method	Cat No.	Size	Storage
	Buckwheat	ELISA	BW1-EK-96	96T	2-8°C
	Buckwheat	ELISA	BW1-EK-48	48T	2-8°C
Seeds	Mustard	ELISA	MS1-EK-96	96T	2-8°C
Seeds	Muscalu	ELISA	MS1-EK-48	48T	2-8°C
	Sesame Seed	ELISA	SE1-EK-96	96T	2-8°C
	Sesame Seed	ELISA	SE1-EK-48	48T	2-8°C

SECTION 14 INSTRUMENTATION

Q

•



Instrumentation

In today's evolving and vast healthcare climate, clinics and laboratories around the globe face the challenges of raising patient care, whilst reducing operating costs.

Fortress Diagnostics' diverse range of analysers offer laboratories the oppurtunity to provide affordable results without compromising quality. We offer:

- **Scalability Solutions:** Meeting the needs of laboratories ranging from small scale independent facilities to large scale clinical settings, Fortress diagnostics' range of instruments can match the specified testing requirements.
- **Efficiency and Productivity:** Our range of analysers can improve sample analysis workflow and turnaround time. Ergonomic connectivity between automation, IT and the user ensures efficiency in all disciplines of analysis.

The Fortress Instrumentation Range:

- Clinical Chemistry Analysers
- Coagulation Analysers
- Electrolyte Analysers
- Food & Feed Analysers
- Gel Card Analysers
- Haematology Analysers
- Immunoassay (CLIA/ELISA) Analysers




Analyst 2010

Semi-Automated Chemistry Analyser (Manual)

- Photometric Accuracy.
- Dual function flow-cell and cuvettes.
- Mono and Bichromatic measurements.
- 99 tests channel available.
- Multi standard curves & user friendly, intuitive software.
- LCD Touchscreen display with integrated thermal printer.
- Low maintenance costs to promote lab efficiency.







Cat No. : MON0240S

Monarch 240+

Fully-Automated 240 Tests/Hour Bench Top Chemistry Analyser

- Fully automated, discrete, STAT priority. 240 tests per hour.
- Colorimetric and Turbidimetric.
- Up to 40 sample positions.
- 120 positions optical plastic cuvettes. Wavelength 340nm 800 nm.
- Low sample and reagent pipetting volumes increasing precision and accuracy of results.

Monarch 480

Fully-Automated 480 Tests/Hour Chemistry Analyser

- Fully automated, discrete, STAT priority.
- 400 tests per hour or 480 tests per hour with optional ISE mode.
- Colorimetric, Turbidimetric and ISE.
- Up to 115 sample positions and up to 112 double reagent disk positions.
- 120 positions optical plastic cuvettes.
- Wavelength 340nm 800 nm.
- Reaction Cuvette Cleaning with 7 stops and 11 step rinsing.
- Liquid level detection, clot detection and collision protection.



Monarch 680

Fully-Automated 680 Tests/Hour Chemistry Analyser

- Fully automated random access, STAT priority.
- 680 tests per hour or 1000 tests per hour with optional ISE mode.
- Colorimetric, Turbidimetric and ISE.
- Up to 115 sample positions and up to 112 double reagent disk positions.
- 120 positions optical plastic cuvettes.
- Wavelength 340nm 800 nm.
- Reaction Cuvette Cleaning with 7 stops and 11 step rinsing.
- Liquid level detection and collision protection.
- Large absorbance linear range up to 3.6Abs.
- Suitable for medium to large laboratories.



Monarch 1600

Ful	y-Automated 1600 Tests/Hour Chemistry Analyser		1		
	Fully automated random access, discrete, STAT priority. Constant speed 1200 tests/ hour for colorimetric, 400 tests/ hour with ISE mode. Long life halogen lamp. Wavelength 340nm - 800 nm. Analysis method: 1- point assay, 2-end point assay, 2-point rate assay, rate - A assay.	-			T
	 Independent sample probes, reagent probes. Automatic liquid level detection, probe liquid level tracing and clot detection function. Probe inner wall with pressure rinsing. 2 independent mixers. 160 new type reaction cuvettes. Temperature of reaction disk is 37C (+/- 0.1C). 140 sample positions, supporting multiple tubes and sample cups. 	34	.Or 40.		U ' #
	2 compartments, 132 reagent positions and 2 detergent positions with continuous refrigeraton.			Cat No : MON	1016005

Cat No. : MON01600S



Accessories

Description	Details	Cat No.	Size
Halogen Lamp	For Monarch 1600	MON00016	1
Halogen Lamp	For Monarch 240/480/680	MON00015	1
ISE Blood Control High	For Monarch Series	MON00014	5 x 4 ml; 5 x Vial Lyo
ISE Blood Control Low	For Monarch Series	MON00013	5 x 4 ml; 5 x Vial Lyo
ISE Blood Sample Calibrator	For Monarch Series	MON00005	5 x 4 ml Diluent; 5 x Vial Lyo
ISE Detergent	For Monarch Series	MON00004	1 x 100 ml
ISE Diluent	For Monarch Series	MON00001	1 x 2 L
ISE Internal Standard Solution	For Monarch Series	MON00003	1 x 2 L
ISE Module	For Monarch Series	MONISE00	1
ISE Module Chloride Electrodes	For Monarch Series	MON0008	1
ISE Module Potassium Electrode Body	For Monarch Series	MON00012	1
ISE Module Potassium Electrode Tip	For Monarch Series	MON00011	1
ISE Module Reference Electrodes	For Monarch Series	MON00010	1
ISE Module Sodium Electrodes	For Monarch Series	MON00009	1
ISE Reference Solution	For Monarch Series	MON0002	1 x 2 L
ISE Standard Solution High	For Monarch Series	MON00006	3 x 10 ml
ISE Standard Solution Low	For Monarch Series	MON00007	3 x 10 ml
Monarch 240 Accessories	Spare Parts Set for Year 1	MON240AC	1 Set
Monarch 480 Accessories	Spare Parts Set for Year 1	MON480AC	1 Set
Monarch 680 Accessories	Spare Parts Set for Year 1	MON680AC	1 Set
Monarch 1600 Accessories	Spare Parts Set for Year 1	MON1600AC	1 Set



Coagulation Analysers

Cascade +

Fully Automatic Coagulometer, Using a Turbodensitometric Detection Principle

- Automatically performs the analysis of various assays.
- Automatic rerun.
- Capacity of 32 samples. 32 reaction cuvettes.
- 6 positions for reagents vials.
- Innovative cleaning program to avoid cross infection. Samples can be run in batches, random or in profiles to promote laboratory flexibility.



Thrombo Read+

Semi-Automated Single Channel Coagulometer

- Measures all clotting assays. Light weight and compact to utilise lab space. User friendly software.
- Test menu includes analysis of various coagulation assays.

Cat No. : COAGA1CH



Thrombo Fast+

Semi-Automated Two Channel Coagulometer

- Two channel coagulometer which enables duplicate tests to be run. Compact and easy to use. Ideal for all laboratories.

- Suitable for a wide variety of clotting assays.

- Suitable for a Wide Variety of cloth 2 measuring channels. 4 reagent positions. Mixing by magnetic stirring bars. 18 cuvette positions. Internal printer.

- ChipCARD® technology.



Thrombo Speed+

Semi-Automated Four Channel Coagulometer

- Four channel coagulometer which enables duplicate tests to be run. Compact and easy to use. Ideal for all laboratories. Suitable for a wide variety of clotting assays. 2 measuring channels. 4 reagent positions

- 4 reagent positions.
- Mixing by magnetic stirring bars.
- 18 cuvette positions. Internal printer. ChipCARD® technology.

Cat No. : COAGA4CH

Accessories

Description	Details	Cat No.	Size
Coagulation Cuvettes Dispo System, Inc Mixer (With Card)	For Thrombo Read + / Fast + / Speed +	COAGCUVD	Box Containing 5 x 100 Cuvettes (with Card)
Coagulation Cuvettes, Inc Mixer (No Card)	For Thrombo Read + Only	COAGCUV1	1 Bags x 500 Cuvettes (No Card)
Coagulation Cuvettes, Inc Mixer (With Card)	For Thrombo Read + / Fast + / Speed +	COAGCUV3	Box Containing 5 Bags x 500 Cuvettes (with Card)
Thermopapier 57mm	For Thrombo Fast + / Thrombo Speed +	COAGROLL	10 Rolls per Carton





Electalyte-300

Electrolyte Analyser

- Built-in integrated reagent pack to reduce biological contamination.
- Auto calibration and slope-offset adjustement to ensure accurate result.
- Excellent measuring range and detection precision.
- User-friendly interface and online fault alarm with troubleshooting.
- One-button maintenance procedure to improve efficiency.
 Measuring cycle less than 25 seconds.
- Results storage > 10,000
- Standard network configuratiopn and online upgradable function.
- Barcode management and reagent volume alert.
- 7 inch colour touch screen.
- Built-in thermal printer.
- Upgradable autoloader.
- Air and liquid mixture flush to eliminate carryover.
- Smart sensor to diagnostic and adjust hardware.
- Competitive cost per test.

Accessories

Description	Details	Cat No.	Size
Accessories Pack	For Electalyte-300	ELECT-ACC	
Calabration Standard Solution	For Electalyte-300	ELECT-CAL	
Cleaning Solution	For Electalyte-300	ELECT-CLN	
Conditioning Solution	For Electalyte-300	ELECT-CON	
De-Proteinizer Kit	For Electalyte-300	ELECT-DEP	
Electalyte Reagent Pack	For Electalyte-200/500	ELECP100	650ml of Sol. A and 350ml of Sol. B
Electalyte Reagent Pack	For Electalyte-300	ELECP300	650ml of Sol. A and 350ml of Sol. B
Electalyte Reagent Pack	For Electalyte-400/600	ELECP400	650ml of Sol. A and 350ml of Sol. B
ISE Electrode	For Electalyte-300	ELECT-ISE	
Quality Control Kit	For Electalyte-300	ELECT-QC1	
Reference Electrode	For Electalyte-300	ELECT-DRE	
Reference Refill Solution	For Electalyte-300	ELECT-REF	
Refill Solution	For Electalyte-300	ELECT-ELC	
Replacement Electrode for Ca		ELECTDCA	
Replacement Electrode for Cl		ELECTDCL	
Replacement Electrode for K		ELECTDK1	
Replacement Electrode for Li		ELECTDLI	
Replacement Electrode for Na		ELECTDNA	
Replacement Electrode for pH		ELECTDPH	
Replacement Reference Electrode		ELECTDRE	

Cat No.: ELECT-300





FlowSense Reader

Rapid Test Lateral Flow Reader (Compatible with FlowSense Products)

The FlowSense Reader is designed to be an accesible and efficient solution for food quality testing, offering significant advantages to end users. It requires a minimal initial investment, making it financially accessible for a wide range of businesses and organizations. The product is user-friendly, requiring no prior experience or extensive technical knowledge, which allows anyone to operate it with ease.

- Easy user accessibility with minimal training needed.
- Portable for on-site testing.
- Provides semi-quantitative results to reduce visual interpretation errors.
- Multiple target nalysis capability.
- Export via USB drive.
- High data storage capacity.
- Easy printing facilities via USB.



Dry Bath Incubator

Dry Bath Incubator (Used in Conjunction with the FlowSense Product Range and Honey Quality Test Products)

- Instant temperature display, both heating and cooling.
- Multiple program running (9 programs can be set).
- Temperature deviation Calibration function.
- Built-in safety device for over-temperature.
- Intelligent algorithm.
- Automated recovery of power cut.
- Multiple modules options (Module DB9 Gluc/Fruc and DB2 FlowSense).
- LED display.
- Failure Detection and Buzzing alarm.

PRISM Food Enzymatic Analyser

Single-Beam Enzymatic Food Analyser

- Can achieve detections limits that meet EU/CODEX requirements for honey
- analysis.
- No expensive analyser required.
- Rapid result time.
- Reduce laboratory out-sourcing costs.
- Detection limits meet EU regulations.
- Minimal technical expertise required.
- Demonstrate the premium quality of your product.
 Strengthen in house control testing prior to blending.



Cat No. : BIOXBFD63





Microscan Mini Microplate Reader

8-Channel Microplate Reader (Including Software)

Its streamlined design and minimal footprint offers superb optics to fit any size lab bench. The Microscan Mini and your PC team up to provide rapid reading and full access to high level software, reporting, curve fitting, and data storage that meet the requirements of modern laboratories. Instrument and software come ready to "plug and play" with your PC.

- The Microscan Mini (including software) is a compact, 8-channel microplate reader.
 Streamlined design and minimal footprint offers superb optics to fit any size lab bench.
- The Microscan Mini and your PC team up to provide rapid reading and full access to high level software, reporting, curve fitting, and data storage that meet the requirements of modern laboratories.
- Instrument and software come ready to "plug and play" with your PC.



Accessories

Description	Details	Cat No.
FlowSense Adaptor	20 x 70 mm Cartridge	BIOXBFD63
FlowSense Adaptor	4.5mm	BIOXBFD51
PRISM Thermal Printer	Data Printer for the PRISM Analyser	BIOXBFD49
Square Quartz Curvettes	10mmm, 2 per Box	BIOXBFD50



Gel Card Centrifuge

Gel Card Centrifuge / Blood Type Card Centrifuge

The Fortress Gel Card Centrifuge specially designed with rotors to suit gel microcolumn blood type cards. It is the perfect choice in blood serology, blood type identification tests before crossmatching, erythrocyte washing, microcolumn agglutination, immunology, and other tests.

- Brushless motor to make sure free maintenance, no powder pollution, microcomputer control and digital display the time, speed, and RCF to let the operation easily.
 - Special centrifugation technical to different reagents according to the requirement.
- 10 Regular centrifugation programs.
- Electrical lid lock, over speed, and imbalance protection.



Gel Card Incubator

24 Card Capacity Gel Card Incubator

- The Fortress Gel Cards incubator offers a temperature range from Room Temperature and 5-60°C range with 1-10000 minutes range.
- It is presented for 24 cards capacity and heating timeof 10-25 minutes for 30°C – 37°C/60°C respectively with LCD display showing false detection and alarm function.





Gel Card Reader

Gel Card Reader (Suitable for ABO/Rh Grouping and Antibody Screening)

- Reads all varieties of gel type cards.
- Fully programmable, high-resolution image analysis. Detects and interprets "weak" positives reliably.
- Batch mode option for maximum throughput.
- Automatic bar-code reading of gel card barcodes.



Cat No. : READER01



Celldiff-3+

3 Part WBC Differential Haematology Analyser (Also Available in 5 Part)

- Rapid test turnaround time at 80 tests per hour.
- Results displayed in the form of histogram.
- Easy to use with intuitive software.
- Reduced consumables cost.
- 18-parameters hematology report.
- Sample type: Whole blood.
- Sampling time: 45 secs.
- 3-part WBC differential, for various patient profiles in 45 seconds with no preparation time.
- Designed to promote lab efficiently.
- 4 USB ports.

Cat No. : CELLDIF3



Bio-2+





CLIA-100 / CLIA-200

Fully Automated Chemiluminescence Immunoassay System

- Wash System: Patent for cleaning, arrayed circle magnestic design.
- Reaction Modules: Controls temperature controls, >4 reaction protocols.
- Reagents Loading: 20 kits/unit, up to 80 kits/4 units, on board fridge.
- System Integration Control Center: Touchscreen, LIS/HIS.
- Samples Loading: Capacity of 150/batch, continuous loading.
- Consumables Loading: Disposable tips with no contamination.

Coming Soon

Cat No. : CLIA-100/200

Innowash

Multi-channel Automatic Microplate Washer

- 96-way manifold, automatic and manual positioning.
- Processing time 55 seconds / microplate.
- Accurate and precise volume dispensing.
- Shaking function where time and speed are adjustable.
- Expansive memory capacity.
- Easy to use with intuitive software.

Microscan*



(*Washer Available on Request)

Four-Filter Microplate Reader


- Touch screen interface and intuitive software.
- Rapid test turnaround time.
- On-board printer.
- Single ELISA Plate read time > 13 secs.
- Bichromatic optics.
- Standard model includes four filters (six filters also available)






Seraqual 365: IQC Management Software

The Seraqual 365 IQC Management Software is a peer data QC programme designed for use with Seraqual Controls. It provides a cost effective and comprehensive package for labs who need to manage their daily QC activities on a centralised platform in real time.




- A dashboard interface that presents a condensed panel of the features of the system showing options most frequented. It also flags any exceptions for review
- A simple easy to use interface that allow labs to configure and enter results easily with a need for minimal training
- Interactive charts for quick and easy performance review including Histogram and Levey –Jennings

- range can be reviewed quickly
- Statistical Comparison Report can be specified by date range at the instrument, method or all method level.
- Peer Group Statistics allowing you to compare your results with other labs using the same QC material
- Advanced Statistical Analysis that includes performance indicators such as Uncertainty of Measurement (UM), Bias, Total Error (TE) and Sigma Scores.



Ordering Information



OC Tests

2

Instrument ID	Cobas				
Location	Chemistry Lab				
Instrument Name	Roche Cobas c311				
Control Type	Clinical Chemistry				
Lots	HSN029 - HSE029				
Show Current Assays		All Assays		Add	Assay
	Method	All Assays Reagent Supplier	Unit	Add Temperature (°C)	
 Current Assays 	Method Naphtlyl with Pentandiol		Unit U/L	_	Curren
Current Assays Analyte		Reagent Supplier		Temperature (°C)	Assay Current

This option allows a lab to quickly see the instruments registered, the programmes and lots associated with these instruments and then the associated QC test.

Here a lab can also set the Target mean and SD of a test (cumulative, user defined or Seraqual 365 (targets available for selected assayed lots)). 3

Result Entry

A simple feature to allow easy manual addition of results.

Instructions for Use (English Only)


OC Data Re

An alarm is activated if a result entered violates a multi rule set.

There are also indicators to show when a result has been edited, commented upon or reviewed.

Target Mean/SD	HSN029				
	Mean	0.0000	Mean	0.0000	
	SD	0.000	SD	0.000	
Your Statistics	Results	192	Results	192	
	Mean	14.9812	Mean	35.2177	
	SD	1.268	SD	2.643	
	CV%	8.46	CV%	7.51	
	Bias%	0.00	Bias%	0.00	

SECTION 15 | IQC & EQA Programmes



The Levey Jennings and Histograms interactive charts can be generated within a specific date range proving invaluable for identifying data shifts and trends.

The charts can be downloaded, saved, annotated and printed for the users convenience.



5

Exceptions Report

The Exceptions Report presents all results that have violated the multi rules set in a date range that can be specified.

Through colour coding it identifies the violated rules that are 'Alert' and those that are 'Reject'.

Any comments associated with a result are also detailed.

Cobas - Ch	emistry La	b - Roche Cobas	c311			
Acid Phos	phatase Pr	ostatic, Naphthyl	with Pentandio	I, Roche Diagnostic	s, U/L,37°C	
			н	SN029	1	HSE029
DATE	TIME	OPERATOR	RESULT	SDI	RESULT	SDI
06/06/2023	11:06:38	FD	15.4000 10	0:x+ 0.5583	36.4000	10:x+ 0.5382
Bilirubin D	irect, DCA,	3P Bioservice S.	R.L., µmol/L			
			н	SN029	I	HSE029
DATE	TIME	OPERATOR	RESULT	SDI	RESULT	SDI
07/06/2023	14:20:48	FD	8.2000 1:	28 2 0474	25.5000	1 8610

6

Statistics Comparison Report

A worldwide and country comparison of statistics at either the Instrument, Method or All Method level is available in this report.

A date range can be specified.

This report includes performance indicators such as Mean, SD, CV%, Bias %, SDI and CVI.

	stics Comparison						00	mparise	on Level: I	nstrume	n								
Blood Ga	s & Electrolytes, Chloride, Enzymat	lic, Roch	e Diagnos	tics, mn	nol/L														
LOT	INSTRUMENT	NO. YOU	OF RESULTS	WORLD	HOU	MEAN	WORLD	YOU	SD COUNTRY	WORLD	YOU	CVS.	WORLD	BIAST COUNTRY	WORLD	SOL	WORLD	CVI	WORL
105-01-005	Roche Cobas c311 - Chemistry Lab 2	109	109	109	14.105	14,104	14.104	0.63	0.63	0.63	4.42	4.42	4.42	NA	NA	NIA	NA	NA	NZ
108-02-008	Roche Cobas c311 - Chemistry Lab 2	109	109	109	72.623	72.620	72.620	1.15	1.15	1.15	1.58	1.50	1.58	NA	NA	NIA	NA	NA	NP
105-03-006	Roche Cobas c311 - Chemistry Lab 2	108	105	105	233.267	233,264	233,264	3.41	3.41	3.41	1.45	1.46	1.46	NA	NA	NA	NA	NA	N
Clinical C	hemistry, Acid Phosphatase Prost	atic, Nap	hthyl with	Pentan	diol, Roc	he Diagno	stics, U/I	., 37°C											
	hemistry, Acid Phosphatase Prost	NO	OF RESULTS			MEAN			80			CV%		BLAST		801		си	
Clinical C LOT HSE029				Pentan HORLD 243	4iol, Roc 100 35.218		works 35.230	-, 37°C	SD COUNTRY 2.54	WORLD 2.54	100 7.51	CV% COUNTRY 7.45	1058LD 7.45	BLAST COUNTRY -0.04	100MLD	SDI COUNTRY 0.00	WORLD 0.00	CVI COLANTRY 1.00	WORLD
LOT	INSTRUMENT	NO. YOU	OF RESULTS	WORLD	YOU	MEAN	WORLD	YOU	COUNTRY			COUNTRY		COUNTRY	WORLD	COUNTRY		COUNTRY	
LOT	INSTRUMENT Roche Cobas c311 - Chemistry Lab	NO. 192	OF RESULTS COLMTRY 243	1808LD 243	900 35.218	MEAN COUNTRY 35.230	WORLD 35.230	YOU 2.64	COUNTRY 2.54	2.64	7.51	COLN/RY 7.48	7.48	COUNTRY -0.04	WORLD -0.04	0.00	0.00	COUNTRY 1.00	1.00

Peer Group Statistics

A lab can select World or Country Statistics for a chosen Control Type and Lot.

The peer group can be at a number of different levels as seen in the image.

The report includes Mean, SD, CV% and the number of labs.

			Group				
O World		0	Analyte				
O United Ki	ngdom		Analyte/Meth	bd			
Control Type	Clinical Chemistry		Analyte/Meth	od/Instrumer	t		
			Analyte/Meth	od/Reagent S	upplier		
Lot	H5N055 Q		Analyte/Meth	od/Instrumer	t/Reagent	Supplier	
eer Group		SI Unit	Results	Mean	SD	CV%	No. Labs
ALT / GPT		U/L	90	27.3941	1.079	3.94	2
		g/L	87	17.0878	17.252	100.96	3
Albumin			100	146.7460	26.409	18.00	2

8

Advanced Statistics Report

This report draws on a number of important statistical metrics that include Mean, SD, CV%, Uncertainty of Measurement (UM), Bias, TE, Sigma (BV) and Sigma (CLIA).

The report can be based at either the Instrument, Method or All Method Level and if required within a specified date range.

9

QC Data Review

A lab selects a date range (instrument, lot and assay selection is optional) and the report generated displays any results that have not yet been reviewed.

Results violating a multirule are highlighted according to their 'Alert' or 'Reject' status.

A reviewer has the ability to set all the results to reviewed or individually look at results and comment.

This function allow quick and easy identification of tests that may be problematic.

Additional Features



Programme available in English, Spanish, Chinese, Farsi and Arabic.



Co-ordinators have the ability to look at the Statistics Comparison Report and Advanced Statistical Analysis Report of the labs they are responsible for.



Instruction for Use available on the Software



Automatic deactivations of peer data to maintain the integrity of the data

For further information please contact admin@seragual365.com

SEQAS: EQA Management Software


SEQAS is an EQA scheme that forms part of the Fortress Quality Control Platform. EQA helps to assure customers, such as physicians, patients, and health authorities, that the laboratory can produce reliable results. Individual laboratories can use EQA to identify problems in laboratory practices, allowing for appropriate corrective action. EQA participation will help to evaluate reliability of methods, materials, and equipment, and to evaluate and monitor training impact.



For laboratories performing public health-related testing, EQA can help to assure that results from different laboratories during surveillance activities are comparable. EQA participation is usually required for accreditation. Samples received for EQA testing, as well as the information shared by the EQA provider, are useful for conducting educational and professional development activities.

Participation in an external quality assessment program provides valuable data and information for the laboratory which:

- allows peer comparison of performance and results among different test sites;
- provides early warning for systematic problems associated with kits or operations;
- provides objective evidence of testing quality;
- indicates areas that need improvement;
- identifies training needs.



Description	Cycle	Cat No.	Size
SEQAS External Quality Assessment Programme		SEQAS00P	
SEQAS Monthly Clinical Chemistry (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS001	1 x 12 x 5ml
SEQAS Monthly Immunoassay Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS002	1 x 12 x 1ml
SEQAS Haematology Programme (Apr-Mar)	2 Cycles = 12 Months (1 Sample/Month)	SEQAS003	2 x 6 x 3ml
SEQAS Monthly HbA1c Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS004	1 x 12 x 0.5ml
SEQAS Monthly Coagulation Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS005	1 x 12 x 1ml
SEQAS Monthly Lipids Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS006	1 x 12 x 1ml
SEQAS Monthly Cardiac Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS007	1 x 12 x 2ml
SEQAS Monthly Blood Gas & Electrolytes Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS008	1 x 12 x 2ml
SEQAS Monthly Specific Protein Panel Programme (Jan-Dec)	1 Cycle = 12 Months (1 Sample/Month)	SEQAS009	1 x 12 x 1ml





Neonatal Screening

Neonatal screening is a critical component of public health and pediatric diagnostics, aimed at identifying congenital disorders and metabolic conditions in newborns before they manifest clinically. This early detection is vital for conditions such as phenylketonuria (PKU), cystic fibrosis, sickle cell disease, and congenital hypothyroidism, among others.

Timely identification of these disorders allows for prompt intervention and treatment, which can significantly improve health outcomes, prevent serious complications, and reduce mortality rates. For example, early treatment of PKU through dietary management can prevent severe intellectual disabilities, while timely intervention in congenital hypothyroidism can avert growth and developmental delays.

Fortress Diagnostics manufactures, develops and supplies an extensive range of high quality neonatal screening tests for the detection of inborn errors of metabolism (IEM).

Featured Products

PKU



A PKU screening test is a blood test given to newborns 24–72 hours after birth. PKU stands for phenylketonuria, a rare disorder that prevents the body from properly breaking down a substance called phenylalanine (Phe). Phe is part of proteins that are found in many foods and in an artificial sweetener called aspartame.

Glucose-6-Phosphate Dehydrogenase



Babies with glucose-6-phosphate dehydrogenase deficiency (G6PD) do not make the enzyme glucose-6-phosphate dehydrogenase (G6PD) properly. This enzyme protects the red blood cells from being destroyed prematurely by harmful substances in the body.

Total Galactose



The Total Galactose kit tests for Galactosemia, meaning "galactose in the blood". Babies with this metabolic condition are not able to metabolize a certain type of sugar (galactose) found primarily in breast milk, cow's milk, and dairy products. When galactose can't be broken down and digested, it builds up in the tissues and blood in large amounts.

Description	Details	Format	Cat No.	Size	Storage
Glucose-6-Phosphate Dehydrogenase (G6PDH)	Digitonin (UV)	Lyo	BXC0571A	R1: 2 x 50ml R2: 1 x 2ml R3:1 x 2ml R4: 1 x 20ml	2-8°C
Glucose-6-Phosphate Dehydrogenase (G6PDH)	UV Visual	Lyo	BXC0574C	100T	2-8°C
Neonatal TSH	ELISA	LS	BXE0682A	96T	2-8°C
Phenylketonuria (PKU) Screening	Enzymatic Colorimetric		BXC0810A	R1: 1x10ml; R2: 1x2ml; R3: 1 vial; STD: 1 Set	2-8°C
Total Galactose	Enzymatic Colorimetric	LS	BXC0812A	100T	2-8°C



SECTION 17 MOLECULAR DIAGNOSTICS

SECTION 17 | Molecular Diagnostics

Molecular Diagnostics

Molecular diagnostics is a cutting-edge field that utilises techniques such as DNA sequencing, polymerase chain reaction (PCR), and other nucleic acid-based methods to detect and analyze genetic material, enabling the diagnosis of various diseases at a molecular level. This approach allows for the identification of pathogens, genetic mutations, and biomarkers associated with diseases, including infectious diseases, cancers, and hereditary disorders.

Molecular diagnostics offers high sensitivity and specificity, leading to accurate and timely diagnoses that can guide personalized treatment strategies, improve patient outcomes, and facilitate targeted therapies. The importance of molecular diagnostics lies in its transformative impact on modern medicine. It provides clinicians with critical information about the underlying mechanisms of diseases, allowing for early detection and intervention, which is particularly crucial in cases of rapidly progressing conditions like infections or cancer. Additionally, molecular diagnostics can aid in the monitoring of treatment effectiveness and disease progression, enabling healthcare providers to make informed decisions about patient management. As technology advances, molecular diagnostics continues to expand its applications, contributing to personalised medicine, public health surveillance, and the development of novel therapeutic approaches, ultimately enhancing patient care and disease prevention efforts.

Featured Products

Fortress Universal Mini Kit (RNA Extraction)



The Fortress Universal Mini Kit is the ideal tool for isolation and purification of highly pure total nucleic acid, like human genomic DNA, bacterial DNA, viral DNA and RNA from 200µl of fresh or frozen plasma, serum, urine, cell free body fluids as well as rinsed liquid from swabs, breast milk, pretreated sputum, BAL and supernatant from stool suspension or whole blood (100 µl), and additionally only for veterinary applications from allantoic fluid or rinse liquid from cloacal or tracheal swabs and from organ abrasions.

COVID-19 RT-PCR

The Fortress Diagnostics COVID-19 Real Time Polymerase Chain Reaction (PCR) test is intended for the detection of SARs-Cov-2. The PCR test is ideal for laboratories with large throughput requirements with accurate results available in 24 hours.

Description	Details	Format	Cat No.	Size	Storage
COVID-19 RT-PCR	For Molecular Detection of COVID-19 by Real Time PCR		MC19-PCR1	96T	2-8°C
Fortress Universal Mini Kit	Nucleic Acid Extraction Kit		GRX00050	50T	2-8°C
Fortress Universal Mini Kit	Nucleic Acid Extraction Kit		GRX00100	100T	2-8°C
Fortress Universal Mini Kit	Nucleic Acid Extraction Kit		GRX00250	250T	2-8°C

SECTION 18 WESTERN BLOT ASSAYS

..........

..........

..........

......

.........

......

....

.......

............

..........

...........

............

Western Blot Assays

Western blot assays are a widely used laboratory technique for detecting and analysing specific proteins in a sample. The method involves separating proteins by gel electrophoresis, transferring them to a membrane, and using specific antibodies to identify the target protein. This assay is particularly valuable for confirming the presence and quantity of proteins of interest, as well as for analysing protein modifications, interactions, and expression levels. Western blotting is commonly employed in various fields, including biomedical research, clinical diagnostics, and the development of vaccines.

The importance of Western blot assays lies in their ability to provide reliable and specific information about protein expression and function. They are especially crucial in the diagnosis of certain diseases, such as HIV, where the presence of specific antibodies against viral proteins can confirm infection. Additionally, Western blotting is essential for validating results obtained from other techniques, such as ELISA, by providing a more detailed characterisation of proteins. The versatility and sensitivity of Western blot assays make them indispensable tools in research and diagnostics, contributing to a better understanding of disease mechanisms and aiding in the development of targeted therapies.

Key Features & Benefits

Western blot assays offers several benefits that make it a widely used technique in various fields, including medical diagnostics, research, and biotechnology:



Quantitative and Qualitative Analysis: Allows for the specific detection of

a particular protein within a complex mixture, using antibodies that bind to the target protein.



High Sensitivity: Western blotting is a technique which is highly sensitive and can detect low levels of protein.



High Throughput: It can provide both qualitative and quantitative data on protein expression levels.



Stability: It can be used for various types of samples, including cell lysates, tissue extracts, and purified proteins.

Description	Details	Format	Cat No.	Size	Storage
HCV Western Blot Assay			HCVWB018	18T	2-8°C
HIV Western Blot Assay			HIVWB018	18T	2-8°C
HTLV Western Blot Assay			HTLVWB18	18T	2-8°C



Stains

Staining is a fundamental technique in diagnostics that enhances the visibility of cellular components and structures in biological samples, facilitating their examination under a microscope. Different staining methods, such as Gram staining, hematoxylin and eosin (H&E) staining, and immunohistochemistry, allow for the differentiation of cell types, the identification of pathogens, and the observation of cellular morphology. Staining plays a crucial role in histopathology, microbiology, and cytology, providing valuable information for the diagnosis of various diseases, including infections, cancers, and inflammatory conditions.

The importance of staining in diagnostics lies in its ability to improve the accuracy and reliability of pathological assessments. By highlighting specific cellular features or antigens, staining helps pathologists and laboratory technicians make informed decisions regarding disease presence, type, and severity. For example, Gram staining differentiates between Gram-positive and Gram-negative bacteria, guiding appropriate antibiotic treatment. Immunohistochemical staining can identify specific markers in tumor tissues, assisting in cancer classification and treatment planning. Overall, staining is an indispensable tool in clinical diagnostics that enhances the interpretation of complex biological samples, ultimately contributing to better patient care and outcomes.

Featured Products

Geimsa Stain



Giemsa stain is a classical cytogenetic and histopathological stain used primarily in microbiology and hematology. It is a mixture of methylene blue, eosin, and Azure B, which stains cells differentially, allowing for the detailed visualization of cellular components. This stain is particularly useful for identifying parasites such as Plasmodium species, responsible for malaria, and for highlighting the chromosomal structures in cytogenetic studies, enabling the detection of genetic abnormalities. Its ability to differentiate between different cellular elements makes it invaluable for diagnosing various blood disorders and infections.

Gordon McLeod Reagent



Gordon McLeod reagent, also known as Gordon and Sweet's silver staining, is a specialised histological stain used primarily to visualise reticular fibers in tissue sections. This staining technique involves the reduction of silver salts to metallic silver, which deposits along the reticular fibers, rendering them visible under a microscope as black or dark brown structures. It is particularly useful in pathology for assessing the architecture of the extracellular matrix and detecting abnormalities in reticulin fiber networks, which can be indicative of various liver diseases, tumors, and other conditions affecting connective tissue.

May Grunwald



The May-Grünwald stain is a vital component of the May-Grünwald-Giemsa (MGG) staining technique, widely used in hematology and cytology for examining blood smears and bone marrow samples. This stain combines eosin and methylene blue, allowing for the differential staining of cellular components, which results in vivid coloration of blood cells and other cellular elements. It helps in the identification and classification of different types of white blood cells, red blood cell morphology, and the presence of parasites or other abnormal cells. The May-Grünwald stain's clarity and contrast make it essential for diagnosing various hematological disorders and infections.



Ordering Information

Description	Details	Format	Cat No.	Size	Storage
Eosin (2% ; Aqueous)	General purpose counter stain.	Stains	BX90125A	1 x 125ml	2-8°C
Eosin (2% ; Aqueous)	General purpose counter stain.	Stains	BX90250A	1 x 250ml	2-8°C
Eosin (2% ; Aqueous)	General purpose counter stain.	Stains	BX90500A	1 x 500ml	2-8°C
Geimsa Stain	For cytogenetics and histopathological diagnosis.	Stains	BX60250A	1 x 250ml	2-8°C
Geimsa Stain	For cytogenetics and histopathological diagnosis.	Stains	BX60500A	1 x 500ml	2-8°C
Geimsa Stain	For cytogenetics and histopathological diagnosis.	Stains	BX61000A	1 x 1000ml	2-8°C
Gordon McLeod Reagent (Oxidase Reagent)	An oxidase reagent to detect the presence of an enzyme cytochrome oxidase found in some bacteria.	Stains	BX80100A	1 x 100ml	2-8°C
Kovacs Reagent	For Indole Test, particularly useful in the identification of E.coli.	Stains	BX70100A	1 x 100ml	2-8°C
Lugol's Iodine	Mordant in gram staining and other Haematology staining.	Stains	BX10100A	1 x 100ml	2-8°C
May Grunwald	For staining of blood, bone marrow smears and clinical cytological specimens.	Stains	BXS0250A	1 x 250ml	2-8°C
May Grunwald	For staining of blood, bone marrow smears and clinical cytological specimens.	Stains	BXS0500A	1 x 500ml	2-8°C

For bulk orders of any stain not listed, please enquire at info@fortressdiagnostics.com

SECTION 20 ANTIBIOTIC DISCS/TABLETS

Antibiotic Discs/Tablets

Antibiotic discs/tablets, commonly used in the disc diffusion method (also known as the Kirby-Bauer test), are a vital tool in microbiology for assessing the susceptibility of bacteria to specific antibiotics. In this technique, paper discs impregnated with known concentrations of antibiotics are placed on an agar plate that has been inoculated with the target bacterial strain. After incubation, the plates are examined for zones of inhibition around the discs, indicating the effectiveness of the antibiotics against the bacteria. This method allows for the determination of bacterial resistance patterns and helps guide appropriate antibiotic therapy.

The importance of antibiotic discs in diagnostics cannot be overstated. They play a crucial role in clinical microbiology by providing rapid and reliable results that inform treatment decisions, especially in the face of rising antibiotic resistance. By identifying which antibiotics are effective against specific bacterial infections, healthcare providers can tailor treatments to individual patients, enhancing the chances of successful outcomes and reducing the risk of further resistance development. Furthermore, antibiotic disc testing contributes to public health surveillance by monitoring resistance trends in pathogens, guiding antibiotic stewardship programs, and informing guidelines for empirical therapy. Overall, antibiotic discs are essential for ensuring effective treatment of bacterial infections and improving patient care.

Featured Products

Ampicillin (AM)



Ampicillin (AM) antibiotic discs are used in microbiology to test the sensitivity of bacteria to the antibiotic ampicillin. Ampicillin is a broad-spectrum penicillin antibiotic that targets bacterial cell wall synthesis, making it effective against a wide range of Gram-positive and some Gram-negative bacteria. In a laboratory setting, the discs, which contain a standardised amount of ampicillin, are placed on agar plates inoculated with bacteria. The diameter of the zone of inhibition around the disc is measured to determine whether the bacteria are sensitive, intermediate, or resistant to ampicillin. This method is crucial for guiding appropriate antibiotic therapy.

Cefoxitin (FOX)

Cefoxitin (FOX) antibiotic discs are commonly used in laboratory testing for the identification of antimicrobial resistance, particularly for detecting Methicillin-Resistant Staphylococcus Aureus (MRSA) and other resistant strains of bacteria. Cefoxitin is a second-generation cephalosporin, and its disc diffusion test is highly sensitive for identifying the presence of resistance mechanisms, especially in beta-lactamase-producing organisms. The benefits of using cefoxitin discs include reliable and accurate detection of resistance, which is crucial for guiding appropriate antibiotic treatment. Additionally, cefoxitin provides a stable and efficient marker for testing in microbiology labs due to its broad spectrum and predictable results in resistance screening.

Neomycin (N)



Neomycin antibiotic discs are used in laboratory settings to test bacterial susceptibility to the antibiotic neomycin. Neomycin is an aminoglycoside antibiotic that inhibits protein synthesis in bacteria, making it effective against a wide range of Gram-negative and some Gram-positive bacteria. These discs contain a standardised concentration of neomycin, and when placed on an agar plate inoculated with bacteria, they create a zone of inhibition where bacterial growth is suppressed. The size of this zone indicates the bacteria's sensitivity or resistance to neomycin. The benefits of using neomycin antibiotic discs include their reliability in determining bacterial resistance, their usefulness in clinical diagnostics, and their role in guiding appropriate antibiotic treatments. Additionally, these discs help in monitoring antibiotic efficacy in research and clinical microbiology labs.

Description	Code	Format	Cat No.	Size	Storage
Amikacin	AN	Discs/ Tablets	BXABT01A	30 ug	Room Temp.
Amoxicillin-clavulanic	AMC	Discs/ Tablets	BXABT02A	30 (20 / 10) ug	Room Temp.
Ampicillin	AM	Discs/ Tablets	BXABT03A	10 ug	Room Temp.
Ampicillin-sulbactam	SAM	Discs/ Tablets	BXABT04A	20 (20 / 10) ug	Room Temp.
Azithromycin	AZM	Discs/ Tablets	BXABT05A	15 ug	Room Temp.
Azlocillin	AZL	Discs/ Tablets	BXABT06A	75 ug	Room Temp.
Aztreonam	ATM	Discs/ Tablets	BXABT07A	30 ug	Room Temp.
Carbenicillin	СВ	Discs/ Tablets	BXABT08A	100 ug	Room Temp.
Cefactor	CEC	Discs/ Tablets	BXABT09A	30 ug	Room Temp.
Cefadroxil	CFR	Discs/ Tablets	BXABT10A	30 ug	Room Temp.
Cefamandole	МА	Discs/ Tablets	BXABT49A	30 ug	Room Temp.
Cefazolin	CZ	Discs/ Tablets	BXABT50A	30 ug	Room Temp.
Cefepime	FEP	Discs/ Tablets	BXABT51A	30 ug	Room Temp.
Cefiderocol	FDC	Discs/ Tablets	BXABT52A	30 ug	Room Temp.
Cefixime	CFM	Discs/ Tablets	BXABT53A	5 ug	Room Temp.
Cefoperazone	CFP	Discs/ Tablets	BXABT54A	75 ug	Room Temp.
Cefotaxime	СТХ	Discs/ Tablets	BXABT55A	30 ug	Room Temp.
Cefotetan	CTT	Discs/ Tablets	BXABT56A	30 ug	Room Temp.
Cefoxitin	FOX	Discs/ Tablets	BXABT57A	30 ug	Room Temp.
Cefpodoxime	CPD	Discs/ Tablets	BXABT58A	10 ug	Room Temp.
Cefprozil	CPR	Discs/ Tablets	BXABT59A	30 ug	Room Temp.
Ceftaroline	СРТ	Discs/ Tablets	BXABT60A	30 ug	Room Temp.
Ceftazidime	CAZ	Discs/ Tablets	BXABT61A	10 ug	Room Temp.
Ceftazidime	CAZ	Discs/ Tablets	BXABT62A	30 ug	Room Temp.
Ceftazidime-avibactam	CZA	Discs/ Tablets	BXABT63A	50 (30 / 20) ug	Room Temp.
Ceftibuten	СТВ	Discs/ Tablets	BXABT64A	30 ug	Room Temp.
Ceftizoxime	ZOX	Discs/ Tablets	BXABT65A	30 ug	Room Temp.
Ceftobiprole	BPR	Discs/ Tablets	BXABT66A	5 ug	Room Temp.
Ceftolozane-tazobactam	СТ	Discs/ Tablets	BXABT67A	40 (30 /10) ug	Room Temp.
Ceftriaxone	CRO	Discs/ Tablets	BXABT68A	30 ug	Room Temp.
Cefuroxime	СХМ	Discs/ Tablets	BXABT69A	30 ug	Room Temp.
Cephalexin	CN	Discs/ Tablets	BXABT70A	30 ug	Room Temp.
Cephalothin	CF	Discs/ Tablets	BXABT71A	30 ug	Room Temp.
Chloramphenicol	С	Discs/ Tablets	BXABT72A	30 ug	Room Temp.
Cinoxacin	CIN	Discs/ Tablets	BXABT73A	100 ug	Room Temp.

SECTION 20 | Antibiotic Discs/Tablets

Description	Code	Format	Cat No.	Size	Storage
Ciprofloxacin	CIP	Discs/ Tablets	BXABT74A	5 ug	Room Temp.
Clarithromycin	CLR	Discs/ Tablets	BXABT75A	15 ug	Room Temp.
Clindamycin	СМ	Discs/ Tablets	BXABT76A	2 ug	Room Temp.
Colistin sulfate	CLR	Discs/ Tablets	BXABT78A	10 ug	Room Temp.
Doripenem	DOR	Discs/ Tablets	BXABT79A	10 ug	Room Temp.
Doxycycline	DO	Discs/ Tablets	BXABT80A	30 ug	Room Temp.
Eravacycline	ERV	Discs/ Tablets	BXABT81A	20 ug	Room Temp.
Ertapenem	ETP	Discs/ Tablets	BXABT82A	10 ug	Room Temp.
Erythromycin	E	Discs/ Tablets	BXABT83A	15 ug	Room Temp.
Fosfomycin (includes G-6-P)	FOS	Discs/ Tablets	BXABT84A	200 ug	Room Temp.
Fusidic acid	FA	Discs/ Tablets	BXABT85A	10 ug	Room Temp.
Gatifloxacin	GAT	Discs/ Tablets	BXABT86A	5 ug	Room Temp.
Gentamicin	GM	Discs/ Tablets	BXABT87A	10 ug	Room Temp.
Imipenem	IPM	Discs/ Tablets	BXABT88A	10 ug	Room Temp
Imipenem-relebactam	IMR	Discs/ Tablets	BXABT89A	35 (10 / 25) ug	Room Temp.
Kanamycin	К	Discs/ Tablets	BXABT90A	30 ug	Room Temp.
Lefamulin	LMU	Discs/ Tablets	BXABT91A	20 ug	Room Temp.
Levofloxacin	LVX	Discs/ Tablets	BXABT92A	5 ug	Room Temp.
Levonadifoxacin	LND	Discs/ Tablets	ВХАВТ93А	10 ug	Room Temp.
Linezolid	LZD	Discs/ Tablets	BXABT94A	30 ug	Room Temp.
Lomefloxacin	LOM	Discs/ Tablets	BXABT95A	10 ug	Room Temp.
Mesillinam	MEC	Discs/ Tablets	BXABT96A	10 ug	Room Temp.
Metopenem	MEM	Discs/ Tablets	BXABT45A	10 ug	Room Temp.
Metronidazole	MET	Discs/ Tablets	BXABT46A	5 ug	Room Temp.
Minocycline	MI	Discs/ Tablets	BXABT47A	30 ug	Room Temp.
Moxifloxacin	MXF	Discs/ Tablets	BXABT48A	5 ug	Room Temp.
Mupirocin	MUP	Discs/ Tablets	BXABT34A	200 ug	Room Temp.
Nafcillin	NF	Discs/ Tablets	BXABT35A	1 ug	Room Temp.
Nalidixic acid	NA	Discs/ Tablets	BXABT36A	30 ug	Room Temp.
Neomycin	Ν	Discs/ Tablets	BXABT37A	10 ug	Room Temp.
Netilmicin	NET	Discs/ Tablets	BXABT38A	10 ug	Room Temp.
Netilmicin	NET	Discs/ Tablets	ВХАВТ39А	30 ug	Room Temp.
Nitrofloxacin	NX	Discs/ Tablets	BXABT42A	10 ug	Room Temp.
Nitrofurantoin	FM	Discs/ Tablets	BXABT40A	300 ug	Room Temp.
Nitroxoline	NI	Discs/ Tablets	BXABT41A	30 ug	Room Temp.
Ofloxacin	OFL	Discs/ Tablets	BXABT43A	5 ug	Room Temp.

Antibiotic Discs/Tablets | SECTION 20

Description	Code	Format	Cat No.	Size	Storage
Omadacycline	OMC	Discs/ Tablets	BXABT44A	30 ug	Room Temp.
Oxacillin	OX	Discs/ Tablets	BXABT30A	1 ug	Room Temp.
Pefloxacin	PEF	Discs/ Tablets	BXABT31A	5 ug	Room Temp.
Penicillin G	Ρ	Discs/ Tablets	BXABT32A	10 units	Room Temp.
Piperacillin	PIP	Discs/ Tablets	ВХАВТЗЗА	100 ug	Room Temp.
Piperacillin-tazobactam	TZP	Discs/ Tablets	BXABT20A	110 (100 / 10) ug	Room Temp.
Polymyxin B	РВ	Discs/ Tablets	BXABT21A	300 units	Room Temp.
Quinupristin-dalfopristin	SYN	Discs/ Tablets	BXABT22A	15 ug	Room Temp.
Rifampicin	RA	Discs/ Tablets	BXABT23A	5 ug	Room Temp.
Spectinomycin	SPT	Discs/ Tablets	BXABT24A	100 ug	Room Temp.
Streptomycin	STS	Discs/ Tablets	BXABT25A	300 ug	Room Temp.
Sulfonamide	SSS	Discs/ Tablets	BXABT26A	300 ug	Room Temp.
Tedizolid	TZD	Discs/ Tablets	BXABT27A	2 ug	Room Temp.
Teicoplanin	TEC	Discs/ Tablets	BXABT28A	30 ug	Room Temp.
Telithromycin	TEL	Discs/ Tablets	BXABT29A	15 ug	Room Temp.
Temocillin	ТМО	Discs/ Tablets	BXABT13A	30 ug	Room Temp.
Tetracycline	TE	Discs/ Tablets	BXABT14A	30 ug	Room Temp.
Ticarcillin	TIC	Discs/ Tablets	BXABT15A	75 ug	Room Temp.
Ticarcillin-clavulanic acid	TIM	Discs/ Tablets	BXABT16A	85 (75 / 10)	Room Temp.
Tigecycline	TGC	Discs/ Tablets	BXABT17A	15 ug	Room Temp.
Tobramycin	ТМ	Discs/ Tablets	BXABT18A	10 ug	Room Temp.
Trimethoprim	ТМР	Discs/ Tablets	BXABT19A	5 ug	Room Temp.
Trimethoprim-sulfamethoxazole	SXT	Discs/ Tablets	BXABT11A	25 (1.25 / 23.75) ug	Room Temp.
Vancomycin	VA	Discs/ Tablets	BXABT12A	30 ug	Room Temp.

Terms & Conditions

1. General

All quotations made, all orders accepted, all goods sold, all service and advice rendered are subject to these conditions and to the exclusion of any previous conditions of purchase. Any variations to these must be expressly accepted by us in writing.

2. Specifications

We reserve the right to supply goods to the specification and/or design current at the date of dispatch. We cannot accept a request for goods as previously supplied as binding.

3. Admin & Handling Charge

An admin & handling charge of £25.00 will be added to all orders.

4. International Bank Charges

For international orders a \pm 15.00 international bank charge will be added to all BACS payments unless the customer agrees to pay these charges at their end.

5. PayPal/Credit Card Transactions

We accept PayPal, credit card and debit card payments. There is a 3.5% PayPal processing fee added to all customer orders who choose to pay via PayPal this will be detailed on your invoice. This is the same charge that Fortress incur to receive this method of payment.

6. Prices

Although we take great care to ensure price stability, we reserve the right to change prices without notice. Goods will be invoiced at the prices valid on the date of confirmation of the Buyer's order. All prices given are exclusive of VAT unless otherwise stated.

Fortress Diagnostics cannot accept responsibility for typographical errors in pricing within our catalogues, advertising literature or quotations.

Prices quoted are "Ex Works" (Incoterms 2010). Freight, insurance and all other respective costs e.g. Certificate of origin, legalised and notarised documents will be invoiced separately.

7. Delivery

Goods are shipped or dispatched at Buyer's risk. Our standard dispatch dates are 21 days after orders are confirmed and paid in full (unless you have credit terms).

Delivery times quoted are to be treated as an estimate only and Fortress Diagnostics shall not be liable for any failure to deliver within such times whether a time for delivery be quoted or not. We shall not be held responsible for delays caused by anything beyond our reasonable control. Any special delivery requirements must be notified to us in writing at the time of placing the order.

We shall have the right at our discretion to make a reasonable charge for special handling and/or delivery.

The Seller is entitled to make delivery in instalments unless such delivery would be unreasonable or unless otherwise agreed upon and confirmed by the Seller in writing. In the case that a fixed period for delivery is agreed upon and the Seller is in default with the supply, the Buyer shall grant the Seller a reasonable additional period of time, normally of 4 weeks.

8. Order Additions

Any additions will result in a new order being created in line with our standard delivery terms. Where possible we will attempt to dispatch your order additions with your original order but cannot guarantee this.

9. Damage/Loss/Shortage

No claim for breakages or missing goods can be recognised unless notified to Fortress Diagnostics and the carriers within 72 hours of receipt of the goods. If damages or shortages are immediately apparent upon delivery please:

1. Notify the carrier when signing for receipt of the goods.

2. Advise Fortress Diagnostics customer services team as soon as possible.

In the case of non-delivery of a complete consignment notification must be received by us within 10 days of invoice date.

10. Return of Goods

Notification of returns must be made within 1 week of receipt of goods. Goods which have been received damaged or in error must be held in the correct storage conditions.

Do not return the goods until you contact our customer services team for a returned goods authorisation number. Clearly mark the authorisation number on the package to ensure your goods can be speedily processed.

You will appreciate that we cannot accept responsibility for loss or damage to goods being returned to us unless we arranged collection. If the mistake was ours there is of course no charge to you. However, if the goods were incorrectly ordered an administration and re-stocking charge of 25% of the invoice value excluding VAT is chargeable. Temperature controlled products incorrectly ordered will not normally be accepted for return.

11. VAT

We are required to charge VAT on all UK orders at the current VAT rate.

For organisations that are VAT exempt each order must be accompanied by an exemption certificate unless special arrangements have been made in writing. Under circumstances where a VAT exemption certificate does not accompany an order we are obliged by law to charge VAT. If subsequently you reclaim a VAT credit it must be completed within 30 days of invoice date.

12. Payment Terms

Payments are to be made within the agreed payment terms set with our customers. We may at our discretion charge interest at the rate of 5% above the base lending rate of Barclays Bank plc on all overdue accounts to the date of actual payment. Payment in advance or financial guarantees may be requested from customers who do not have an account.

13. Quotations

Quotations are valid for 30 days.

14. Cancellation

Orders may not be cancelled without our prior written agreement and may be subject to a cancellation charge. Standing orders must be cancelled in writing at least 8 weeks prior to what becomes the last shipment date.

15. Liability

Fortress Diagnostics' liability for any loss or damage suffered by you and arising by way of defects in the goods or otherwise, howsoever shall be limited to the invoice price of the goods in respect of or in relation to which loss or damage is claimed. Our liability under this clause shall be in lieu of any warranty or condition implied by law as to the quality or fitness for any particular purpose of the goods and save as provided.

16. Ownership

All goods remain the property of Fortress Diagnostics Limited until payment is received in full. The responsibility for insurance and care of goods, including 3rd party risks lies with the purchaser.

17. Health & Safety

All products sold by Fortress Diagnostics Limited, should be handled by qualified personnel, exercising care during handling, usage and disposal. For further information please contact Fortress Diagnostics Limited.

Control of Substances Hazardous to Health 2002 (COSHH)

All Fortress products are developed and designed for 'IN VITRO' use only.

COSHH safety data sheets are available for all products on request.





Q	9A, The Technology Park, Antrim, United Kingdom, BT41 1QS
L	+44 (0)28 9448 7676
\leq	info@fortressdiagnostics.com
k	fortressdiagnostics.com





BR001FOROCT24

Copyright © 2024 Fortress Diagnostics Ltd. All rights reserved. VAT No.: GB 424802023. Product availability may vary from country to country and some products may be for Research Use Only (RUO). For more information on product application or availability, please contact us.